

Breeding Birds in the Wider Countryside: their conservation status 2008

Baillie, S., Marchant, J., Leech, D., Joys, A., Noble, D., Barimore, C., Grantham, M., Risely, K.
& Robinson, R..



Breeding Birds in the Wider Countryside: their conservation status 2008

Trends in numbers and breeding performance for UK birds

- [Summary of key findings](#)
- [Species list](#)
- [Using this web site](#)
- [Contents page](#)



Spotted Flycatchers, among the long-distance migrants that are of conservation concern, have decreased by 81% over the past 25 years: nest losses have increased greatly

Using this web site

This web site is a one-stop shop for information about the population status of our common terrestrial birds. It is based on data gathered by many thousands of volunteers who contribute to BTO-led surveys. With one web page per species, users can quickly find all the key information about trends in population size and breeding performance over the period 1967–2007, as measured by BTO monitoring schemes.

The [summary of key findings](#) provides a brief overview of our main findings this year. For each species, we provide:

- General information concerning species' conservation listings and UK population sizes
- A brief summary of observed changes in the size of the population and information concerning the possible causes of these changes
- A series of graphs and tables showing the trends and changes in population size and breeding performance over the past 39 years
- Trends calculated from BTO/JNCC/RSPB Breeding Bird Survey (BBS) data, not only for the UK as a whole but also for each of its constituent countries (England, Scotland, Wales and Northern Ireland)
- Alerts that highlight population declines in any census scheme of greater than 25% or greater than 50% that have occurred over the past 5 years, 10 years, 25 years and the maximum period available (usually 39 years).

Other pages provide details of the field and analytical [methods](#) that were used to produce the results for each species and of the methods used to identify [alerts](#). We [discuss](#) overall patterns of trends in abundance and breeding success, and compare the latest trend information and alerts with the Population Status of Birds list ([Gregory et al. 2002](#)). Four [appendices](#) list alerts and population changes by scheme, and there is also a facility to select and display your own [tables of population change](#). A detailed [references](#) section lists almost 300 of the most relevant recent publications, with onward links to abstracts or full text where available, and is a valuable key to recent scientific work by BTO and other researchers.

You can navigate your way around the site using links from the [contents page](#), from the [species index](#), and between sections. Alternatively, use the drop-down menus accessible from the menu bar at the top of each page. 'Species quick links', on the right-hand side of the menu bar, provides a drop-down list (in taxonomic order) with quick access to the species accounts.

The website covers the majority of British breeding birds, over 100 species in total, but excludes (with a few exceptions) colonial seabirds, which are well covered by the JNCC's [Seabird Monitoring Programme](#) ([Mavor et al. 2008](#)), and rare species that are included in the reports of the [Rare Breeding Birds Panel](#) (e.g. [Holling & RBBP 2007b, 2008](#)).

We value your comments on this report and particularly any suggestions on how it can be improved.

[Email your comments](#)

Authors

This report was written by Stephen Baillie, John Marchant, David Leech, Andrew Joys, David Noble, Carl Barimore, Mark Grantham, Kate Risely and Rob Robinson. The formal citation for the report is given in the page footer.

[Next page – Key findings](#)

Key findings

- Declining species
- New alerts
- Positive changes
- Reduced breeding success
- Increased breeding success
- Early breeding

Declining species

Best trend estimates over the longest available time period (usually 39 years) provide alerts to population declines of greater than 50% for 22 species.

These are **Grey Partridge**, **Little Grebe**, **Woodcock**, **Turtle Dove**, **Cuckoo**, **Lesser Spotted Woodpecker**, **Skylark**, **Tree Pipit**, **Yellow Wagtail**, **Song Thrush**, **Whitethroat**, **Willow Warbler**, **Spotted Flycatcher**, **Willow Tit**, **Marsh Tit**, **Starling**, **House Sparrow**, **Tree Sparrow**, **Linnet**, **Lesser Redpoll**, **Yellowhammer** and **Corn Bunting**. In addition, **Lapwing** has declined by 53% in the last 25 years, although its 39-year decline is measured at 34%.



Corn Buntings have declined by 87% over the last 39 years

The **Whitethroat** decline results from the severe crash between 1968 and 1969 linked to conditions on the wintering grounds. The **Little Grebe** decline should be treated with caution as we have long-term data from only a subset of its breeding habitats. Apart from these two, all these rapidly declining species are already red or amber listed on the Population Status of Birds (**PSoB**) list (**Gregory et al. 2002**).

For several of the species listed here long-term trend data are only available for England, where BTO has more volunteers to record information. Different long-term trends could be operating in other parts of the UK. **Lesser Redpoll**, **Tree Pipit** and **Woodcock**, in particular, have limited data.

A further ten species trigger alerts as a result of long-term declines of between 25% and 50% over periods of 22 to 39 years. These are **Common Sandpiper**, **Redshank**, **Little Owl**, **Meadow Pipit**, **Duncock**, **Mistle Thrush**, **Sedge Warbler**, **Reed Warbler**, **Lesser Whitethroat** and **Bullfinch**. Most of these species are already on the PSoB list on account of their population declines.

Recent alerts and alert changes



Yellow Wagtails have declined by 70% over the last 25 years

We draw special attention to the alerts for four species that have recently crossed the 50% decline threshold in the 25-year period. These are **Yellow Wagtail** (-70%), **Cuckoo** (-61%), **Willow Warbler** (-58%), and **Lapwing** (-53%). These species, all currently amber listed, may be candidates for addition to the red list at the next PSoB revision. Lapwing is an addition to this set of species this year; no species has fallen out.

We also identify three species that may be candidates to join the amber list (from green) owing to declines of between 25% and 50%. These are **Common Sandpiper** (-28% over 25 years), **Sedge Warbler** (-33% over 22 years) and **Lesser Whitethroat** (-59% over 22 years). **Sedge Warbler** is an addition to this set of species this year; one species, **Tawny Owl**, has fallen out because its decline no longer meets alert thresholds. **Little Owl** has also decreased strongly (-46% over 25 years) but is not eligible for amber listing because it is not native to the UK.

The alerts for **Sedge Warbler** and **Lesser Whitethroat** are raised by CES data, with census results showing shallower, non-significant declines. In addition, a CES decline of 33% in **Reed Warbler** may warrant investigation, although CBC, BBS and WBS results indicate that the species has increased.

Bullfinch is a currently red-listed species, but its long-term population decline is now marginally under the red-list threshold, at -50% over 39 years, due to a population increase of 7% over the past five

years. Unless the decrease resumes, this species may be a candidate for changing from the red to the amber list. Similarly the red-listed **Reed Bunting** now shows only a 19% decline over the last 39 years, with a significant upturn recorded over the past ten years. It is now questionable whether conservation listing is still warranted for this species.

Three currently amber-listed species, **Kestrel**, **Grey Wagtail** and **Goldcrest**, would no longer meet the population change criterion for amber listing, although **Kestrel** is also amber listed for reasons to do with its European status.

Positive changes

Only a few of those species that have declined previously show evidence of improvements in status, with eight formerly declining species showing clear positive trends over the last ten years. These are **Snipe**, **Grey Wagtail**, **Dunnock**, **Song Thrush**, **Whitethroat**, **Goldcrest**, **Tree Sparrow** and **Reed Bunting**.

The increases in the red-listed **Reed Bunting** and **Song Thrush** are particularly encouraging, as are the positive trends for the amber-listed **Grey Wagtail** and **Dunnock**. However, the most recent figures for **Song Thrush** and **Grey Wagtail** suggest that their recoveries may be levelling off well short of their previous population levels. Similarly while the BBS shows a 29% increase in **Snipe** over the last ten years, much of the former range across lowland Britain lost since the 1960s remains unoccupied, and moreover the population has been declining again since 2003.



Reed Buntings have begun to recover from their long-term decline

Seventeen species have more than doubled over the longest time period for which data are available (usually 39 years). These are **Mute Swan**, **Shelduck**, **Mallard**, **Goosander**, **Oystercatcher**, **Sparrowhawk**, **Buzzard**, **Stock Dove**, **Collared Dove**, **Woodpigeon**, **Green Woodpecker**, **Great Spotted Woodpecker**, **Nuthatch**, **Blackcap**, **Great Tit**, **Magpie** and **Carrion Crow**. **Canada Goose** has been lost to this group since the last report, WBS counts having levelled off, but BBS data indicate a continuing strong increase.

Reduced breeding success



Linnets have declined due to reduced breeding success

There are a number of species for which declines in breeding performance are likely to be driving the population declines (**Linnet** and **Lapwing**) or helping to inhibit recovery (possibly **Reed Bunting**). The importance of decreases in individual aspects of breeding performance for declining **Nightjar**, **Willow Warbler**, **Spotted Flycatcher**, **House Sparrow** and **Bullfinch** remain to be determined, as do the implications of the large reductions in CES productivity measures recorded for **Nightingale**, **Song Thrush**, **Sedge Warbler** and **Willow Warbler**. Many declining species show improving productivity, probably as a consequence of density-dependent processes (there are more resources available to feed the young when population numbers are low).

Increased breeding success

Increasing breeding performance may be helping to drive population expansion of a number of rapidly increasing species: the predatory **Grey Heron**, **Sparrowhawk**, **Buzzard** and **Barn Owl**; the corvids **Jackdaw**, **Magpie**, **Carrion Crow** and **Raven**; the resident seed-eaters **Collared Dove**, **Stock Dove** and **Greenfinch**; the resident insectivores **Great Spotted Woodpecker**, **Robin**, **Stonechat**, **Wren**, **Nuthatch**; and one migrant, **Redstart**.



Breeding success is improving for Redstarts

Early breeding

Data from the Nest Record Scheme provide strong evidence of shifts towards earlier laying in a range of



*On average, Grey Herons are now laying
28 days earlier than in 1968*

species, linked to climate change ([Crick et al. 1997](#), [Crick & Sparks 1999](#)). We have now identified 40 species that, on average, are laying up to 31 days earlier than they did 38 years earlier. The species involved represent a wide range of taxonomic and ecological groups, including raptors ([Kestrel](#) – 6 days), waterbirds ([Moorhen](#) – 5 days), waders ([Oystercatcher](#) – 7 days), owls ([Tawny Owl](#) – 7 days), migrant insectivores ([Willow Warbler](#) – 6 days), resident insectivores ([Blue Tit](#) – 7 days), corvids ([Magpie](#) – 31 days) and resident seed-eaters ([Chaffinch](#) – 8 days).

[Next page – Contents](#)

Contents

[Home](#)

[Key findings](#)

[Contents](#)

1. [General introduction](#)
 - 1.1 [The BTO's monitoring of breeding birds in the UK](#)
 - 1.2 [The value of combining results from different monitoring schemes](#)
 - 1.3 [The aims of this report](#)
2. [Methodology](#)
 - 2.1 [Breeding Bird Survey](#)
 - 2.2 [Common Birds Census](#)
 - 2.3 [Combined CBC/BBS trends](#)
 - 2.4 [Waterways Bird Survey](#)
 - 2.5 [Heronries Census](#)
 - 2.6 [Constant Effort Sites Scheme](#)
 - 2.7 [Nest Record Scheme](#)
 - 2.8 [The alert system](#)
 - 2.9 [Statistical methods used for alerts](#)
3. [Species index](#)
 - [Help on species accounts](#)
4. [Discussion](#)
 - 4.1 [The alert system](#)
 - 4.2 [Latest long-term alerts](#)
 - 4.3 [Ten-year trends and evidence of species recovery](#)
 - 4.4 [Increasing species](#)
 - 4.5 [Changes in breeding performance](#)
 - 4.6 [Conclusion](#)
5. [Acknowledgements](#)
6. [References](#)
7. [Appendix – Summary tables of changes in population size and breeding performance](#)
 - 7.1 [Tables of alerts and population increases from CBC/BBS](#)
 - 7.2 [Tables of alerts and population increases from WBS](#)
 - 7.3 [Tables of alerts and population increases from CES](#)
 - 7.4 [Tables of population declines and increases from BBS](#)
8. [Select your own table of population changes](#)

[Previous reports](#)

1. INTRODUCTION

Since its formation in 1933, BTO has been deeply committed to gathering quantitative information on the bird populations of the UK. Its nationwide network of skilled volunteers, many of whom are long-term contributors to survey schemes, provides the ideal way to monitor the bird populations that are widely distributed across the countryside. BTO data, from such schemes as the [BTO/JNCC/RSPB Breeding Bird Survey](#), the [Common Birds Census](#) and [Nest Record Scheme](#), have been increasingly influential in determining nature conservation policy in the UK. The partnership between JNCC and BTO has ensured that these schemes are operated and developed so as to provide high-quality information for nature conservation.

The value of the monitoring work undertaken by the BTO was recognised in the Government's Biodiversity Steering Group report ([Anon. 1995](#)). The BTO's results, particularly those regarding declining farmland species, are highlighted as an example of the way in which broad-scale surveillance techniques can identify important new trends. More generally, the report states that monitoring is essential if the broad aims, specific objectives and precise targets of the Government's Biodiversity Action Plans are to be achieved. It notes that:

- baselines must be established;
- regular and systematic recording must be made, to detect change; and
- the reasons for change should be studied, to inform action.

The BTO's monitoring schemes fulfil a considerable portion of these needs for a wide range of bird species in the UK.

The current system of [alerts](#) derived from the BTO's census and nest record data ensures that conservation bodies are quickly made aware of important demographic changes. Multi-species [indicators](#), making extensive use of BTO census data, track how bird populations are faring generally across the countryside, UK-wide and within specific regions or habitats. These indicators were developed in association with Government and some have been adopted by them as policy drivers. More recently, indicators have been developed on the European scale ([click here](#)).

1.1 [The BTO's monitoring of breeding birds in the UK](#)

1.2 [The value of combining results from different monitoring schemes](#)

1.3 [The aims of this report](#)

1.1 The BTO's monitoring of breeding birds in the UK


The Integrated Population Monitoring Programme has been developed by the BTO, in partnership with JNCC, to monitor the numbers, breeding performance and survival rates of a wide range of bird species. It has the following specific aims ([Baillie 1990, 1991](#)):

- (a) to establish thresholds that will be used to notify conservation bodies of requirements for further research or conservation action;
- (b) to identify the stage of the life cycle at which demographic changes are taking place;
- (c) to provide data that will assist in identifying the causes of such changes; and
- (d) to distinguish changes in population sizes or demographic rates induced by human activities from those that are due to natural fluctuations.

The programme brings together data from several long-running BTO schemes.

- Changes in numbers of breeding birds are measured by:
 - the [BTO/JNCC/RSPB Breeding Bird Survey \(BBS\)](#) – which began in 1994 and replaced the CBC (below) as the major monitoring scheme for landbirds, after a seven-year overlap. BBS is based on around 3000 1-km squares, within each of which birdwatchers count and record birds in a standardised manner along a 2-km transect. Because the survey squares are chosen randomly, the results are not biased towards particular habitats or regions. Combined CBC/BBS indices now provide long-running and ongoing population monitoring for many common birds.
 - the [Common Birds Census \(CBC\)](#) – which ran from 1962 to 2000. This scheme mapped the breeding territories of common birds on 200–300 mainly farmland and woodland plots each year, averaging about 70 and 20 ha respectively.
 - the [Waterways Bird Survey \(WBS\)](#) – which ran from 1974 to 2007. WBS observers mapped the territories of birds along rivers, streams and canals on 80–130 plots each year, each on average 4.5 km long. WBS has now been replaced by [WBBS](#), a transect scheme akin to BBS but with transects running alongside linear waterways, which started in 1998.
 - the [Constant Effort Sites Scheme \(CES\)](#) – which began in 1983 and is based on bird ringing at over 100 sites. The catching effort is kept constant at each site during each year, so that changes in numbers of birds caught will reflect population changes and not variation in catching effort.
 - the [Heronries Census](#) – through which counts of 'apparently occupied nests' have been collected from a high proportion of the UK's heronries every year since 1928.
- Changes in breeding performance are measured by:
 - the [Nest Record Scheme](#) – which began in 1939 and collates standardised information on up to 35,000 individual nesting attempts per year. This allows the measurement of:
 - laying dates
 - clutch sizes
 - brood sizes
 - nesting success during egg and chick stages.
 - the [CES](#) (see above) – which provides information on overall productivity for a range of species by measuring the ratio of juveniles to adults caught each year.
- Changes in survival are measured by:
 - the [British and Irish Ringing Scheme](#) – which provides information on the finding circumstances and longevity of ringed birds found dead by members of the public.
 - The [CES](#) can also provide information on survival rates, based on the recapture of ringed birds at CES sites. In future further information on survival rates will be provided through the [Retrapping Adults for Survival \(RAS\)](#) scheme.

The ways in which the schemes fit together are shown in the diagram below, which also demonstrates the way in which the BTO aims to combine all this information to understand the mechanisms behind changes in population sizes using population models.



Next section –1.2 The value of combining results from different monitoring schemes

[Back to Introduction Index](#)

1.2 The value of combining results from different monitoring schemes

Monitoring the changes in the size of a population does not in itself provide sufficient information on which to base an effective conservation strategy ([Goss-Custard 1993](#), [Furness & Greenwood 1993](#)). Concurrent monitoring of breeding performance and survival rates is necessary to allow changes in population size to be properly interpreted ([Temple & Wiens 1989](#), [Crick *et al.* 2003](#)) and, for long-lived species, can provide early warning of impending conservation problems ([Pienkowski 1991](#)).

Where good long-term data sets for breeding performance and survival are lacking, conservation action may have to be taken without an adequate understanding of the mechanisms involved or need to wait for detailed research to be undertaken. For many species, however, BTO already has the necessary data, collected by its volunteers over periods of several decades ([Greenwood 2000](#)).

For a long-lived species, a decline in population may not begin until a long period of low survival or reduced reproductive output has elapsed. The classic example is that of the [Peregrine](#), which in the UK suffered from poor breeding performance during the 1940s and 1950s due to sub-lethal DDT contamination. This decreased the capacity of the non-breeding population to buffer the severe mortality of breeding adults that occurred due to cyclodiene poisoning from the mid 1950s onwards ([Ratcliffe 1993](#)). Monitoring of breeding performance gave an early warning of subsequent numerical decline ([Pienkowski 1991](#)). Another example of a decline in breeding performance that presaged population decline is the catastrophic breeding failures of seabirds, particularly Arctic Terns, in Shetland ([Monaghan *et al.* 1989, 1992](#), [Walsh *et al.* 1995](#), [Mavor *et al.* 2003, 2004](#), [Wanless *et al.* 2005](#)).

Farmland birds

During the mid 1980s, the BTO identified rapid declines in the population sizes of several farmland bird species ([O'Connor & Shrubbs 1986](#), [Fuller *et al.* 1995](#)). The BTO has since been able to investigate the demographic mechanisms underlying these declines, using its long-term historical data sets ([Siriwardena *et al.* 1998a, 2000a](#)).

This investigation, which was Government-funded and undertaken jointly with Oxford University, looked at changes in population size, breeding performance and survival rates of a variety of species in relation to changing farming practice. It showed that species responded to different aspects of the agricultural environment, but that typically these aspects were linked to intensification or regional specialisation. Declines in survival rates were found to be the main factor driving population decline in these species, with the exception of [Linnet](#), for which the main factor appears to have been a decline in nesting success at the egg stage ([Siriwardena *et al.* 2000b](#)). The study was therefore able to eliminate some possible causes of change, and identify areas for future research, thus helping conservation bodies to use their scarce resources productively. This work made an important contribution to the wider programme of work on farmland birds undertaken by many research and conservation organisations ([Aebischer *et al.* 2000](#), [Vickery *et al.* 2004](#)).

This report describes a number of other cases where the combined analysis of BTO data sets has helped to identify the causes of population declines, for example on the pages for [Lapwing](#) ([Peach *et al.* 1994](#)), [Song Thrush](#) ([Baillie 1990](#), [Thomson *et al.* 1997](#), [Robinson *et al.* 2004](#)), [Sedge Warbler](#) ([Peach *et al.* 1991](#)), [Willow Warbler](#) ([Peach *et al.* 1995a](#)), [Spotted Flycatcher](#) ([Freeman & Crick 2003](#)), [Starling](#) ([Freeman *et al.* 2002, 2007b](#)), and [House Sparrow](#) ([Freeman & Crick 2002](#)). A fully integrated approach, estimating trends in numbers and demographic parameters through a single model containing data from various BTO surveys, is introduced by [Besbeas *et al.* \(2002\)](#).

Biodiversity Action Plans

The ability to quickly determine the stage of the life-cycle most heavily involved during population declines is particularly important for the conservation agencies when considering the plight of species on the lists of conservation concern ([JNCC 1996](#), [Anon. 1995, 1998](#)). Analysis of BTO data sets, which has already helped to build these lists, is a key point in several of the UK Government's [Biodiversity Action Plans](#) for rapidly declining species. Once conservation actions have been initiated, the BTO's Integrated Population Monitoring programme has a further function, because the success of these actions will be measured and assessed by continued BTO monitoring.

[Next section – 1.3 The aims of this report](#)

[Back to Introduction Index](#)

1.3 The aims of this report

This report is the latest in a series of reports that are used by conservation practitioners as a ready-reference guide to recent changes in status of breeding birds in the UK. By publishing it on the BTO website, we aim to make it available to a much wider audience, especially to BTO members and the general birdwatching public. We hope that it also provides a useful resource for schools, colleges and universities, the media, ecological consultants, decision-makers, local government, and the more general world of industry and commerce. In summary, its aims are:

- 1) To provide, to as wide a readership as possible, a species-by-species overview of the trends in breeding population and reproductive success of birds covered by BTO monitoring schemes since the 1960s, at the UK or UK-country scale.
- 2) To provide warning alerts to JNCC and Country Agencies and to other conservation bodies about worrying declines in population size or reproductive success, with special reference to species on the UK red and amber lists.

This document is the result of the sustained fieldwork of many thousands of the BTO's volunteer supporters. Without their enthusiasm for collecting these hard-won facts, the cause of conservation in the UK would be very much the poorer. The data we present here include information on distributions, from breeding-season and winter atlas projects, and on estimates of the absolute size of breeding populations, which are reported at intervals by the Avian Population Estimates Panel ([Stone *et al.* 1997](#), [Baker *et al.* 2006](#)). Colonial seabirds, which are well covered by the recently published results of Seabird 2000 ([Mitchell *et al.* 2004](#)) and by the JNCC's [Seabird Monitoring Programme \(Mavor *et al.* 2008\)](#), and the majority of species covered by the [Rare Breeding Birds Panel \(Holling & RBBP 2007b, 2008\)](#), are not included here. Wintering populations of waterfowl are covered by the Wetland Bird Survey annual reports (e.g. [Austin *et al.* 2008](#)) and by the WeBS alerts system ([Maclean & Austin 2008](#)).

The main emphasis of this report is on trends in the abundance and demography of individual species. The data on trends in abundance also provide the basis for multi-species [indicators](#) of bird population changes ([Gregory *et al.* 2004](#)). The [Wild Bird Indicator](#) has been adopted as one of the UK Government's 15 headline Quality of Life indicators. Furthermore, the related [Farmland Bird Indicator](#) is now being used as the basis of the Government's target for farmland bird recovery. This approach is now being extended more widely through a collaboration between EBCC, BirdLife and RSPB to produce [pan-European bird indicators](#).

The report is the latest in a series, begun in 1997, produced under the BTO's partnership with the Joint Nature Conservation Committee (on behalf of Natural England, Scottish Natural Heritage, the Countryside Council for Wales, and the Environment and Heritage Service in Northern Ireland) as part of its programme of research into nature conservation. Only the first two reports were published as paper reports, with subsequent ones being produced solely as web documents. A complete list of all the previous reports and links to those published online can be found [here](#).

[Section 2 – Methodology](#)

[Back to Introduction Index](#)

2. METHODOLOGY

Six monitoring schemes have contributed data to this report. Five provide data on changes in abundance: these are the **Breeding Bird Survey**, **Common Birds Census**, **Waterways Bird Survey**, **Heronries Census** and **Constant Effort Sites** ringing scheme. Two schemes, the **Nest Record Scheme** and Constant Effort Sites, provide data on changes in breeding productivity. In addition, information from detailed analyses of the recoveries of ringed birds, from the Ringing Scheme, is included where relevant.

The methodologies of the monitoring schemes are described below, including information on fieldwork, data preparation, sampling considerations and the statistical methods used in analysis.

- 2.1 [Breeding Bird Survey](#)
- 2.2 [Common Birds Census](#)
- 2.3 [Combined CBC/BBS trends](#)
- 2.4 [Waterways Bird Survey](#)
- 2.5 [Heronries Census](#)
- 2.6 [Constant Effort Sites Scheme](#)
- 2.7 [Nest Record Scheme](#)
- 2.8 [The alert system](#)
- 2.9 [Statistical methods used for alerts](#)

[Next page – Breeding Bird Survey](#)

[Back to Contents](#)

2.1 Breeding Bird Survey



The BTO/JNCC/RSPB Breeding Bird Survey (BBS) was launched in 1994, following two years of extensive pilot work and earlier desk-based studies. The introduction of the BBS was a move designed to overcome the limitations of the [Common Birds Census](#) (CBC), which had monitored bird populations since 1962. In particular, it improves the geographical representativeness of UK bird monitoring, thus boosting coverage of species and of habitats.

The BBS uses line transects rather than the more intensive territory-mapping method that was used by the CBC. This makes the survey relatively quick to undertake, and has been successful in encouraging a large number of volunteers to take part. The average time observers spend per visit is only around 90 minutes. Sampling units are the 1x1-km squares of the Ordnance Survey national grid, of which there are some 254,000 in the UK. From these we make random selections, by computer, for inclusion in the scheme (see [Square selection](#), below). The BBS requires a relatively large sample of survey squares, and the initial aim was to achieve coverage of about 2,500 squares.

An important aspect of BBS is its coordination through a network of volunteer BBS Regional Organisers. Information and survey forms are distributed first to these organisers, who contact volunteers willing to survey the squares every year. After the field season, forms are returned to BTO headquarters again via the Regional Organisers, but an alternative, on-line method for submission of BBS data was introduced in 2003 is already used by the majority of observers – see the [BBS pages](#) of the main BTO website for details.

Fieldwork involves three visits to each survey square each year. The first is to record details of habitat and to establish or re-check the survey route, while the second and third (termed 'early' and 'late') are to count birds. A survey route is composed of two roughly parallel lines, each 1 km in length, although for practical reasons routes typically deviate somewhat from the ideal. Each of these lines is divided into five sections, making a total of ten 200-m sections, and birds and habitats are recorded within these ten units. The two bird-count visits are made about four weeks apart (ideally in early May and early June), ensuring that late-arriving migrants are recorded. Volunteers record all the birds they see or hear as they walk along their transect routes. Birds are noted in three distance categories (within 25 m, 25–100 m, or more than 100 m on either side of the line, measured at right angles to the transect line), or as in flight. Recording birds within distance bands provides a measure of bird detectability in different habitats and thus allows population densities to be estimated more accurately. The total numbers of each species, excluding juveniles, are recorded in each 200-m transect section and distance category, as well as the timing of the survey and weather conditions.

By 1998, more than 2,300 BBS squares were being surveyed annually, close to the original target of 2,500. Only around a quarter of these plots were covered in 2001, owing to Foot & Mouth Disease access restrictions, but (thanks to our keen observers) the sample recovered immediately to over 2,100 in 2002 and had increased further to 2,254 squares in 2003, 2,526 in 2004, 2,879 in 2005 and 3,295 in 2006. The sample soared to 3,604 in 2007 ([Risely *et al.* 2008](#)). Squares are distributed throughout the UK and cover a broad range of habitats, including uplands and urban areas. There are around 100 species that are present on 40 or more BBS squares annually and so can be monitored with good precision at the UK scale ([Joys *et al.* 2003](#)), although a few present special difficulties because of their colonial or flocking habit or their wide-ranging behaviour. For most of these 100 species, BBS can also assess annual population changes within [England](#) alone, using data from 30 or more squares, and for about half the species also within [Scotland](#) and [Wales](#) as separate units. Sample sizes in [Northern Ireland](#) currently allow about 25 species to be indexed annually.

Square selection

Survey squares are chosen randomly using a stratified random sampling approach from within 83 sampling regions. These sampling regions, which in most cases are the standard BTO regions, are the 'strata' (literally layers) of the sample. Survey squares are chosen at random within each region, to a density that varies with the number of BTO members resident there. Regions with larger numbers of potential volunteers are thereby allotted a larger number of squares, enabling more birdwatchers to become involved in these areas. This does not introduce bias into the results because the analysis takes the differences in regional sampling density into account (see below).

Data analysis

Change measures between years are assessed using a log-linear model with Poisson error terms. For each species and square, counts are summed across all sections and distance bands for each visit ('early' and 'late') and the higher value is used in the model (or the single count if the square was visited only once). Counts are modelled as a function of square and year effects. Each observation is weighted by the number of 1-km squares in each region divided by the number of squares counted in that region, to correct for the differences in sampling density between regions. The upper and lower confidence limits of the changes indicate the certainty that can be attached to each change measure. When the limits are both positive or both negative, we can be 95% confident that a real change has taken place. Note that this presentation and its interpretation differs from the 85% confidence limits shown on most graphs within this report (see [section 2.8.4](#) for details).

Trends are presented as graphs in which annual population indices are shown in blue and their 95% confidence limits in green. A caveat, 'small sample', is provided against the trends for England, Northern Ireland, Wales and Scotland where the mean sample size is between 30 and 40 plots per year. A minimum sample size of 40 plots is required for the UK trends.

[Next section – 2.2 Common Birds Census](#)

[Back to Methodology Index](#)

[Click here](#) to go to the BBS section of the main BTO website

2.2 Common Birds Census

The Common Birds Census (CBC), which ran from 1962 to 2000, was the first of the BTO's schemes for monitoring population trends among widespread breeding birds, but has now been superseded for this purpose by [BBS](#).

The CBC was instigated to provide sound information on farmland bird populations in the face of rapid changes in agricultural practice. Although the original emphasis was on farmland, woodland plots were added by 1964. Fieldwork was carried out by a team of 250–300 volunteers. The same observers surveyed the same plots using the same methods year after year. On average, plots were censused for around seven consecutive years but a few dedicated observers surveyed the same sites for more than 30 years. Farmland plots averaged around 70 hectares in extent. Woodland plots were generally smaller, averaging just over 20 hectares. A small number of plots of other habitats, including heathlands and small wetlands, were also surveyed annually, especially before 1985.

A territory-mapping approach was used to estimate the number and positions of territories of each species present on each survey plot during the breeding season. Volunteers visited their survey plots typically eight to ten times between late March and early July and all contacts with birds, either by sight or sound, were plotted on outline maps at a standard scale of 1:2,500. Codes were used to note each bird's species, with sex and age where possible, and also to record activity such as song or nest-building. The registrations were then transferred to species maps and returned to BTO headquarters for analysis. The pattern of registrations on the species maps reveals the numbers of territories for each species. All assessments of territory number were made by trained BTO staff, applying rigorous guidelines, to ensure consistency between estimates across sites and years. Observers also provided maps and other details of the habitat on their plots. This makes it possible to match the distribution of bird territories with habitat features, providing the potential for detailed studies of bird–habitat relationships.

In 1990, the results from the Common Birds Census were brought together in the book *Population Trends in British Breeding Birds* ([Marchant *et al.* 1990](#)). This landmark publication discussed long-term population trends for the years 1962 to 1988 for 164 species, with CBC or WBS population graphs for around two-thirds of these.

The results from the Common Birds Census (CBC) provided reliable population trends for more than 60 of the UK's commoner breeding species and, through the linking of CBC with BBS, continue to be hugely influential in determining conservation priorities in the UK countryside. The store of detailed maps of almost a million birds' territories, collected through the CBC and maintained by BTO since the early 1960s, is a uniquely valuable resource for investigating the relationships between breeding birds and their environment, over wide temporal and spatial scales.

The weaknesses of the CBC as a monitor of UK bird populations were largely related to the time-consuming nature of both fieldwork and analysis. This inevitably limited the number of volunteers able to participate in the scheme, with the result that areas with few birdwatchers were under-represented. Constrained by the relatively small sample size, CBC concentrated on farmland and woodland habitats. Bird population trends in built-up areas and the uplands were therefore poorly represented. Furthermore, as the plots were chosen by the observers, some may not have been representative of the surrounding countryside and some bias towards bird-rich habitats might be suspected. It is for these reasons that the BBS was introduced in 1994. The two surveys were run in parallel for seven years to allow calibration between the results: for many species, CBC and BBS trends can be linked to form joint CBC/BBS trends that provide ongoing monitoring, continuous since the 1960s ([Freeman *et al.* 2003, 2007a](#); [section 2.3](#) of this report).

Validation studies

The CBC was the first national breeding bird monitoring scheme of its kind anywhere in the world and its value has been widely recognised internationally. The territory-mapping method adopted by the CBC is acknowledged as the most efficient and practical way of estimating breeding bird numbers in small areas, and has been well validated. Although intensive nest searches may sometimes reveal more birds, a comparison by [Snow \(1965\)](#) concluded that mapping censuses were a good measure of the true breeding population for 70% of species. Experiments to test differences between observers' abilities to detect birds found that, although there was considerable variation between individual abilities, the observers were consistent from year to year ([O'Connor & Marchant 1981](#)). As the CBC relies on data from plots covered by the same observer in consecutive years, this source of bias has no implications for the CBC's ability to identify population trends. It has also been confirmed that the sample of plots from which CBC results are drawn changed little in composition or character over the years ([Marchant *et al.* 1990](#)) and that the results of territory analysis are not affected by changes in analysts, once trained ([O'Connor & Marchant 1981](#)). [Fuller *et al.* \(1985\)](#) found that farmland CBC plots were representative of ITE lowland land-classes throughout England (excluding the extreme north and southwest), and closely reflected the agricultural statistics for southern and eastern Britain.

Data analysis

Population changes are modelled using a generalised additive model (GAM), a type of log–linear regression model that incorporates a smoothing function ([Fewster *et al.* 2000](#)). This replaces the Mountford model that employed a six-year moving window ([Mountford 1982, 1985, Peach & Baillie](#)

1994) and was used to produce annual population indices until 1999, but the principles are similar. These models are also very similar to log-linear poisson regression as implemented by program TRIM (Pannekoek & van Strien 1996). Counts are modelled as the product of site and year effects on the assumption that between-year changes are homogeneous across plots. Smoothing is used to remove short-term fluctuations (e.g. those caused by periods of severe weather or measurement error) and thus reveal the underlying pattern of population change. This is achieved by setting the degrees of freedom to about 0.3 times the number of years in the series. Confidence limits on the indices are estimated by bootstrapping (a resampling method; Manly 1991) and thus do not make any assumptions about the underlying distribution of counts.

Indices are plotted as the blue line on the graphs, and provide a relative measure of population size on an arithmetic scale relative to an arbitrary value of 100 in one of the years of the sequence. If an index value increases from 100 to 200, the population has doubled; if it declines from 100 to 50, it has halved. The two green lines on the graphs, above and below the index line, are the upper and lower 85% confidence limits. A narrow confidence interval indicates that the index series is estimated precisely, and a wider interval indicates that it is less precise. The use of 85% confidence limits allows relatively straightforward comparison of points along the modelled line: non-overlap of the 85% confidence limits is equivalent to a significant difference at approximately the 5% level (Anganuzzi 1993).

Caveats are provided to show where the data suffer from a 'Small sample' if the mean number of plots was less than 20. Data are flagged as 'Unrepresentative?' if the average abundance of a species in 10-km squares containing CBC plots was less than that in other 10-km squares of the species' distribution in the UK (as measured from 1988–91 *Breeding Atlas* data (Gibbons *et al.* 1993)), or, where average abundances could not be calculated, if expert opinion judged that CBC data may not be representative.

In practice most CBC data included in this report have been combined with BBS data to provide joint CBC/BBS trends, using the methods described in the next section. These methods for producing joint trends represent an extension of those described above.

[Next section – 2.3 Joint CBC/BBS trends](#)

[Back to Methodology Index](#)

[CLICK HERE](#) to go to the CBC section of the main BTO website

2.3 Combined Common Birds Census (CBC) and Breeding Bird Survey (BBS) trends

The field protocols for the two surveys are described in sections 2.1 and 2.2. As previously noted, the **CBC** has been an enormously influential project, providing the main source of information on national population levels in the UK since its inception in 1962. Because observers chose their own plots, however, its coverage had always been uneven. Coverage was predominantly in lowland, southeastern Britain, where the numbers of potential volunteers are greatest, while coverage was more patchy in more sparsely populated regions and especially the uplands ([Marchant *et al.* 1990](#)). CBC plots were situated in a limited number of habitats, predominantly farmland and woodland. Within a large rectangle of southeastern Britain (England and Wales south and east from Seascale, Scarborough and Exeter), the plots are nevertheless believed to be broadly representative, at least of lowland land-classes ([Fuller *et al.* 1985](#)). For species such as Wood Warbler and Meadow Pipit that have the greater part of their numbers in northern or western Britain, however, the CBC may not have accurately reflected national trends.

The **BBS**, on account of its more rigorous, stratified random sampling design, and its simplicity in the field, produces data that better cover the previously under-represented regions and habitats. In some **early editions** of 'Breeding Birds in the Wider Countryside' (e.g. [Baillie *et al.* 2002](#)), separate indices were published from CBC and BBS data, for those species with sufficiently large sample sizes. There being no new CBC data since 2000, however, it is futile to present a CBC-only trend – except for those few species where no joint or BBS index is available.

For most purposes, the presentation and analysis of longer time-series is required, dating back to before the establishment of the BBS but coming right up to the present day. The calculation of 25-year alert designations, as in this report, provides just one example. This need led to the BTO carrying out research into the compatibility of indices from BBS and CBC data in various years and regions, and the possibility of deriving trustworthy long-term indices from the two data sources in combination ([Freeman *et al.* 2003, 2007a](#)). This research suggested that for the vast majority of species considered there was no significant difference between population trends, calculated from the two surveys, based on that part of the country where CBC data are sufficient to support a meaningful comparison. Where a statistically significant difference was found, this was sometimes for very abundant species for which the power to detect even a biologically insubstantial difference was considerable. Within this region, therefore, long-term trends based on CBC and BBS data can be produced for almost all species previously monitored by the CBC alone. For ([Freeman *et al.* 2003, 2007a](#)) this was the area covered by [Fuller *et al.* \(1985\)](#), because CBC plots in that region were shown to be representative of lowland farmland there. As this region covers the bulk of England, and for consistency with the rest of this report, we have produced joint indices for CBC/BBS for the whole of England (the CBC/BBS England index), rather than just the English part of the 'Fuller rectangle'.

A second question then is whether one can obtain reliable trends over the same period for the entire UK. That is, since prior to 1994 only CBC data are available, are the population trends within the region well covered by the CBC typical of those for the UK as a whole? The shortage of CBC data in the north and west means that the only way of investigating this is via the BBS data. Significant differences in trends between the area well covered by the CBC and the rest of the UK were found for approximately half the species (see [Freeman *et al.* 2003, 2007a](#), for full details). For such species, a regional bias in CBC data means that no reliable UK index can be produced prior to 1994. In summary, joint population indices dating back to the start of the CBC can continue to be produced for that part of the country well served by the CBC (essentially England) for almost all common species. However, a similar UK index can be produced for only about 50% of species (CBC/BBS UK index).

This report presents joint CBC/BBS trends for the UK and/or England, as appropriate. Ideally the trends would have been estimated using generalised additive models ([Fewster *et al.* 2000](#)) but these were too computationally intensive, given the large number of sites involved. Therefore we fitted a generalised linear model, with counts assumed to follow a Poisson distribution, and a logarithmic link function, to the combined CBC/BBS data. Standard errors were calculated via a bootstrapping procedure and there is therefore no need to model overdispersion, as it does not affect the parameter estimates. BBS squares were weighted by the number of 1-km squares in each sampling region divided by the number of squares counted in that region as in standard BBS trend analyses. CBC plots were assigned the average weight of all BBS squares as this allows them to be incorporated within the analysis while retaining the convention of not applying weights within the BBS sample. The population trend was smoothed using a thin-plate smoothing spline with degrees of freedom about one third the number of years. Confidence intervals were calculated via a bootstrap procedure. Bootstrap samples were generated by resampling sites from the original data set, with replacement. A generalised linear model was then fitted to each bootstrap replicate and a smoothing spline fitted to the annual population indices as described above. Confidence limits were then calculated as the appropriate percentiles from the sets of smoothed estimates. The overall result is a smoothed trend that is mathematically equivalent to that produced from a generalised additive model. The method of estimation is less statistically efficient because the smoothing is not incorporated within the estimation procedure, and is likely to have resulted in more conservative statistical tests and wider confidence limits. However this compromise was necessary to make it possible to fit the trends within a reasonable amount of computer time (still several weeks).

Indices are plotted as the blue line on the graphs, and provide a relative measure of population size on

an arithmetic scale relative to an arbitrary value of 100 in one of the recent years of the sequence. If an index value increases from 100 to 200, the population has doubled; if it declines from 100 to 50, it has halved. Note that positive and negative percentage changes are not directly equivalent: for example, a decrease of 20% would require an increase of 25% to restore the population to its former level. The two green lines on the graphs, above and below the index line, are the upper and lower 85% confidence limits. A narrow confidence interval indicates that the index series is estimated precisely, and a wider interval indicates that it is less precise. The use of 85% confidence limits allows relatively straightforward comparison of points along the modelled line: non-overlap of the 85% confidence limits is equivalent to a significant difference at approximately the 5% level ([Anganuzzi 1993](#)).

[Next section – 2.4 Waterways Bird Survey](#)

[Back to Methodology Index](#)

2.4 Waterways Bird Survey

The Waterways Bird Survey (WBS) has monitored the population trends of up to 24 riparian bird species on canals and rivers throughout the UK since 1974. WBS uses a territory-mapping method like that of its parent scheme, the [Common Birds Census](#), to estimate the breeding population of waterbirds on each plot. Detailed territory maps are prepared that can be compared with habitat data to show which features of linear waterways are important to breeding birds. The plots average 4.4 km in length; almost half are slow-flowing lowland rivers with the rest either fast-flowing rivers/streams or canals. In recent years there have been around 90 plots distributed throughout the UK. The proportion of plots in the north and west of England is higher than existed in the CBC ([Marchant *et al.* 1990](#)). As with CBC, coverage outside England has been relatively poor.

All fieldwork has been carried out by volunteers. Observers are asked to survey their plots on nine occasions between March and July, mapping all the birds seen or heard onto 1:10,000-scale maps. Registrations are then transferred to species maps, which are analysed to reveal the numbers and positions of territories for each species. For the first 20 years all territory analysis was performed by trained headquarters staff but, during 1994–2007, observers completed their own territory analysis, based on the scheme's written guidelines, with results checked by BTO staff. As WBS has employed very similar methods to those of CBC, the validation studies carried out for the latter generally hold true for WBS (see [section 2.2](#)). [Marchant *et al.* \(1990\)](#) found that there had been little change in the composition of the WBS sample in terms of waterway type or geographical spread.

Population changes along waterways have been reported in *BTO News* for around 20 riparian species, of which Goosander is not covered by BBS monitoring. For specialist waterbirds, including [Little Grebe](#), [Mute Swan](#), [Common Sandpiper](#), [Kingfisher](#), [Sand Martin](#), [Grey Wagtail](#), [Dipper](#) and [Reed Warbler](#), targeted surveys along waterways can provide a better precision of monitoring than is possible through the more generalised BBS surveys. WBS indices can also add a new perspective on trends in waterbirds that are monitored, largely in different habitats, by CBC/BBS. For [Lapwing](#), populations declined rapidly on arable farmland during the late 1980s while numbers on WBS plots, typically representing populations along river floodplains, were more stable. [Yellow Wagtails](#) have declined much more steeply in WBS habitats than elsewhere.

WBS has similar limitations as a monitoring scheme that led to the CBC's replacement by BBS. In particular, plot distribution is biased geographically and possibly also towards sites that are good for birds, and an intensive survey method is used that severely limits the sample size ([Marchant *et al.* 1990](#)). A drawback specific to WBS is that it has covered only waterbirds. BTO has addressed these issues by setting up the [Waterways Breeding Bird Survey \(WBBS\)](#), which has been running since 1998 in parallel with WBS. WBBS uses BBS-style transect methods along random waterways, and includes all species of birds. WBS closed after the 2007 season and it is now expected that WBBS will become an ongoing scheme, providing useful monitoring data to supplement BBS.

Data analysis

Smoothed population trends are estimated using generalised additive models, with confidence intervals calculated by bootstrapping ([Fewster *et al.* 2000](#)). Trend analysis and presentation follows the same pattern as CBC ([section 2.2](#)), except that the 'Unrepresentative?' caveat has not been used. A caveat of 'Small samples' is provided when the number of plots falls between 10 and 20.

[Next section – 2.5 Heronries Census](#)

[Back to Methodology Index](#)

[CLICK HERE](#) to go to the WBS section of the main BTO website

2.5 Heronries Census

The BTO Heronries Census began in 1928 and is the longest-running breeding-season bird monitoring scheme in the world. As predators at the top of the freshwater food chain, Grey Herons are excellent indicators of environmental health in the countryside. They build large stick nests, mostly in colonies at traditional sites. The aim of this census is to collect annual nest counts of Grey Herons from as many sites as possible in the United Kingdom. Volunteer observers make counts of 'apparently occupied nests' at heron colonies each year. Changes in the numbers of nests, especially over periods of several years, provide a clear measure of the population trend. In recent seasons, observers have also counted the nests of [Little Egrets](#) *Egretta garzetta*, which have been appearing in an increasing number of southern heronries since the first breeding records in 1996. Counts of Cormorant colonies, which often occur alongside heronries, are also welcome ([Newson et al. 2007](#)).

Coverage is coordinated through a network of regional organisers. A core of birdwatchers and ringers monitor their local colonies annually, providing a backbone of regular counts. Around two-thirds of the heronries in England and Wales are currently counted each year, with more-complete censuses carried out in 1929, 1954, 1964, 1985 and 2003. Historically rather few counts have been made of heronries in Scotland and Northern Ireland, except during the special surveys, but support for the Heronries Census has been growing fast in recent years. Counts are submitted mostly on cards and the data are entered onto computer at BTO headquarters. The number of heronries counted each year has grown in recent years to around 550–600.

Data analysis

Population changes are estimated using a ratio-estimators approach derived from that of [Thomas \(1993\)](#). Essentially, the ratios of the populations in any two (not necessarily consecutive) years of the survey are estimated from counts at sites visited in each of those years. These ratios can be used to estimate the counts at sites that were not visited, and hence build an estimate of the total population. Further modifications have been made to allow for the extinction of colonies and the establishment of new ones ([Marchant et al. 2004](#)).

On the [Grey Heron](#) page of this report, the UK trend is presented graphically with annual estimates in blue and their 85% confidence limits in green. A smooth trend line in red is based on a non-parametric regression model, using thin-plate smoothing splines with 24 degrees of freedom. Trends are also shown for England and Wales together, and for England, Wales and Scotland alone.

[Next section – 2.6 Constant Effort Sites Scheme](#)

[Back to Methodology Index](#)

[CLICK HERE](#) to visit the Heronries Census page of the main BTO website

[CLICK HERE](#) to visit the Little Egret page of the main BTO website (WeBS)

2.6 Constant Effort Sites Scheme

The Constant Effort Sites (CES) Scheme uses changes in catch sizes across a network of standardised mist-netting sites to monitor changes in the abundance and breeding success of common passerines in scrub and wetland habitats. At each constant effort site, licensed ringers erect a series of mist nets in the same positions, for the same amount of time, during 12 visits evenly spaced between May and August. Year-to-year changes in the number of adults caught provide a measure of changing population size, while the ratio of young birds to adults in the total catch is used to monitor annual productivity (breeding success). By monitoring the abundance of young birds between May and August, the CES method should integrate contributions to annual productivity from the entire nesting season, including second and third broods for multi-brooded species, but will also include a small component of mortality during the immediate post-fledging period. Between-year recaptures of ringed birds can also be used to calculate annual survival rates of adult birds, although this requires specialised analytical techniques (e.g. [Peach 1993](#)) and is not considered further here. Further details of the CES Scheme are presented by [Peach *et al.* \(1996\)](#) and methods of analysis are detailed in [Peach *et al.* \(1998\)](#) for abundance measures and [Robinson *et al.* \(2007\)](#) for productivity measures.

The CES Scheme began in 1983 with 46 sites and now has around 120. The distribution of CES sites tends to reflect the distribution of ringers within Britain and Ireland. The majority are operated in England, and there are small numbers in Scotland, Wales, Northern Ireland and the Republic of Ireland. The CES routinely monitors the populations of 25 species of passerines in scrub and wetland habitats.

Data analysis

Smoothed trends in the abundance of adults and young are separately assessed using a generalised additive model (GAM), with 85% confidence intervals calculated by bootstrapping ([Fewster *et al.* 2000](#)). At sites where catching effort in a year falls below the required 12 visits, but eight or more visits have been completed, annual catch sizes are corrected according to experience during years with complete coverage, by incorporating an offset into the GAM (see [Peach *et al.* \(1998\)](#) for full details). Sites with fewer than eight visits in a given year are omitted for the year in question. Annual indices of productivity (young per adult) are estimated from logistic regression models applied to the proportions of juvenile birds in the catch, the year-effects then being transformed to measures of productivity relative to an arbitrary value of 100 in the most recent year. As above, catch sizes are corrected where small numbers of visits have been missed. It should be noted that these indices are relative, and are not estimates of the actual numbers of young produced per adult ([Robinson *et al.* 2007](#)).

Data are presented graphically with the smoothed trend in blue and their 85% confidence limits in green. A caveat is provided for 'Small samples' when the average number of plots per year is between 10 and 20.

[Next section – 2.7 Nest Record Scheme](#)

[Back to Methodology Index](#)

[CLICK HERE](#) to go to the CES section of the main BTO website

2.7 Nest Record Scheme

The BTO's Nest Record Scheme is the largest, longest-running and most highly computerised of such schemes in the world and possesses the most advanced and efficient techniques of data gathering, data capture and analysis ([Crick *et al.* 2003](#)). There are now more than a million nest records held by the Trust, of which 35% are computerised.

The primary aim of the Nest Record Scheme is to monitor the breeding performance of a wide range of UK birds annually as a key part of the BTO's data collection. Annual reports are published in *BTO News* (e.g. [Leech & Barimore 2008](#)) and the significant results communicated immediately to JNCC. Another primary aim is to undertake detailed analyses of breeding performance of species of conservation interest (e.g. [Crick *et al.* 1994](#), [Brown *et al.* 1995](#), [Peach *et al.* 1995a](#), [Crick 1997](#), [Chamberlain & Crick 1999](#), [Siriwardena *et al.* 2001](#), [Crick *et al.* 2002](#), [Chamberlain & Crick 2003](#), [Freeman & Crick 2003](#), [Browne *et al.* 2005](#), [Tryjanowski *et al.* 2006](#)).

The Nest Record Scheme gathers data on the breeding performance of birds in the UK through a network of volunteer ornithologists. Each observer is given a code of conduct that emphasises the responsibility of recorders towards the safety of the birds they record and explains their legal responsibilities. These observers complete standard nest record cards for each nest they find, giving details of nest site, habitat, contents of the nest at each visit and evidence for success or failure. When received by the BTO staff, the cards are checked, sorted and filed away ready for input and analysis. Those for Schedule 1 species are kept confidential. (These are species protected from disturbance at the nest by Schedule 1 of the [Wildlife & Countryside Act 1981](#): they are generally rare species and the location of their nests may need to be protected from egg collecting (an illegal activity) or other potential disturbance. To visit the nests of these species a special licence is required.) Computer programs developed by BTO check the data for errors and calculate first-egg date, clutch size, nest loss rates at egg and chick stages. Data are computerised according to priorities for population monitoring and for specific research projects.

Currently the BTO collects a total of more than 30,000 records each year for around 180 species. Typically, there are more than 150 records for 55 species and more than 100 for a further 10–15 species. The quality of records improved substantially in 1990 with the introduction of a new recording card, which promotes greater standardisation and clarity in the information recorded by observers. The general distribution of Nest Record Cards is patchy at the county scale but is more even over larger regions of the UK. Overall, Northern Ireland and parts of Scotland (southeast, Western Isles) and parts of England (West Midlands, southwest) have relatively low coverage, often reflecting observer density. A major analysis of trends over time in various aspects of breeding performance found relatively few differences between major regions in the UK, when analysed using analysis of covariance ([Crick *et al.* 1993](#)). Habitat coverage is broad, as the scheme receives records from all the UK's major habitats. Most records come from woodland, farmland and freshwater sites, but the scheme also receives data from scrub, grassland, heathland and coastal areas.

Data analysis

Five different variables were analysed for this report: laying date (where day 1 = January 1); clutch size; brood size; and daily nest failure rates during egg and nestling stages, calculated using the methods of [Mayfield \(1961, 1975\)](#) and [Johnson \(1979\)](#) (see [Crick *et al.* 2003](#) for a review).

To minimise the incidence of errors and inaccurately recorded nests, a set of rejection criteria was applied to the data: laying date included only cases where precision was within ± 5 days; clutch size was not estimated for nests which had been visited only once, for nests which were visited when laying could still have been in progress, or for nests which were visited only after hatching; and maximum brood size was calculated only for nests which were observed after hatching. The last variable is an underestimate of brood size at hatching, because observers may miss early losses of individual chicks; it differs from clutch size because eggs may be lost during incubation and hatching success may be incomplete.

Daily failure rates of whole nests were calculated using a formulation of [Mayfield's \(1961, 1975\)](#) method as a logit-linear model with a binomial error term, in which success or failure over a given number of days (as a binary variable) was modelled, with the number of days over which the nest was exposed during the egg and nestling periods as the binomial denominator ([Crawley 1993](#), [Etheridge *et al.* 1997](#), [Aebischer 1999](#)). Number of exposure days during the egg and nestling periods was calculated as the midpoint between the maximum and minimum possible, given the timing of nest visits recorded on each Nest Record Card (note that exposure days refer only to the time span for which data were recorded for each nest and do not represent the full length of the egg or nestling periods). Each calculation assumes that failure rates were constant during the period considered. Violations of this assumption of the Mayfield method can lead to biased estimates if sampling of nests is uneven over the course of each period. It is unlikely that any such bias would vary from year to year, so although absolute failure rates may be biased, annual comparisons should be unaffected ([Crick *et al.* 2003](#)). In this report, therefore, we present only temporal trends in daily nest failure rates.

Statistical analyses of nest record data were undertaken using SAS programs ([SAS 1990](#)). Regressions through annual mean laying dates, clutch sizes and brood sizes were weighted by sample size. Nest survival was analysed by logistic regression. Quadratic regressions were used when the inclusion of a quadratic term provided a significant improvement over linear regression. These are

described as 'curvilinear' in the tables on species pages. Significant linear trends are described as 'linear'. The best-fitting regressions (i.e. quadratic or linear) are presented on the figures in this report. Where neither regression is significant the linear regression line is shown for illustrative purposes.

Results are presented only if the mean sample size of records for a particular variable and species exceeds ten per year, and are presented with a caveat for small sample sizes if the mean number of records contributing data was between ten and 30 per year.

[Next section – 2.8 The alert system](#)

[Back to Methodology Index](#)

[CLICK HERE](#) to go to the NRS section of the main BTO website

2.8 The Alert System

- 2.8.1 [General approach](#)
- 2.8.2 [Smoothing population trends](#)
- 2.8.3 [Years used for analysis](#)
- 2.8.4 [Confidence limits and statistical testing](#)
- 2.8.5 [Data-deficient species](#)

2.8.1 General approach

The alert system used within this report is designed to draw attention to developing population declines that may be of conservation concern, and is described in detail by [Baillie & Rehfisch \(2006\)](#). It also identifies situations where long-term declines have reversed, leading to an improvement in conservation status. It must be stressed that the changes reported here are advisory and do not supersede the agreed UK conservation listings ([Gregory *et al.* 2002](#); see [PSoB](#) pages). They are based on similar criteria to *The Population Status of Birds in the UK*, however, and so provide an indication of likely changes at future revisions.

The system is based on statistical analyses of the population trend data for individual species. Alerts seek to identify rapid declines (>50%) and moderate declines (>25% but <50%). These declines are measured over a number of time-scales, depending on the availability of data – the full length of the available time series, and the most recent 25 years, 10 years and 5 years for which change can be estimated. The conservation emphasis is particularly on the longer periods, but short-term changes help to separate declines that are continuing – or accelerating – from those that have ceased or reversed.

The alerts are calculated annually using standard automated procedures. Where species are at the margin of two categories (e.g. a decline of about 25%) they may fire alerts in some years but not others, or different levels of alert in different years.

Data on some species might be biased, owing to possibly unrepresentative monitoring, or imprecise, owing to small sample sizes. Because these data often provide the only information that is available, our general approach is to report all the alerts raised but to flag up clearly any deficiencies in the data.

2.8.2 Smoothing population trends

Bird populations show long-term changes that do not follow simple mathematical trajectories. In addition to the long-term trends, unsmoothed population indices also show short-term fluctuations resulting from a combination of natural population variability and statistical error. We use smoothing techniques that aim to extract the long-term pattern of population change, without forcing it to follow any particular shape (such as a straight line or a polynomial curve). These methods remove most of the effects of short-term fluctuations (including any natural year-to-year variability) so that the long-term trend is revealed more clearly.

[Technical details available here](#)

2.8.3 Years used for analysis

Once a smoothed population trend has been calculated, change measures are calculated from the ratio of the smoothed population indices for the two years of interest. Population indices for the first and last years of a smoothed time series are less reliable than the others, and so we always drop them before calculating alerts. Because the latest year is not included, the alerts are therefore less up-to-date than they could be, but fewer false alarms are generated. The latest year's data points do contribute to the smoothed curve and are dropped only after the smoothing has taken place.

The time it takes BTO to collate and analyse bird monitoring data is another factor affecting the years that can be included in these analyses. Full analyses of data sets are not usually all available until 12–15 months after the end of a particular breeding season. Thus for a report prepared for year x (e.g. 2008) we have analyses of monitoring data up to year $(x-1)$ (e.g. 2007). As we drop the final year of the smoothed time series, we report here on change measures up to year $(x-2)$ (e.g. 2006).

Long-term changes for most of the species included in this report are calculated from joint Common Birds Census and Breeding Bird Survey data (CBC/BBS indices). The CBC started on farmland in 1962 and on woodland in 1964. However, the early years of the CBC population indices are strongly influenced by the effects of the unusually severe winters of 1961/62 and 1962/63, as well as by developments in methodology ([Marchant *et al.* 1990](#)). Joint CBC/BBS indices have been calculated using only the data from 1966 onwards, therefore, and population changes are calculated back to 1967.

2.8.4 Confidence limits and statistical testing

We show 90% confidence limits for population change measures wherever possible. Any decline where the confidence limits do not overlap zero (no change) is regarded as statistically significant and will trigger an alert if it is of sufficient magnitude. Note that, because we are seeking to detect only declines, we are using a one-tailed test – with a P value of 0.05. These confidence limits therefore do

not indicate whether increases are statistically significant.

The graphs of population trends show 85% confidence limits because these allow an approximate visual test of whether the difference between the index values for any two given years is statistically significant: if the index values for two given years are assumed to be independent, and normally distributed with standard errors of comparable size (standard errors differing by a factor of up to about 2 are quite acceptable), then to a good approximation the difference between them is significant at the 5% level if there is no overlap in their 85% confidence intervals (Buckland *et al.* 1992, Anganuzzi 1993). This test is fairly robust, and the independence assumption is reasonable if the years are well separated.

[Technical details available here](#)

2.8.5 Data-deficient species

There is uncertainty about the reliability of the results for some species, either because data may be unrepresentative or because they are based on a very small sample of plots. In these cases the cause of the uncertainty is recorded in the comment column of the population change table.

Unrepresentative data

In this report we present joint UK or England CBC/BBS trends only if there was no substantial or statistical difference between the trends from the two schemes over the period when they ran in parallel. Thus, since BBS results are drawn from a random sample, the trends are always considered to be representative of the region concerned.

In previous reports representativeness was assessed using the criteria developed by Gibbons *et al.* (1993). Data from the 1988–91 Breeding Atlas were used to compare the average abundance of a given species in 10-km squares with and without CBC plots. If average abundance is higher in squares without CBC plots, it is likely that much of the population is not well sampled by the CBC. In past reports, CBC data for such species were labelled as "unrepresentative". Where there are insufficient data to undertake such calculations, expert opinion was used instead.

Sample size

Sample size is assessed from the average number of plots contributing to the population indices for a given species in each year. A plot with a zero count would be included provided that the species had been recorded there in at least one year and that records for that plot were available for at least two years. Plots where a species has never been recorded do not enter the index calculations. These average sample sizes are shown in column four (plots) of the population change tables. For CBC, WBS and CES, a mean of between 10 and 19 plots is flagged as a small sample. For BBS indices for individual countries a mean in the range 30–39 plots is flagged as a small sample. UK BBS indices are presented only where samples reach at least 40 plots.

[Technical details available here](#)

[Next section – 2.9 Statistical methods used for alerts](#)

[Back to Methodology Index](#)

2.9 Statistical methods used for alerts

The [Alert System](#) page contains a general overview of how the alert system works. More detailed information is given below about the statistical methods used to estimate population indices, population changes and their confidence intervals.

2.9.1 General structure of data and models

2.9.2 Fitting smoothed models

2.9.3 CBC/BBS trends

2.9.4 Waterways Bird Survey

2.9.5 Constant Effort Sites Scheme

2.9.6 Heronries Census

2.9.1 General structure of data

The data for all of the schemes reported here consist of annual counts made over a period of years at a series of sites. They can thus be summarised as a data matrix of sites x years, within which a proportion of the cells contain missing values because not all of the sites are covered every year. Such data can be represented as a simple model:

$$\log(\text{count}) = \text{site effect} + \text{year effect}$$

Each site has a single site-effect parameter. These site parameters are not usually of biological interest but they are important because abundance is likely to differ between sites. The main parameters of interest are the year effects. These can be modelled either with as many parameters as years (an annual model), or with a smaller number of parameters, representing a smoothed curve.

A simple annual model would be fitted as a generalised linear model with Poisson errors and a log link function. This is the main model provided by the program TRIM ([Pannekoek & van Strien 1996](#)), which is widely used for population monitoring.

2.9.2 Fitting smoothed models

Our preferred method for generating a smoothed population trend is to fit a smoothed curve to the data directly using a generalised additive model (GAM) ([Hastie & Tibshirani 1990](#), [Fewster et al. 2000](#)). Thus the model from the previous section becomes:

$$\log(\text{count}) = \text{site effect} + \text{smooth}(\text{year})$$

where smooth(year) represents some smoothing function of year. It was not straightforward to fit GAMs to the CBC/BBS or Heronries Census data and we have therefore fitted smooth curves with a similar degree of smoothing to the annual indices (details below).

The non-parametric smooth curve fitted in our models is based on a smoothing spline. The degree of smoothing is specified by the number of degrees of freedom (df). A simple linear trend has $df = 1$, whereas the full annual model has $df = t - 1$, where t is the number of years in the time series. Here we set df to be approximately 0.3 times the number of years in the time series ([Fewster et al. 2000](#)). The degrees of freedom used for the main data sets presented in this report are summarised below.

	Years	Length of time series	df for smoothed index
CBC/BBS	1966–2007	43	13
Waterways Bird Survey	1974–2007	35	11
Constant Effort Sites	1983–2007	26	8
Heronries Census	1928–2007	81	24

Note that the numbers of years shown here are different from those available for calculating change measures, because we use the whole time series available for analysis (i.e. prior to the truncation of end points), and because we count the number of years in the time series rather than the number of annual change measures.

2.9.3 CBC/BBS trends

The model fitted to the combined CBC and BBS data is that historically employed for the BBS, a Generalised Linear Model with counts assumed to follow a Poisson distribution and a logarithmic link function. Standard errors were calculated via a bootstrapping procedure. For presentation in the figures, both the population trend and its confidence limits were also subsequently smoothed using a thin-plate smoothing spline. The overall result is a smoothed trend that is mathematically equivalent to that produced from a generalised additive model.

A similar method as employed for the joint CBC/BBS trend has been used for the BBS alone. This adopted a Generalised Linear Model with counts assumed to follow a Poisson distribution and a logarithmic link function. Standard errors were calculated via a bootstrapping procedure involving 199 bootstraps. For presentation in the figures, both the population trend and its confidence limits were also subsequently smoothed using a thin-plate smoothing spline.

2.9.4 Waterways Bird Survey

GAMs were fitted to the WBS data using the approach described above ([Fewster *et al.* 2000](#)). Confidence limits were fitted using a bootstrap technique to avoid restrictive assumptions about the distribution of the data. Bootstrap samples were drawn from the data by sampling plots with replacement. We generated 199 bootstrap samples from each data set and fitted a GAM to each of them. Confidence limits for the smoothed population indices (85% ci) and change measures (90% ci) were determined by taking the appropriate percentiles from the distributions of the bootstrap estimates. The section on [confidence limits and statistical testing \(2.8.4\)](#) gives the reasons for choosing these particular confidence limits.

The GAMs were fitted using a modified version of the FORTRAN program GAIM ([Hastie & Tibshirani 1990](#)).

2.9.5 Constant Effort Sites

GAMs were fitted to the CES data for catches of adults and juveniles separately with the addition of an offset to correct for missing visits. Confidence limits were fitted using a bootstrap technique to avoid restrictive assumptions about the distribution of the data. Bootstrap samples were drawn from the data by sampling plots with replacement. We generated 199 bootstrap samples from each data set and fitted a GAM to each of them. Confidence limits for the smoothed population indices (85% ci) and change measures (90% ci) were determined by taking the appropriate percentiles from the distributions of the bootstrap estimates, in a similar manner to that employed for the Waterways Bird Survey.

2.9.6 Heronries Census

The Heronries Census data were analysed using a modified sites x years model based on ratio estimation which incorporates information about new colonies (sites) that have been established and other colonies from the sample that are known to have gone extinct. The method was developed by [Thomas \(1993\)](#) specifically in relation to the heronries data set. Since then the heronries database has been substantially upgraded and the method has been applied to the full data set ([Marchant *et al.* 2004](#)).

The above method of analysis cannot be easily applied within a GAM framework. Therefore we fitted a smooth curve to the annual indices. This was done using PROC TSPLINE of SAS. This procedure should give very similar estimates to a GAM analysis but it does not provide confidence intervals for the smoothed population trend or the change measures derived from it. This is not a serious limitation as there are no potential alerts for [Grey Heron](#), whose populations have generally been increasing.

[Section 3 – Species pages](#)

[Back to Methodology Index](#)

SPECIES LIST

Jump to

[Wildfowl](#)
[Gamebirds](#)
[Waterbirds](#)
[Raptors](#)
[Waders](#)
[Pigeons](#)
[Owls](#)
[Larks](#)

[Thrushes](#)
[Warblers](#)
[Tits](#)
[Crows](#)
[Sparrows](#)
[Finches](#)
[Buntings](#)

List of species (in BOU taxonomic order)

WILDFOWL

[Mute Swan](#)
[Greylag Goose](#)
[Canada Goose](#)
[Shelduck](#)
[Mallard](#)
[Tufted Duck](#)
[Goosander](#)

GAMEBIRDS

[Red Grouse](#)
[Red-legged Partridge](#)
[Grey Partridge](#)
[Pheasant](#)

WATERBIRDS

[Red-throated Diver](#)
[Little Grebe](#)
[Great Crested Grebe](#)
[Cormorant](#)
[Grey Heron](#)

RAPTORS

[Hen Harrier](#)
[Sparrowhawk](#)
[Buzzard](#)
[Kestrel](#)
[Merlin](#)
[Hobby](#)
[Peregrine](#)
[Moorhen](#)
[Coot](#)

WADERS

[Oystercatcher](#)
[Ringed Plover](#)
[Golden Plover](#)
[Lapwing](#)
[Snipe](#)
[Woodcock](#)
[Curlew](#)
[Common Sandpiper](#)
[Redshank](#)

PIGEONS

[Feral Pigeon](#)
[Stock Dove](#)
[Woodpigeon](#)
[Collared Dove](#)
[Turtle Dove](#)
[Ring-necked Parakeet](#)
[Cuckoo](#)

OWLS

[Barn Owl](#)
[Little Owl](#)
[Tawny Owl](#)
[Nightjar](#)

[Swift](#)
[Kingfisher](#)

[Green Woodpecker](#)
[Great Spotted Woodpecker](#)
[Lesser Spotted Woodpecker](#)

LARKS

[Woodlark](#)
[Skylark](#)

[Yellow Wagtail](#)
[Grey Wagtail](#)
[Pied Wagtail](#)

[Dipper](#)

[Wren](#)

[Dunnock](#)

THRUSHES

[Robin](#)

[Nightingale](#)

[Redstart](#)

[Whinchat](#)

[Stonechat](#)

[Wheatear](#)

[Ring Ouzel](#)

[Blackbird](#)

[Song Thrush](#)

[Mistle Thrush](#)

WARBLERS

[Cetti's Warbler](#)

[Grasshopper Warbler](#)

[Sedge Warbler](#)

[Reed Warbler](#)

[Blackcap](#)

[Garden Warbler](#)

[Lesser Whitethroat](#)

[Whitethroat](#)

[Wood Warbler](#)

[Chiffchaff](#)

[Willow Warbler](#)

[Goldcrest](#)

[Spotted Flycatcher](#)

[Pied Flycatcher](#)

TITS

[Long-tailed Tit](#)

[Blue Tit](#)

[Great Tit](#)

[Coal Tit](#)

[Willow Tit](#)

[Marsh Tit](#)

[Nuthatch](#)

[Treecreeper](#)

CROWS

[Jay](#)

[Magpie](#)

[Jackdaw](#)

[Rook](#)

[Carrion Crow](#)

[Hooded Crow](#)

[Raven](#)

[Starling](#)

SPARROWS

[House Sparrow](#)

[Tree Sparrow](#)

FINCHES

[Chaffinch](#)

[Greenfinch](#)

[Goldfinch](#)

[Siskin](#)

[Linnet](#)

[Lesser Redpoll](#)

Sand Martin
Swallow
House Martin
Tree Pipit
Meadow Pipit

Bullfinch
BUNTINGS
Yellowhammer
Reed Bunting
Corn Bunting

Information to aid interpretation of the pages for individual species
can be found on the [Species Help Page](#)

MUTE SWAN *Cygnus olor*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: amber (>20% of European breeding population)

Long-term trend

UK, England: rapid increase

UK population size

28,000–30,000 adults in 1990 (Delany *et al.* 1992: APEP06); 23,900–25,600 pairs in 2000 (updated using CBC/BBS trend: BIE04); 28,600–35,200 birds in Britain in 2002 (Rowell & Spray 2004)

Status summary

Mute Swan populations, which had been fairly stable since the 1960s, have increased progressively since the mid 1980s, perhaps reflecting warmer winter weather and the replacement of anglers' lead weights, which had earlier caused many casualties, with non-toxic alternatives (Rowell & Spray 2004, Ward *et al.* 2007). WBS plots, likely to be a preferred habitat for breeding swans, show a more moderate rate of increase than CBC/BBS. Winter trends as measured by WeBS have shown a parallel upturn (Austin *et al.* 2008). The reductions in breeding performance, although statistically significant, may be to some extent artefacts of the relatively small and perhaps unrepresentative annual samples in the 1990s. The recent change of conservation listing from green to amber is unconnected with its UK trend.



Population changes

CBC/BBS UK 1966–2007

Mute Swan




Table of population changes for Mute Swan

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	80	199	33	473		
	25	1981-2006	117	139	64	277		
	10	1996-2006	234	7	-9	25		
	5	2001-2006	254	-14	-25	-4		
CBC/BBS England	39	1967-2006	69	181	13	560		Small CBC sample
	25	1981-2006	100	132	56	255		Small CBC sample
	10	1996-2006	201	1	-15	18		
	5	2001-2006	215	-7	-17	9		
WBS waterways	31	1975-2006	45	52	-1	152		
	25	1981-2006	47	33	3	89		
	10	1996-2006	55	2	-13	23		
	5	2001-2006	48	-7	-17	6		
BBS UK	11	1995-2006	213	10	-7	40		
	10	1996-2006	221	6	-10	24		
	5	2001-2006	254	-12	-23	-2		
BBS England	11	1995-2006	182	1	-18	21		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007

Mute Swan




Waterways Bird Survey 1974–2007

Mute Swan




BBS UK 1994–2007

Mute Swan



BBS England 1994–2007

Mute Swan




Productivity trends

Table of productivity changes for Mute Swan

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	22	Curvilinear	5.89 eggs	5.64 eggs	-4.2%	Small sample
Brood size	38	1968-2006	38	None				
Daily failure rate (eggs)	38	1968-2006	30	Curvilinear	0.61% nests/day	1.41% nests/day	131.1%	Small sample
Daily failure rate (chicks)	38	1968-2006	25	None				Small sample
Laying date	38	1968-2006	13	None				Small sample


Clutch size 1966–2007

Mute Swan




Brood size 1966–2007

Mute Swan




Egg stage nest failure rate

Mute Swan




Chick stage nest failure rate

Mute Swan



Laying date 1966–2007

Mute Swan



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GREYLAG GOOSE

Anser anser

• Population changes

• Productivity trends

• Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
 UK: not listed (introduced population);
 amber (localised NW Scottish population);
 amber (in winter, localised and >20% of NW European Flyway population)

Long-term trend

UK: rapid increase

UK population size

3,200 indigenous pairs in 1997, and 30,900 introduced adults in 1999 (Mitchell *et al.* 2000, Rehfishch *et al.* 2002, APEP06); 15,600–15,800 pairs in 2000 (BIE04)

Status summary

Apart from a small indigenous population in northwest Scotland and the Western Isles, and winter visitors mainly from Iceland, the Greylag Goose is an introduced species throughout the UK. Introduced Greylags have increased very rapidly, at a rate estimated at 12% per annum in southern Britain between the 1988–91 Atlas period and 1999 (Rehfishch *et al.* 2002). This equates across Britain to 170%, or 9.4% per annum, in the period to 2000 (Austin *et al.* 2007). The WBS sample became large enough for annual monitoring in 1992, since when further steep increase has been recorded along linear waterways. Annual breeding-season monitoring in a wider range of habitats through BBS has shown similar strong increases. Winter counts confirm that the introduced population is likely to be already much larger than the latest agreed population size estimates from 1999 and 2000 (Austin *et al.* 2008).

Population changes

Waterways Bird Survey 1992–2007 Greylag Goose





Table of population changes for Greylag Goose

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	13	1993-2006	10	507	106	1427		Small sample
	10	1996-2006	12	245	35	679		Small sample
	5	2001-2006	11	113	0	221		Small sample
BBS UK	11	1995-2006	127	120	21	281		
	10	1996-2006	133	116	33	235		
	5	2001-2006	173	58	23	89		
BBS England	11	1995-2006	103	126	64	257		
	10	1996-2006	109	116	56	215		
	5	2001-2006	142	39	16	66		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

CANADA GOOSE

Branta canadensis

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: not listed (introduced)

Long-term trend

UK: rapid increase

UK population size

82,550 adults in 1999 (Rehfishch *et al.* 2002: APEP06);
88,866 adults in Britain in 2000 (Austin *et al.* 2007)



Status summary

Canada Geese have increased rapidly, at a rate estimated at 9.3% per annum in Britain between the 1988–91 Atlas period and 2000, with no sign of any slowing in the rate of increase (Austin *et al.* 2007). Most of this increase, amounting to 166% during that period, has been in areas previously with low goose densities. The WBS sample became large enough for annual monitoring in 1980, since when further, apparently accelerating, increase on linear waterways occurred during the 1990s. WBS results for recent seasons suggest little change in this habitat since 2004. Annual breeding-season monitoring in a wider range of habitats through BBS has shown similar strong increases in England and in the UK as a whole. Winter monitoring by WeBS shows a continuing long-term increase (Austin *et al.* 2008).

Population changes

Waterways Bird Survey 1980–2007

Canada Goose






Table of population changes for Canada Goose

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	25	1981-2006	30	90	-9	698		
	10	1996-2006	38	80	41	169		
	5	2001-2006	36	12	-7	46		
BBS UK	11	1995-2006	393	115	72	159		
	10	1996-2006	409	103	64	141		
	5	2001-2006	490	46	25	64		
BBS England	11	1995-2006	367	102	58	152		
	10	1996-2006	382	92	52	141		
	5	2001-2006	453	41	17	57		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

SHELDUCK *Tadorna tadorna*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: amber (localised in winter, >20% of NW European population in winter)

Long term-trend

UK: rapid increase

UK population size

10,900 pairs in 1990 (1988–91 Atlas: [APEP06](#)); 5,800–10,800 pairs in 2000 (updated using CBC and BBS trends: [BIE04](#))

Status summary

Shelducks occurred on relatively few CBC plots, most of which were close to a coast or an estuary, and it is unclear how well the CBC trend represented the UK breeding population. The CBC showed a substantial increase from the mid 1960s until the early 1980s, some decrease during the 1980s, and stability during the 1990s, although the wide confidence intervals provide scope for other interpretations. Population increase was associated with expansion of range, measured as an additional 20% of occupied 10-km squares in Britain between 1968–72 and 1988–91 ([Gibbons *et al.* 1993](#)). The UK winter Shelduck population rose during the 1960s and 1970s, alongside the rise in breeding numbers, but has been falling again since the mid 1990s ([Austin *et al.* 2008](#)). The BBS index is affected by occasional large counts, and therefore its confidence intervals are relatively wide. BBS results suggest an accelerating increase since 1994.



Population changes

CBC all habitats 1966–2000 Shelduck





Table of population changes for Shelduck

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC all habitats	31	1968-1999	18	300	94	787		Small sample
	25	1974-1999	21	12	-40	118		
	10	1989-1999	21	3	-21	40		
	5	1994-1999	23	4	-18	39		
BBS UK	11	1995-2006	131	-2	-30	35		
	10	1996-2006	135	8	-17	38		
	5	2001-2006	144	24	0	53		
BBS England	11	1995-2006	108	21	-11	48		
	10	1996-2006	111	25	-9	52		
	5	2001-2006	116	33	-2	65		



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

MALLARD
Anas platyrhynchos

- Population changes
- Productivity trends
- Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK, England: rapid increase

UK population size

50,400–127,100 pairs in 1990 (1988–91 Atlas: [APEP06](#)); 63,000–158,900 pairs in 2000 (updated using CBC/BBS trend: [BiE04](#))

Status summary

The Mallard has increased steadily as a breeding bird in the UK since the 1960s, and especially in England, a trend to which ongoing large-scale releases for shooting may have contributed ([Marchant et al. 1990](#)). Mallards originating from domesticated birds and not resembling wild-type birds in either plumage or behaviour are very abundant but perhaps under-represented in survey data, especially since many individuals appear to be semi-captive. A large part of the increase in breeding numbers may be attributable to such birds, rather than to true-bred stock. Winter populations have declined since at least the late 1980s ([Austin et al. 2008](#)), linked apparently to a decrease in continental immigration ([Mitchell et al. 2002](#)).

Population changes

CBC/BBS UK 1966–2007
Mallard




Table of population changes for Mallard

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	427	166	111	226		
	25	1981-2006	608	34	14	55		
	10	1996-2006	1204	16	7	24		
	5	2001-2006	1286	-5	-10	0		
CBC/BBS England	39	1967-2006	360	205	138	279		
	25	1981-2006	510	44	20	71		
	10	1996-2006	1009	27	18	36		
	5	2001-2006	1078	2	-4	6		
WBS waterways	31	1975-2006	92	165	88	258		
	25	1981-2006	97	105	45	166		
	10	1996-2006	96	-3	-13	8		
	5	2001-2006	84	-4	-12	4		
BBS UK	11	1995-2006	1115	19	10	28		
	10	1996-2006	1151	16	8	24		
	5	2001-2006	1286	-3	-8	2		
BBS England	11	1995-2006	932	30	22	41		
	10	1996-2006	963	27	20	36		
	5	2001-2006	1078	4	-2	8		


BBS S	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
		5	2001-2006	96	-26	-37	-17	>25	
BBS Wales		11	1995-2006	62	-9	-45	50		
		10	1996-2006	65	-1	-37	50		
		5	2001-2006	75	25	2	54		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Mallard




Waterways Bird Survey 1974—2007

Mallard




BBS UK 1994—2007

Mallard




BBS England 1994—2007

Mallard




BBS Scotland 1994—2007

Mallard



BBS Wales 1994—2007

Mallard



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

TUFTED DUCK *Aythya fuligula*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 3 (declining)
UK: green

Long-term trend

UK: shallow increase

UK population size

7,000–8,000 pairs in GB in 1979–83 ([Owen et al. 1986](#): [APEP06](#)); 10,200–11,500 pairs in UK in 2000 (1988–91 Atlas estimate updated using WBS trend: [BIE04](#))

Status summary

The colonisation of the UK by Tufted Ducks, which began in 1849, was aided by the spread of the zebra mussel *Dreissena polymorpha*, a non-native invasive species that had been introduced accidentally to Britain a few decades earlier. The long-term shallow increase shown by WBS, and the 15% increase in range in Britain between the two atlas periods ([Gibbons et al. 1993](#)) may indicate that population expansion and in-filling of range are still occurring. BBS data suggest significant further increase since 1994, in England and in the UK as a whole. The species' winter trend in the UK since the 1960s, which includes many continental visitors, is also shallowly upward overall ([Austin et al. 2008](#)). In contrast, moderate recent declines elsewhere in northern Europe have resulted in reclassification as a species of conservation concern ([BirdLife International 2004](#)).



Population changes

Waterways Bird Survey 1974–2007 Tufted Duck






Table of population changes for Tufted Duck

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	23	15	-57	168		
	25	1981-2006	24	4	-53	127		
	10	1996-2006	25	-6	-42	56		
BBS UK	5	2001-2006	17	-12	-36	23		Small sample
	11	1995-2006	136	46	14	81		
	10	1996-2006	140	40	10	67		
BBS England	5	2001-2006	145	16	-8	39		
	11	1995-2006	117	27	-1	64		
	10	1996-2006	121	21	-3	54		
	5	2001-2006	125	11	-6	30		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

GOOSANDER

Mergus merganser

• Population changes

• Productivity trends

• Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK: rapid increase

UK population size

2,600 (2,300–2,900) pairs in 1987 (Gregory *et al.* 1997: APEP06); 2,900–3,600 pairs in 2000 (updated using WBS trend: BiE04)

Status summary

Goosanders were first discovered to have colonised the UK in Perthshire in 1871, and spread from Scotland into northern England in the 1940s (Holloway 1996). Between the two breeding atlases, the species expanded its range in northern England, and colonised Wales and southwest England. WBS samples became large enough for annual monitoring in 1980, and have shown sustained population increase. The BTO's two national surveys of sawbills demonstrated an average increase in population size of 3% per annum between 1987 and 1997 (Rehfishch *et al.* 1999). Reasons for the colonisation of the UK, and the subsequent range expansion and population increase, are unknown. The species' winter trend in Britain, comprising British breeders and continental visitors, rose at an accelerating rate from the late 1960s to the mid 1990s, but subsequently began to decline (Austin *et al.* 2008).

Population changes

Waterways Bird Survey 1980–2007 Goosander




Table of population changes for Goosander

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	25	1981-2006	23	144	18	405		
	10	1996-2006	25	32	-12	88		
	5	2001-2006	24	26	-2	60		

Productivity trends

Productivity information is not currently available for this species

Additional information

- Distribution maps for this species are not currently available online (see [Atlases species help](#))
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

RED GROUSE *Lagopus lagopus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
 UK: amber (25–50% population decline)
 UK Biodiversity Action Plan: priority species

Long-term trend

UK: decline

UK population size

155,000 pairs in 2000 (1988–91 Atlas estimate updated using GCT gamebag data: [BiE04](#), [APEP06](#))



Status summary

The distinctive dark-winged race *scotica* is endemic to Britain and Ireland and has the vast bulk of its population within the UK. BBS shows no overall trend since 1994. **Shooting bags** have revealed long-term declines, apparently driven by loss of heather moorland, increased predation from corvids and foxes, and an increasing incidence of viral disease ([Hudson 1992](#), [Newton 2004](#)), which have prompted the move of the species from the Green to the Amber List. Raptor predation is believed not to affect breeding populations significantly, although it can reduce numbers in the post-breeding period ([Redpath & Thirgood 1997](#)). Longer-term trends in Red Grouse abundance are overlain by cycles, with periods that vary regionally, linked to the dynamics of infection by a nematode parasite ([Dobson & Hudson 1992](#), [Gibbons et al. 1993](#)).

Population changes

BBS UK 1994–2007 Red Grouse




Table of population changes for Red Grouse


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	106	-14	-27	-2		
	10	1996-2006	108	-20	-31	-7		
	5	2001-2006	105	-24	-35	-12		
BBS England	11	1995-2006	48	-14	-30	3		
	10	1996-2006	50	-16	-31	-1		
	5	2001-2006	56	-19	-32	-7		
BBS Scotland	11	1995-2006	53	-15	-32	10		
	10	1996-2006	52	-22	-38	0		
	5	2001-2006	43	-26	-44	-6	>25	



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS England 1994—2007

Red Grouse



BBS Scotland 1994—2007

Red Grouse



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

RED-LEGGED PARTRIDGE

Alectoris rufa

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 2 (declining)

UK: not listed (introduced)

Long-term trend

UK, England: shallow decline

UK population size

72,000–200,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

Since Red-legged Partridge is a non-native species released in the UK for the purpose of being shot by hunters, its population decrease over the recent 25-year period raises no conservation concern. Moreover, BBS data indicate that significant increase has occurred in the UK since 1994. [Game-bag data](#) show that the numbers released per unit area onto shooting estates, and the numbers shot, have both increased by a remarkable eightfold since 1980. Around 2 million birds were being released each year during the 1990s ([Tapper 1999](#)). The effects of large-scale releases of this species and [Pheasant](#) on native fauna have been little studied. It is now believed, however, that shooting operations based on large-scale releases of Red-legged Partridges can lead to local extinction of the native [Grey Partridge](#) ([Watson et al. 2007](#)).



Population changes

CBC/BBS UK 1966–2007 Red-legged Partridge




Table of population changes for Red-legged Partridge


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	167	-16	-46	35		
	25	1981-2006	243	-21	-45	7		
	10	1996-2006	497	24	13	35		
	5	2001-2006	549	18	10	25		
CBC/BBS England	39	1967-2006	164	-19	-47	18		
	25	1981-2006	238	-24	-44	-3		
	10	1996-2006	487	20	10	31		
	5	2001-2006	535	17	9	25		
BBS UK	11	1995-2006	465	29	17	42		
	10	1996-2006	481	24	14	36		
	5	2001-2006	549	16	9	22		
BBS England	11	1995-2006	456	24	13	36		
	10	1996-2006	472	21	10	32		
	5	2001-2006	535	14	8	20		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Red-legged Partridge




BBS UK 1994—2007

Red-legged Partridge



BBS England 1994—2007

Red-legged Partridge



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GREY PARTRIDGE

Perdix perdix

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (vulnerable)

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK, England: rapid decline

UK population size

70,000–75,000 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))



Status summary

This species has declined enormously, probably because of the effects of agricultural intensification (specifically herbicides) on the food plants of young chicks' insect prey ([Potts 1986](#)). Despite years of research and the application of a government Biodiversity Action Plan ([Aebischer & Ewald 2004](#)), the continuing decline shown by CBC/BBS suggests that all efforts to boost the population have so far been unsuccessful. Local extinctions are now likely to be widespread, but masked in some areas by continuing releases of hand-reared birds onto shooting estates. Artificial rearing has increased since the mid 1980s, despite the failure of restocking as a means of restoring breeding numbers (see [here](#)), while releases of non-native gamebirds, which have increased greatly, can be detrimental to the native species. Infection with caecal nematodes from farm-reared [Pheasants](#) may be contributing to the decline of Grey Partridges in Britain ([Tompkins et al. 2002](#)). The practice of releasing [Red-legged Partridges](#) in large numbers can lead to Grey Partridge extinction, in part because shooters are unable to distinguish these two species ([Watson et al. 2007](#)): these authors conclude that overshooting has greater implications for Grey Partridge conservation than raptor predation. Grey Partridge is one of the most strongly declining bird species in Europe, having decreased at an annual rate of 7% during 1980–2005 ([PECBMS 2007](#)).

Population changes

CBC/BBS UK 1966–2007

Grey Partridge




Table of population changes for Grey Partridge

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	119	-88	-91	-83	>50	
	25	1981-2006	142	-78	-85	-71	>50	
	10	1996-2006	245	-39	-48	-31	>25	
	5	2001-2006	222	-13	-24	-1		
CBC/BBS England	39	1967-2006	107	-88	-91	-82	>50	
	25	1981-2006	126	-78	-84	-71	>50	
	10	1996-2006	220	-33	-41	-25	>25	
	5	2001-2006	202	-4	-14	6		
BBS UK	11	1995-2006	227	-40	-49	-32	>25	
	10	1996-2006	230	-38	-47	-30	>25	
	5	2001-2006	222	-14	-25	-5		
BBS England	11	1995-2006	202	-35	-44	-26	>25	
	10	1996-2006	205	-31	-41	-23	>25	


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Grey Partridge




BBS UK 1994—2007

Grey Partridge



BBS England 1994—2007

Grey Partridge



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

PHEASANT *Phasianus colchicus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: not listed (introduced)

Long-term trend

England: moderate increase

UK population size

1,800,000–1,900,000 females in 2000 (Robertson *et al.* 1989, updated using CBC/BBS trend: BiE04, APEP06)

Status summary

Pheasants have increased in abundance since the 1960s, at a rate that appears to be accelerating, but it must be noted that numbers of this introduced gamebird are determined principally by releases of reared birds for shooting (Marchant *et al.* 1990). The Game Conservancy Trust estimates that about 20–22 million birds are released in the UK each autumn, a figure that has increased approximately four-fold since the mid 1960s (Tapper 1999; also [here](#)). More than two million newly released birds are expected to survive until spring, when they must form the major part of the breeding population. The BBS records increase in England and Wales, but little change in Scotland since 1994. During 1968–88, a period when the total biomass of birds in Britain fell by an estimated 10%, CBC data indicate that Pheasant biomass rose by about 2,500 tonnes – more than ten times more than any other species (Dolton & Brooke 1999). High Pheasant densities potentially have negative effects, that have not been adequately studied, on native UK birds: these include the effect on the structure of the field layer, the spread of disease and parasites, and competition for food (Fuller *et al.* 2005). Infection with caecal nematodes from farm-reared Pheasants may be contributing to the decline of Grey Partridges in Britain (Tompkins *et al.* 2002).



Population changes

CBC/BBS England 1966–2007 Pheasant








Table of population changes for Pheasant

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	475	99	53	187		
	25	1981-2006	686	92	58	130		
	10	1996-2006	1378	36	29	42		
	5	2001-2006	1484	20	16	24		
BBS UK	11	1995-2006	1511	36	28	43		
	10	1996-2006	1561	35	29	42		
	5	2001-2006	1764	19	15	23		
BBS England	11	1995-2006	1276	39	30	47		
	10	1996-2006	1317	37	29	43		
	5	2001-2006	1484	20	14	23		
BBS Scotland	11	1995-2006	114	6	-15	30		
	10	1996-2006	116	10	-8	33		
	5	2001-2006	125	8	-3	23		

BBS W	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
		5	2001-2006	103	52	29	69		
BBS N.Ireland		11	1995-2006	30	148	46	239		
		10	1996-2006	33	111	40	167		
		5	2001-2006	40	41	12	75		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

RED-THROATED DIVER

Gavia stellata

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (depleted)
UK: amber (25–50% population decline)

Long-term trend

Shetland: moderate decline

UK population size

935–1,500 pairs in 1994 (Gibbons *et al.* 1997: BIE04, APEP06)

Status summary

Population trends are not monitored by the BTO, but JNCC's [Seabird Monitoring Programme](#) shows that breeding numbers at sample study areas in Shetland fluctuated without long-term change during 1980–2005, with low points in 1980, 2000 and 2004 (Mavor *et al.* 2008). Complete surveys of Shetland indicated a decrease of 36% there between 1983 and 1994, however (Gibbons *et al.* 1997). Since in 1994 Shetland held 28–45% of the total UK population, this warrants amber listing for Red-throated Diver, in addition to its depleted status in Europe as a whole. Since the 1980s, there may have been some tendency for more pairs to hatch a second chick, although two-chick broods are only occasional in Orkney and the proportion of nest records from there could have changed over time.



Population changes

Annual breeding population changes are not currently monitored by BTO for this species


Productivity trends

Table of productivity changes for Red-throated Diver

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	26	1980-2006	21	None				Small sample
Brood size	26	1980-2006	32	Linear increase	1.25 chicks	1.52 chicks	22.3%	
Daily failure rate (eggs)	26	1980-2006	12	Linear increase	0.59% nests/day	2.26% nests/day	283.1%	Small sample
Daily failure rate (chicks)	26	1980-2006	17	None				Small sample


Clutch size 1966–2007

Red-throated Diver




Brood size 1966–2007

Red-throated Diver




Egg stage nest failure rate

Red-throated Diver



Chick stage nest failure rate

Red-throated Diver



Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- Distribution maps for this species are not currently available online (see [Atlases species help](#))

- [BirdFacts page on species biology](#)

LITTLE GREBE *Tachybaptus ruficollis*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK: uncertain

UK population size

5,900–12,000 pairs in 1990 (1988–91 Atlas: [APEP06](#)); 3,800–13,000 pairs in 2000 (updated using CBC and WBS trends: [BiE04](#))

Status summary

The rapid decline shown by the WBS may reveal problems among birds on linear waterways during the early 1980s and since the late 1990s, while shallow increases shown by the CBC and by BBS may suggest that wider populations (including birds on small still waters) are healthy. Because of the shortage of data, and the conflict between WBS and BBS assessments, the rapid decline indicated by WBS has not triggered a conservation listing. In an analysis of nest record cards, [Moss & Moss \(1993\)](#) found that nests on ponds and lakes were significantly more successful than those on rivers and streams and that nests on rivers, subject to fluctuating water levels, experienced significantly higher failure rates through flooding than those on canals, where water levels are artificially maintained. Winter numbers, monitored by [WeBS](#), have shown sustained shallow increase ([Austin et al. 2008](#)).



Population changes

Waterways Bird Survey 1974–2007 Little Grebe





Table of population changes for Little Grebe

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	16	-59	-87	-9	>50	Small sample
	25	1981-2006	15	-64	-87	-23	>50	Small sample
	10	1996-2006	11	-30	-72	22		Small sample
BBS UK	11	1995-2006	62	25	-4	61		
	10	1996-2006	65	22	-6	52		
	5	2001-2006	77	4	-13	24		
BBS England	11	1995-2006	51	12	-21	65		
	10	1996-2006	53	6	-22	51		
	5	2001-2006	62	-5	-23	23		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GREAT CRESTED GREBE

Podiceps cristatus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK: probable increase

UK population size

9,400 adults in 1990 (1988–91 Atlas: [APEP06](#)); 6,100 pairs in 2000 (updated using BBS trend: [BiE04](#))

Status summary

This species was believed to be on the verge of extinction in Britain around 1860, when only 32–72 pairs were known in England ([Holloway 1996](#)). A subsequent increase followed reductions in persecution, aided by statutory protection, and the creation of habitat in the form of gravel pits ([Gibbons et al. 1993](#)). Increase was tracked by special surveys to around 7,000 adult birds in Britain by 1975 ([Hughes et al. 1979](#)). The BBS provides the first annual, national monitoring of this species and indicates shallow increase since 1994. Winter numbers, monitored by [WeBS](#), have shown sustained shallow increase ([Austin et al. 2008](#)).



Population changes

BBS UK 1994–2007 Great Crested Grebe




Table of population changes for Great Crested Grebe


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	63	31	-23	112		
	10	1996-2006	65	38	-19	124		
	5	2001-2006	71	54	-8	160		
BBS England	11	1995-2006	58	-11	-27	15		
	10	1996-2006	58	-8	-25	13		
	5	2001-2006	65	5	-13	23		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

BBS England 1994–2007

Great Crested Grebe



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

CORMORANT *Phalacrocorax carbo*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: amber (breeding localised, >20% of European population in winter)

Long-term trend

UK: increase

UK population size

9,018 pairs in 1998–2002 (Mitchell *et al.* 2004: APEP06); 9,100 pairs including Channel Islands (BiE04)

Status summary

The Cormorant was almost exclusively a coastal breeder in the UK until 1981, but has since established colonies in many inland areas of eastern and central England (Rehfishch *et al.* 1999; Newson *et al.* 2006). Breeding numbers and productivity at sample colonies have been monitored annually since 1986 by JNCC's **Seabird Monitoring Programme**. Overall in Britain and Ireland there was a 15% increase in the population between full surveys in 1985–88 and 1998–2002 (Mitchell *et al.* 2004). Trends during 1986–2005 show decreases in Scotland and in northeast and southwest England, but no trend in Wales, and steep increases inland in England and in regions bordering the northern part of the Irish Sea (Mavor *et al.* 2008). By 2005, breeding had been recorded at 58 inland sites, and the inland population had risen to about 2,130 pairs (Newson *et al.* 2007). Inland breeding in England is thought to have been sparked by birds of the continental race *sinensis* from the Netherlands and Denmark, although many nominate *carbo* from coastal colonies in Wales and England have contributed to its development. The winter trend in Britain, comprising British and Irish breeders and continental visitors, has shown strong increase since the late 1980s but now appears more stable (Austin *et al.* 2008). The species has recently been moved from the green to the amber list, for reasons unconnected with its UK trend.



Population changes

BBS UK 1994–2007 Cormorant




Table of population changes for Cormorant


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	201	23	-5	58		Non-breeders included
	10	1996-2006	209	23	2	52		Non-breeders included
	5	2001-2006	257	-8	-20	5		Non-breeders included
BBS England	11	1995-2006	165	22	1	49		Non-breeders included
	10	1996-2006	172	22	2	46		Non-breeders included
	5	2001-2006	213	5	-7	18		Non-breeders included



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

BBS England 1994–2007

Cormorant



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)



Page not found

[home](#) » [birdtrends2008](#) » Page not found



Terribly sorry...


The page you are looking for could not be found.

- Try searching using the search bar (top right)
- Navigate to the area you are looking for using the main menu (above)
- [Return to our homepage](#)
- Email us about what you are trying to find, or to tell us about a broken link - info@bto.org

Share this page



Related content



ARTICLE

Results

See live NBC results and watch the breeding season unfold



EVENT

Northern Ireland Seabird Report Launch

The NI Seabird Report is unique in the UK, presenting the most up-to-date seabird monitoring data collected by volunteers, NGOs and NIEA from around Northern Ireland. Each March we present freshly produced hard copies...



BLOG

Widening BTO's appeal

Andy Clements, BTO's Chief Executive, looks at how BTO can engage new audiences.

BTO - Looking out for birds

[About us](#) [Jobs](#) [National offices](#) [Community](#) [Events](#) [Blog](#) [Shop](#)

✉ info@bto.org

[Press releases](#) [Terms & Conditions](#) [Privacy](#) [Cookies](#)



Registered Charity Number 216652 (England & Wales), SC039193 (Scotland)

© British Trust for Ornithology, BTO, The Nunnery, Thetford, Norfolk IP24 2PU, Tel: +44 (0)1842 750050 Fax: +44 (0)1842 750030

Seen a ringed / dead / diseased bird?

[REPORT IT TO US](#) ▶

Keep up to date with BTO

[E-NEWSLETTER SIGN UP](#)

Flock together



May we use cookies to monitor performance and improve your experience?

May we use cookies to monitor performance and improve your experience?

These optional cookies are used for collecting anonymous visit data only.

[READ OUR COOKIE POLICY](#)

ACCEPT ALL COOKIES

Set preferences

HEN HARRIER

Circus cyaneus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3, vulnerable

UK: red (historical decline)

Long-term trend

UK: stable (between 1988–89 and 1998)

UK population size

570 (500–640) territorial pairs in 1998 (Sim *et al.* 2001: BiE04, APEP06); 806 (732–889) territorial pairs in 2004 (Sim *et al.* 2007a)

Status summary

Red listed because of substantial declines over the last two centuries, this species has suffered in recent decades from loss of habitat as forestry plantations have matured (Bibby & Etheridge 1993) but more especially from continuing illegal persecution on grouse moors (Etheridge *et al.* 1997). Although the Hen Harrier and other raptors have been protected under UK law since 1961, many are still killed unlawfully to protect Red Grouse stocks. The UK population was unchanged between surveys in 1988–89 and 1998, with declines in Orkney and England but increases in Northern Ireland and the Isle of Man (Sim *et al.* 2001). A decrease of 70% in the Orkney population over the last 20 years has been linked to reductions in the area of unmanaged grassland (Amar & Redpath 2005); the demographic drivers of this decline have been a decrease in polygyny and reduced nesting success among secondary females (Amar *et al.* 2005). The latest survey reveals a 41% increase in the UK and Isle of Man during 1998–2004, but with decreases in the Southern Uplands, east Highlands and England, all being areas with many managed grouse moors (Sim *et al.* 2007a). Although average clutch size declined substantially during the 1980s, further investigation has shown that this trend is due to the increased proportions in recent years of records from Orkney, where clutch sizes tend to be smaller than on the mainland (Summers 1998, Crick 1998). Recent results confirm that rough grass is a critical habitat for Orkney Hen Harriers, providing the necessary food during the incubation period (Amar *et al.* 2008).



Population changes


Annual breeding population changes for this species are not currently monitored by BTO

Productivity trends


Table of productivity changes for Hen Harrier

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	Curvilinear	5.48 eggs	4.73 eggs	-13.6%	Small sample
Brood size	38	1968-2006	20	None				Small sample
Daily failure rate (eggs)	38	1968-2006	11	Curvilinear	0.03% nests/day	0.22% nests/day	633.3%	Small sample
Daily failure rate (chicks)	38	1968-2006	14	None				Small sample


Clutch size 1966–2007




Brood size 1966–2007



Egg stage nest failure rate



Chick stage nest failure rate



Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- Distribution maps for this species are not currently available online (see [Atlases species help](#))
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

SPARROWHAWK

Accipiter nisus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

England: rapid increase

UK population size

40,100 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary



Sparrowhawks suffered a severe population crash caused by organochlorine pesticides in the 1950s and 1960s, when the species was extinguished from large areas of lowland Britain ([Newton 1986](#)). Following a ban on the use of organochlorines, the species increased and spread, and became common enough on CBC plots for annual monitoring in the early 1970s. Between then and the mid 1990s, the CBC charted a steep increase. Many former haunts especially in the Midlands and east of England were reoccupied between the two atlas periods ([Gibbons *et al.* 1993](#)). Improving breeding performance is likely to have contributed to this remarkable period of success: failure rates at the egg stage (c.44 days from laying the first egg) fell markedly from high initial values, and brood sizes increased throughout. The population has stabilised since the mid 1990s and, possibly through the effects of intraspecific competition, average brood size has begun to drop again.

Population changes

CBC/BBS England 1974–2007

Sparrowhawk






Table of population changes for Sparrowhawk

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	31	1975-2006	131	166	78	360		
	25	1981-2006	156	69	28	139		
	10	1996-2006	291	2	-8	12		
	5	2001-2006	300	-2	-10	7		
BBS UK	11	1995-2006	315	5	-7	16		
	10	1996-2006	325	5	-5	16		
	5	2001-2006	362	0	-9	10		
BBS England	11	1995-2006	262	0	-11	12		
	10	1996-2006	269	2	-8	12		
	5	2001-2006	300	-3	-10	5		








The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Table of productivity changes for Sparrowhawk

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	36	None				
Brood size	38	1968-2006	71	Curvilinear	3.11 chicks	3.45 chicks	10.9%	
Daily failure rate (eggs)	38	1968-2006	34	Linear decline	0.47% nests/day	0.09% nests/day	-80.9%	
Daily failure rate (chicks)	38	1968-2006	48	None				
Laying date	38	1968-2006	14	None				Small sample



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

BUZZARD
Buteo buteo

- Population changes
- Productivity trends
- Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK, England: rapid increase

UK population size

31,100–44,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#)); 44,000–61,000 territorial pairs in GB in 2001 ([Clements 2002](#))

Status summary

The Common Buzzard has shown a substantial eastward range expansion since the 1988–91 Atlas, and is arguably now the most abundant diurnal raptor in Britain ([Clements 2002](#)). The increasing trend identified by the CBC relates especially to the spread of range into central and eastern Britain, where CBC was more strongly represented. If anything, however, the upsurge has been amplified with the addition of the more geographically representative BBS data since 1994. The increase has been associated with improving nesting success, perhaps through reduced persecution, the recovery of rabbit populations from the effects of myxomatosis and release from the deleterious effects of organochlorine pesticides ([Elliott & Avery 1991](#), [Clements 2002](#)).

Population changes

CBC/BBS UK 1966–2007
Buzzard




Table of population changes for Buzzard

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	216	419	251	1372		
	25	1981-2006	328	197	105	392		
	10	1996-2006	725	39	29	50		
	5	2001-2006	887	7	2	14		
CBC/BBS England	39	1967-2006	130	483	268	1367		Small CBC sample
	25	1981-2006	198	286	167	681		
	10	1996-2006	440	77	61	100		
	5	2001-2006	559	32	24	41		
BBS UK	11	1995-2006	669	47	37	60		
	10	1996-2006	703	39	31	50		
	5	2001-2006	887	9	3	16		
BBS England	11	1995-2006	400	90	68	117		
	10	1996-2006	424	78	58	96		
	5	2001-2006	559	32	24	40		
BBS Scotland	11	1995-2006	115	24	6	51		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Wales	11	1995-2006	132	5	-8	23		
	10	1996-2006	138	7	-4	22		
	5	2001-2006	161	2	-6	11		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Buzzard




BBS UK 1994—2007

Buzzard




BBS England 1994—2007

Buzzard




BBS Scotland 1994—2007

Buzzard



BBS Wales 1994—2007

Buzzard




Productivity trends

Table of productivity changes for Buzzard

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	31	None				
Brood size	38	1968-2006	94	Curvilinear	1.87 chicks	1.95 chicks	4.6%	
Daily failure rate (eggs)	38	1968-2006	26	Linear decline	0.7% nests/day	0.12% nests/day	-82.9%	Small sample
Daily failure rate (chicks)	38	1968-2006	47	None				


Clutch size 1966—2007

Buzzard




Brood size 1966—2007

Buzzard



Egg stage nest failure rate


Buzzard



Insufficient data on laying dates available for this species

Chick stage nest failure rate

Buzzard



Insufficient data on CES available for this species

Additional information

- Distribution maps for this species are not currently available online (see [Atlases species help](#))
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

KESTREL *Falco tinnunculus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 3, declining
UK: amber (25–50% population decline)

Long-term trend

England: fluctuating, with no long-term trend

UK population size

36,800 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

Kestrels had recovered from the lethal and sublethal effects of organochlorine pesticides by the mid 1970s, the recovery probably driven by improving nesting success, but subsequently entered a decline which has been linked to the effects of agricultural intensification on farmland habitats and their populations of small mammals ([Gibbons *et al.* 1993](#)). Since the mid 1980s, the English population has fluctuated without a long-term trend being apparent. In Scotland, however, there has been a significant decline since 1994. The failure rate at the egg stage (c.28 days from laying the first egg) has declined substantially since the 1970s; brood sizes increased up to 1990, but a subsequent decline has resulted in the inclusion of Kestrel in the NRS concern list ([Leech & Barimore 2008](#)). Despite its decline since the mid 1970s, the Kestrel breeds at high density in mixed farmland across much of England, suggesting that the British population may number more than 50,000 pairs ([Clements 2008](#)).



Population changes

CBC/BBS England 1966–2007 Kestrel





Table of population changes for Kestrel

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	216	6	-23	53		
	25	1981-2006	293	-21	-35	-1		
	10	1996-2006	548	5	-2	14		
BBS UK	5	2001-2006	570	5	-2	11		
	11	1995-2006	586	-12	-20	-5		
	10	1996-2006	599	-7	-16	0		
BBS England	5	2001-2006	657	1	-6	8		
	11	1995-2006	507	0	-8	8		
	10	1996-2006	517	4	-3	12		
BBS Scotland	5	2001-2006	570	5	-1	10		
	11	1995-2006	42	-40	-56	-17	>25	
	10	1996-2006	42	-34	-50	-12	>25	
	5	2001-2006	44	-13	-34	7		


BBS UK 1994–2007

Kestrel




BBS England 1994–2007

Kestrel



BBS Scotland 1994–2007

Kestrel




Productivity trends

Table of productivity changes for Kestrel

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	54	None				
Brood size	38	1968-2006	121	Curvilinear	3.74 chicks	3.83 chicks	2.5%	
Daily failure rate (eggs)	38	1968-2006	40	Linear decline	0.55% nests/day	0.09% nests/day	-83.6%	
Daily failure rate (chicks)	38	1968-2006	65	Linear decline	0.22% nests/day	0.09% nests/day	-59.1%	
Laying date	38	1968-2006	21	Linear decline	May 4	Apr 28	-6 days	Small sample


Clutch size 1966–2007

Kestrel




Brood size 1966–2007

Kestrel




Egg stage nest failure rate

Kestrel




Chick stage nest failure rate

Kestrel



Laying date 1966–2007

Kestrel



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

MERLIN
Falco columbarius

- Population changes
- Productivity trends
- Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
 UK: amber (historical decline)

Long-term trend

UK: probable increase

UK population size

1,300 pairs in 1990–94 (Rebecca & Bainbridge 1998: BiE04, APEP06)

Status summary

Having declined substantially over the past two centuries, Merlin shows indications of a recent doubling of population (Rebecca & Bainbridge 1998). This increase may be associated with an increased use of forest edge as a nesting habitat (Parr 1994). Because of its recent population upturn, the species has been moved from the red to the amber list. It remains much too scarce, however, for annual population monitoring via BBS: dedicated observers and specialised field methods are required, as described by Hardey et al. (2006). Submissions to the Rare Breeding Birds Panel fall well short of the estimated UK total population but show an average of 1.86 young fledged per occupied territory during 1996–2004 (Holling & RBBP 2007a). Breeding performance has tended to improve since the 1960s, probably linked to the declining influence of organochlorine pesticides (Crick 1993). Hatching rates in the southeast Yorkshire Dales were consistently higher than had been recorded in earlier studies in Northumberland (Wright 2005).





Population changes

Annual breeding population changes for this species are not currently monitored by BTO

Productivity trends

Table of productivity changes for Merlin

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	38	None				
Brood size	38	1968-2006	55	Linear increase	3.52 chicks	3.79 chicks	7.8%	
Daily failure rate (eggs)	38	1968-2006	26	Linear decline	0.67% nests/day	0.24% nests/day	-64.2%	Small sample
Daily failure rate (chicks)	38	1968-2006	29	Linear decline	0.94% nests/day	0.24% nests/day	-74.5%	Small sample



Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

HOBBY *Falco subbuteo*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK: increase

UK population size

2,200 pairs in 2000 (Clements 2001: BiE04, APEP06)

Status summary

This species cannot be monitored by any of the standard monitoring schemes, due to its low population density and unobtrusive habits. Many sightings must refer to migrants, first-summer non-breeders, or to breeding birds from distant nests, and do not help to establish whether nesting occurs in the locality. Dedicated observers and specialised field methods are required, as described by [Hardey *et al.* \(2006\)](#). The Rare Breeding Birds Panel collects annual data, which under-represent the true population to unknown degrees, but adequately establish the long-term upward trend ([Holling & RBBP 2008](#)). The Hobby's distribution has spread markedly northwards in England since the 1970s ([Gibbons *et al.* 1993](#)), perhaps linked to increases in its dragonfly prey supplies ([Prince & Clarke 1993](#)) and to a decreasing dependency on its traditional heathland habitat, but the reasons underlying the increase are still only speculative ([Clements 2001](#)). A success rate of more than 90% was recorded for nests in Derbyshire during 1992–2001, with successful nests fledging a mean of 2.44 young ([Messenger & Roome 2007](#)). The small annual samples of nest record cards indicate no long-term change in either brood size or nest success.



Population changes


Annual breeding population changes for this species are not currently monitored by BTO

Productivity trends

Table of productivity changes for Hobby

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size	38	1968-2006	18	None				Small sample
Daily failure rate (chicks)	38	1968-2006	12	None				Small sample

Brood size 1966–2007
Hobby



Insufficient data on clutch size available for this species

Insufficient data on nest failure available for this species

Insufficient data on laying date available for this species

Insufficient data on nestling failure available for this species

Insufficient data on CES available for this species

Additional information

- Distribution maps for this species are not currently available online (see [Atlases species help](#))
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

PEREGRINE *Falco peregrinus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: amber (European status)

Long-term trend

UK, England: increase
Northwest Scotland: decline since 1991

UK population size

1,283 pairs in 1991 (Crick & Ratcliffe 1995: APEP06);
1,400 pairs in 2002 (Banks *et al.* 2003: BiE04)

Status summary

The UK population size, distribution and breeding performance have all largely recovered from the detrimental effects of organochlorine pesticides in the 1950s and 1960s. Populations and breeding performance have declined recently, however, in northwest Scotland and the Northern Isles (Crick & Ratcliffe 1995), and nest record information for the UK as a whole shows a significant decline in clutch size, although samples for the first ten years are small. The number of UK breeding pairs has been censused every ten years since 1961 by BTO/JNCC/RSPB/Raptor Study Groups, and has been estimated as follows: 1961 – 385 pairs; 1971 – 489 pairs; 1981 – 728 pairs; 1991 – 1,283 pairs (Ratcliffe 1993). The National Peregrine Survey 2002 found 1,402 breeding pairs, a further 10% increase overall since 1991 but with declines in north and west Scotland, North Wales and Northern Ireland (Banks *et al.* 2003); around 50 pairs were missed in Wales, however (Dixon *et al.* 2008). Similar increases across Europe have resulted in a downgrading of conservation listing from 'SPEC 3 (rare)' to 'secure' (BirdLife International 2004).



Population changes


Annual population changes are not monitored for this species

Productivity trends


Table of productivity changes for Peregrine

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	16	Linear decline	3.58 eggs	3.09 eggs	-13.7%	Small sample
Brood size	38	1968-2006	41	None				
Daily failure rate (eggs)	38	1968-2006	21	Curvilinear	0.14% nests/day	0.28% nests/day	100%	Small sample
Daily failure rate (chicks)	38	1968-2006	23	None				Small sample


Clutch size 1966–2007
Peregrine




Brood size 1966–2007
Peregrine



Egg stage nest failure rate
Peregrine



Chick stage nest failure rate
Peregrine



Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- Distribution maps for this species are not currently available online (see [Atlases species help](#))
- [BirdFacts page on species biology](#)

MOORHEN *Gallinula chloropus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK: fluctuating, with no long-term trend

UK population size

270,000 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

While the long-term CBC/BBS trend is of shallow increase, much of the population increase took place before 1974, when WBS monitoring began, and may have been a recovery from heavy mortality during the cold winters of the early 1960s. On both CBC/BBS and WBS evidence, there was decrease during the 1970s and 1980s, but this has been followed by a partial recovery. A decline in the number and quality of farmland ponds, and the spread of American mink *Mustela vison*, which is an important predator especially along watercourses, have been suggested as possible causes of decline. The decline has been associated with significant reductions in breeding performance. Average clutch size has declined and the failure rate of nests over the full 25-day egg period (20 days for incubation and 5 days for laying) has increased, earning the species a place on the NRS concern list ([Leach & Barimore 2008](#)). Average brood sizes have increased, however.



Population changes

Waterways Bird Survey 1974–2007 Moorhen




Table of population changes for Moorhen


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	260	27	0	59		
	25	1981-2006	347	-3	-18	11		
	10	1996-2006	640	16	8	25		
	5	2001-2006	656	0	-6	5		
CBC/BBS England	39	1967-2006	237	31	2	62		
	25	1981-2006	318	-2	-14	16		
	10	1996-2006	587	14	6	23		
	5	2001-2006	605	-2	-8	3		
WBS waterways	31	1975-2006	78	-8	-32	31		
	25	1981-2006	81	3	-23	38		
	10	1996-2006	80	3	-16	24		
	5	2001-2006	69	4	-10	18		
BBS UK	11	1995-2006	586	17	8	26		
	10	1996-2006	599	17	9	25		
	5	2001-2006	656	1	-5	7		
BBS England	11	1995-2006	538	13	3	23		
	10	1996-2006	549	14	5	23		

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS UK 1966—2007
Moorhen




CBC/BBS England 1966—2007
Moorhen



BBS UK 1994—2007
Moorhen



BBS England 1994—2007
Moorhen




Productivity trends


Table of productivity changes for Moorhen

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	91	Linear decline	6.52 eggs	5.96 eggs	-8.6%	
Brood size	38	1968-2006	77	Curvilinear	3.12 chicks	4.3 chicks	38.1%	
Daily failure rate (eggs)	38	1968-2006	110	Curvilinear	1.35% nests/day	2.15% nests/day	59.3%	
Daily failure rate (chicks)	38	1968-2006	36	None				
Laying date	38	1968-2006	68	Linear decline	May 10	May 5	-5 days	


Clutch size 1966—2007
Moorhen



Brood size 1966—2007
Moorhen




Egg stage nest failure rate
Moorhen



Insufficient data on nestling failure available for this species

Laying date 1966—2007
Moorhen



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

COOT *Fulica atra*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK: moderate increase

UK population size

22,600–28,800 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS and WBS trends: [BIE04](#), [APEP06](#))



Status summary

WBS and CBC/BBS trends for Coot indicate a long-term increase, although the magnitude of the change is not clear. Small CBC samples, mainly of birds on small water-bodies, suggested a rapid rise in the late 1960s. WBS and BBS both include more birds on larger waters, and so may be more representative of Coot populations, but WBS has not recorded the strong increase found by BBS observers since 1994. The combination of CBC and BBS data suggests that the long-term increase in the UK and England may have been rapid. Winter abundance on large still waters, as monitored by WeBS, showed shallow increase from the mid 1980s to around 2000/01 but has since declined, especially in Northern Ireland ([Austin et al. 2008](#)).

Population changes

Waterways Bird Survey 1974–2007 Coot




Table of population changes for Coot


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	95	193	84	538		
	25	1981-2006	134	38	-6	89		
	10	1996-2006	259	26	6	52		
	5	2001-2006	274	-4	-15	11		
CBC/BBS England	39	1967-2006	87	186	86	484		
	25	1981-2006	121	36	-3	87		
	10	1996-2006	234	25	8	49		
	5	2001-2006	248	-5	-15	9		
WBS waterways	31	1975-2006	39	66	-16	223		
	25	1981-2006	42	40	-24	170		
	10	1996-2006	45	4	-42	41		
	5	2001-2006	36	8	-20	34		
BBS UK	11	1995-2006	235	34	15	64		
	10	1996-2006	243	27	11	52		
	5	2001-2006	274	-4	-11	8		
BBS England	11	1995-2006	212	34	11	56		
	10	1996-2006	219	27	8	46		
	5	2001-2006	248	-5	-16	7		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS UK 1966—2007

Coot




CBC/BBS England 1966—2007

Coot




BBS UK 1994—2007

Coot



BBS England 1994—2007

Coot



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

OYSTERCATCHER *Haematopus ostralegus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: amber (>20% of European breeding population, >20% of East Atlantic Flyway population in winter, localised wintering population)

Long-term trend

UK: rapid increase

UK population size

113,000 (98,500–127,000) pairs in 1985–99 ([O'Brien 2005](#): [BiE04](#), [APEP06](#))

Status summary

Oystercatchers increased along linear waterways between 1974 and about 1986, as the species colonised inland sites across England and Wales ([Gibbons et al. 1993](#)). Thereafter, the WBS index stabilised, so showing a pattern similar to that in winter abundance revealed by WeBS ([Banks et al. 2006](#)). Surveys in England and Wales revealed an increase of 47% in breeding birds in wet meadows between 1982 and 2002 ([Wilson et al. 2005](#)). BBS data since 1994, which include birds in a broader range of locations and habitats, show strong increase in England but apparently a significant decline in Scotland. The increase in nest failure rates during the 27-day egg stage (25 days for incubation and 2 days for laying) probably results from the spread of the species into less favourable habitats, where nest losses through predation or trampling may be more likely. The trend towards earlier laying can be partly explained by recent climate change ([Crick & Sparks 1999](#)).



Population changes

Waterways Bird Survey 1974–2007 Oystercatcher





Table of population changes for Oystercatcher

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	23	113	67	213		
	25	1981-2006	25	39	12	104		
	10	1996-2006	27	4	-9	42		
	5	2001-2006	24	0	-5	19		
BBS UK	11	1995-2006	274	-12	-22	-5		
	10	1996-2006	281	-10	-18	-3		
	5	2001-2006	309	-5	-13	1		
BBS England	11	1995-2006	137	48	26	79		
	10	1996-2006	143	44	26	67		
	5	2001-2006	169	21	9	30		
BBS Scotland	11	1995-2006	123	-22	-32	-14		
	10	1996-2006	123	-19	-29	-10		
	5	2001-2006	123	-9	-18	-1		


BBS UK 1994—2007

Oystercatcher




BBS England 1994—2007

Oystercatcher



BBS Scotland 1994—2007

Oystercatcher




Productivity trends

Table of productivity changes for Oystercatcher

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	102	None				
Daily failure rate (eggs)	38	1968-2006	112	Curvilinear	1.43% nests/day	2.72% nests/day	90.2%	
Laying date	38	1968-2006	46	Linear decline	May 17	May 10	-7 days	

Clutch size 1966—2007


Oystercatcher



Insufficient data on brood size available for this species

Egg stage nest failure rate


Oystercatcher



Insufficient data on nestling failure available for this species

Laying date 1966—2007

Oystercatcher



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

RINGED PLOVER *Charadrius hiaticula*

- Population changes
- Productivity trends
- Additional information



Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
 UK: amber (25–50% decline in winter population, >20% East Atlantic Flyway population in winter)

Long-term trend

UK: decline

UK population size

8,540 pairs in 1984 (Prater 1989: APEP06, rounded to 8,600 BiE04); 5,438 (5,257–5,622) pairs in 2007 (Conway *et al.* 2008)

Status summary

This species was already amber-listed on the strength of its concentration within UK in the winter, but a decline in winter numbers since the late 1980s (Austin *et al.* 2008) adds a further amber criterion. The breeding population is not monitored annually, but a BTO survey in 1984 showed increases throughout the UK since the previous survey in 1973–74 (Prater 1989). The spread of the breeding distribution inland between the two atlas periods, especially in England, was probably associated with the increase in number of gravel pits and reservoirs (Gibbons *et al.* 1993). The 1984 survey revealed that over 25% of the UK population nested on the Western Isles, especially on the machair, but breeding waders there have subsequently suffered greatly from predation by introduced hedgehogs (Jackson *et al.* 2004) – a problem that appears increasingly severe (Jackson 2007). Surveys in England and Wales revealed an increase of 12% in breeding birds in wet meadows between 1982 and 2002 (Wilson *et al.* 2005). The BTO's repeat national survey in 2007 found an overall decrease in UK population of around 37% since 1984, with the greatest decreases in inland areas (Burton & Conway 2008, Conway *et al.* 2008, Conway & Burton 2009; click [here](#)). Ringed Plovers that choose beaches for nesting are especially vulnerable to disturbance, however, and were in some regions in 1984 largely confined to wardened reserves (Prater 1989). Human usage of beach areas severely restricts the availability of this habitat to nesting plovers (Liley & Sutherland 2007). The marked increase in nest failures at the egg stage has earned Ringed Plover a place on the NRS concern list (Leech & Barimore 2008).

Population changes

Annual breeding population changes for this species are not currently monitored by BTO


Productivity trends

Table of productivity changes for Ringed Plover

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	87	None				
Daily failure rate (eggs)	38	1968-2006	125	Linear increase	2.36% nests/day	2.85% nests/day	20.8%	
Laying date	38	1968-2006	39	None				

Clutch size 1966–2007


Ringed Plover



Insufficient data on brood size available for this species

Egg stage nest failure rate


Ringed Plover



Insufficient data on nestling failure available for this species

Laying date 1966–2007

Ringed Plover



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GOLDEN PLOVER

Pluvialis apricaria

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK: possible decline

UK population size

22,600 pairs in 1981–84 (Reed 1985, Stroud *et al.* 1987: APEP06); 38,400–59,400 pairs in 1980–2000 (BiE04)



Status summary

The species has recently been moved from the amber to the green list because new data suggest that it does not qualify as internationally important during the breeding season. There was no annual monitoring of the breeding population before the inception of BBS. Since 1994, BBS has shown some increase in Scotland and the UK, but this is believed to follow an earlier decline (Gibbons *et al.* 1993). A detailed survey has confirmed a sharp decline in Wales since the 1980s, with just 36 pairs located in 2007 (Johnstone *et al.* 2008). Nest survival on grass moors, unlike that on heather moors, may have declined over time (Crick 1992), perhaps linked to increased stocking densities of sheep (Fuller 1996). There is no clear trend in clutch size; a large number of late-season nest records, which provide higher proportions of two- and three-egg clutches, were submitted from an intensive study during 1996–98 (J.W. Pearce-Higgins, pers. comm.). Warmer springs are reported to advance the breeding phenology of Golden Plovers and of their tipulid prey (Pearce-Higgins *et al.* 2005). Winter numbers counted by WeBS, although mainly at coastal sites and omitting some big concentrations inland, have increased sharply in Britain since the mid 1980s (Austin *et al.* 2008); these birds are mainly of Fennoscandian or Russian origin.

Population changes

BBS UK 1994–2007

Golden Plover




Table of population changes for Golden Plover


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	54	7	-12	28		
	10	1996-2006	53	9	-7	27		
	5	2001-2006	49	19	-3	45		
BBS Scotland	11	1995-2006	41	4	-16	26		
	10	1996-2006	40	7	-12	29		
	5	2001-2006	34	22	-7	53		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

BBS Scotland 1994–2007

Golden Plover




Productivity trends

Table of productivity changes for Golden Plover

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	None				Small sample

Clutch size 1966–2007

Golden Plover



Insufficient data on brood size available for this species

Insufficient data on nest failure available for this species

Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

LAPWING

Vanellus vanellus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 2, vulnerable
UK: amber (25–50% population decline, >20% European wintering population)
UK Biodiversity Action Plan: priority species

Long-term trend

UK: moderate decline

UK population size

156,000 (137,000–174,000) pairs in 1985–99 (O'Brien 2005: BiE04, APEP06)

Status summary

Although CBC recorded some increase in its early years, Lapwings have declined continuously on lowland farmland since the mid 1980s, probably because changes in agricultural practice have led to their breeding productivity dropping below a sustainable level (Galbraith 1988, Hudson *et al.* 1994, Siriwardena *et al.* 2000a, Besbeas *et al.* 2002, Milsom 2005). National surveys in England and Wales showed a 49% population decline between 1987 and 1998 (Wilson *et al.* 2001). Population declines of more than 50% over 15 years in Northern Ireland (Henderson *et al.* 2002) mirror similar declines throughout grassland areas of Wales and southeast England (Wilson *et al.* 2001, 2005). BBS data indicate shallow increase in England since 1994, but steep decline in Scotland. Adult and first-year survival rates show no trend through time (Peach *et al.* 1994, Catchpole *et al.* 1999). Mean clutch size increased significantly as the population fell. Using NRS data for 1962–99, Chamberlain & Crick (2003) found that marginal upland had relatively low reproductive performance, and arable relatively high, while grazed grass had higher failure rates and lower clutch sizes than ungrazed grass: their results suggest that recent population change may have been influenced by changes in clutch failure rates, perhaps mediated by an increase in grazing intensity in marginal uplands and by increased predation, possibly associated with habitat change. There have been several very poor years for egg-stage survival since 1996, and the species is therefore now of NRS concern (Leech & Barimore 2008). A recent study has indicated that 88% of nest predations occurred during darkness, suggesting that nocturnal mammals were to blame (Bolton *et al.* 2007). Nests with close neighbours and furthest from field edges were most likely to survive (MacDonald & Bolton 2008). Sharpe *et al.* (2008), however, conclude that chick mortality is the main determinant of poor Lapwing productivity and therefore of population decline.

The amber listing of this species is now based on UK decline, as well as the original criterion of international importance. Winter numbers counted by WeBS, mainly at coastal sites and omitting some big concentrations inland, increased in Britain during the 1980s and early 1990s and are now stable, but have decreased in Northern Ireland (Austin *et al.* 2008); these birds are mainly of continental origin. Lapwing is one of the most strongly declining bird species in Europe, having decreased at an annual rate of 4% during 1980–2005 (PECBMS 2007).



Population changes

CBC/BBS UK 1966–2007

Lapwing




Table of population changes for Lapwing


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	231	-34	-64	-5	>25	
	25	1981-2006	324	-53	-65	-37	>50	
	10	1996-2006	642	-12	-23	-3		
	5	2001-2006	685	-6	-12	3		
CBC/BBS England	39	1967-2006	190	-11	-59	29		
	25	1981-2006	266	-43	-58	-16	>25	
	10	1996-2006	529	10	0	20		
	5	2001-2006	575	3	-4	10		
WBS waterways	26	1980-2006	37	-8	-58	69		
	25	1981-2006	38	-14	-59	53		
	10	1996-2006	34	-13	-46	41		
	5	2001-2006	29	-7	-32	36		
BBS UK	11	1995-2006	613	-16	-25	-7		
	10	1996-2006	625	-13	-22	-4		
	5	2001-2006	685	-5	-13	3		
BBS England	11	1995-2006	503	7	-3	16		
	10	1996-2006	514	9	-1	17		
	5	2001-2006	575	4	-3	9		
BBS Scotland	11	1995-2006	86	-38	-50	-25	>25	
	10	1996-2006	87	-34	-47	-20	>25	
	5	2001-2006	85	-15	-30	2		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Lapwing




Waterways Bird Survey 1979—2007

Lapwing




BBS UK 1994—2007

Lapwing




BBS England 1994—2007

Lapwing



BBS Scotland 1994—2007

Lapwing




Productivity trends

Table of productivity changes for Lapwing

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	122	Linear increase	3.69 eggs	3.82 eggs	3.4%	
Daily failure rate (eggs)	38	1968-2006	133	Curvilinear	1.67% nests/day	2.45% nests/day	46.7%	
Laying date	38	1968-2006	30	None				Small sample

Clutch size 1966—2007


Lapwing



Insufficient data on brood size available for this species

Egg stage nest failure rate


Lapwing



Insufficient data on nestling failure available for this species

Laying date 1966—2007

Lapwing



Insufficient data on CES available for this species<

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

SNIPE *Gallinago gallinago*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 3 (declining)
UK: amber (>50% population decline, but data possibly unrepresentative)

Long-term trend

UK: probable decline

UK population size

59,300 (52,600–69,000) pairs in 1985–99 (O'Brien 2005: BiE04, APEP06)

Status summary

Snipe were monitored by the CBC mainly in lowland England, where numbers have fallen rapidly since the 1970s as farmland has been drained (Gibbons *et al.* 1993, Siriwardena *et al.* 2000a). The CBC index fell from the early 1970s until 1984, when the number of occupied plots became too small for further monitoring (Marchant *et al.* 1990), and the graph is not shown here. In Northern Ireland, a breeding decline of around 30% occurred between the mid 1980s and 1999 (Henderson *et al.* 2002). Surveys in England and Wales revealed a decrease of 62% in breeding birds in wet meadows between 1982 and 2002, with the remaining birds becoming highly aggregated into a tiny number of suitable sites (Wilson *et al.* 2005). Birds were more likely to persist where soils remained soft and wet; the fact that Snipe have continued to decline, despite soil conditions being improved for them at many lowland wetland reserves, suggests that other key aspects of habitat quality, such as prey abundance, are more likely to be driving the decline (Smart *et al.* 2008). The trend in the upland and moorland strongholds of the species is not fully known, but the 1988–91 atlas documented range loss widely in Wales, Northern Ireland and Scotland, as well as lowland England, and a general decrease is therefore highly probable. The BBS shows increases in England and especially in Scotland since 1994, falling back in recent seasons. Daily nest failure rates at the egg stage appear to have halved. Following declines across much of Europe during the 1990s, this previously 'secure' species is now provisionally evaluated as 'declining' (BirdLife International 2004).



Population changes

BBS UK 1994–2007 Snipe





Table of population changes for Snipe

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	132	31	17	51		
	10	1996-2006	134	29	14	47		
	5	2001-2006	147	0	-12	15		
BBS England	11	1995-2006	62	14	-8	36		
	10	1996-2006	63	15	-4	35		
	5	2001-2006	74	-5	-19	7		
BBS Scotland	11	1995-2006	53	39	21	68		
	10	1996-2006	52	35	18	61		
	5	2001-2006	52	2	-12	19		

BBS England 1994—2007



BBS Scotland 1994—2007




Productivity trends

Table of productivity changes for Snipe


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	None				Small sample
Daily failure rate (eggs)	38	1968-2006	16	Linear decline	3.3% nests/day	1.36% nests/day	-58.8%	Small sample

Clutch size 1966—2007



Insufficient data on brood size available for this species

Egg stage nest failure rate



Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WOODCOCK *Scolopax rusticola*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 3 (declining)
UK: amber (>50% population decline, but data possibly unrepresentative)

Long-term trend

UK: rapid decline

UK population size

5,400–13,700 pairs in 2000 (1988–91 Atlas estimate updated using CBC trend: [BiE04](#), [APEP06](#)); 78,000 pairs in 2003 (unpublished estimate from [GCT](#))



Status summary

The Woodcock has declined rapidly and significantly on CBC plots. Because CBC did not include many coniferous forests and was concentrated in lowland Britain, however, it is not certain how well this trend represents the whole population. Provisionally, therefore, the results warrant only an amber listing. Range contractions, that may have the same cause as the decline in abundance, were recorded concurrently with part of the CBC decline ([Gibbons *et al.* 1993](#)). Recreational disturbance, the drying out of natural woodlands, overgrazing by deer, declining woodland management, and the maturation of new plantations are possible causes of the Woodcock's decline, but there is no strong hypothesis as yet ([Fuller *et al.* 2005](#)). BBS is inefficient at recording this species, and cannot continue the index series. The first special survey aimed at monitoring the UK's breeding Woodcock took place in 2003 and has provided a baseline for future monitoring ([Fuller & Hoodless 2004](#); also, [here](#)). Annual [numbers shot](#) in the UK, which include winter visitors from declining populations in Europe, increased during the 1970s and have since been maintained around the higher level. The possible effects of hunting on breeding populations of Woodcock across Europe are little understood.

Population changes

CBC all habitats 1966–2000 Woodcock




Table of population changes for Woodcock

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC all habitats	31	1968-1999	20	-74	-88	-49	>50	Small sample
	25	1974-1999	20	-76	-88	-51	>50	Small sample
	10	1989-1999	13	-40	-62	-11	>25	Small sample
	5	1994-1999	13	-24	-44	-3		Small sample

Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

CURLEW *Numenius arquata*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 2 (declining)
 UK: amber (>20% of European breeding and winter populations)
 UK Biodiversity Action Plan: priority species

Long-term trend

England: probable decline

UK population size

107,000 (99,500–125,000) pairs in 1985–99 (O'Brien 2005: BiE04, APEP06)

Status summary

Curlews monitored by CBC were mostly in lowland habitats and may have been affected primarily by drainage of farmland (Gibbons *et al.* 1993). Surveys of breeding birds in wet meadows in England and Wales revealed a decrease of 39% between 1982 and 2002 (Wilson *et al.* 2005). A 2006 survey highlighted the rapid decline of the species across all habitats in Wales, with low breeding success as a plausible mechanism (Johnstone *et al.* 2007). In Northern Ireland, a breeding decline of around 60% occurred between the mid 1980s and 1999 (Henderson *et al.* 2002). BBS data also show that decline has been widespread. WBS data, in contrast, indicate a moderate increase during the 1980s in Curlews nesting alongside waterways. Wintering Curlew abundance showed a shallow long-term increase to around 2000, but has since declined (Austin *et al.* 2008).



Population changes

CBC/BBS England 1966–2007 Curlew




Table of population changes for Curlew


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	95	-29	-68	25		Small CBC sample
	25	1981-2006	139	-31	-66	10		Small CBC sample
	10	1996-2006	284	-19	-26	-12		
	5	2001-2006	285	-10	-16	-3		
WBS waterways	26	1980-2006	22	22	-23	85		
	25	1981-2006	22	19	-22	75		
	10	1996-2006	20	6	-14	29		
	5	2001-2006	15	3	-14	14		Small sample
BBS UK	11	1995-2006	453	-38	-43	-33	>25	
	10	1996-2006	460	-36	-40	-31	>25	
	5	2001-2006	455	-23	-28	-18		
BBS England	11	1995-2006	270	-21	-27	-12		
	10	1996-2006	275	-18	-25	-10		
	5	2001-2006	285	-9	-15	-3		
BBS Scotland	11	1995-2006	120	-50	-55	-44	>25	
	10	1996-2006	120	-47	-53	-41	>25	
	5	2001-2006	110	-31	-40	-24	>25	

BBS V	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
			5 2001-2006	37	-26	-40	-10	>25	




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


Waterways Bird Survey 1979—2007




BBS UK 1994—2007




BBS England 1994—2007



BBS Scotland 1994—2007



BBS Wales 1994—2007




Productivity trends

Table of productivity changes for Curlew


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	21	None				Small sample
Daily failure rate (eggs)	38	1968-2006	24	None				Small sample

Clutch size 1966—2007



Insufficient data on brood size available for this species

Egg stage nest failure rate



Insufficient data on nestling failure available for this species

Insufficient data on laying date
available for this species

Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

COMMON SANDPIPER *Actitis hypoleucos*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (declining)
UK: green

Long-term trend

UK: shallow decline

UK population size

12,000 pairs in 2000 (1988–91 Atlas estimate updated using WBS trend: [BiE04](#), [APEP06](#)); about 24,000 pairs in Britain ([Dougall et al. 2004](#))



Status summary

WBS results for this species show a decline from 1985 onwards (after a more gradual increase) that has yet to be explained. The recent decrease is matched by BBS data from Scotland and from the UK as a whole, and warrants a BTO alert. Poorer breeding success and reduced survival of first-year birds over the winter in West Africa were both suggested as possible reasons for the failure of the Peak District population to recover after a hard-weather event in 1989 ([Holland & Yalden 2002](#)). Following declines during the 1990s in the large Swedish and Finnish populations, and more widely in Europe, the European status of this species is no longer considered 'secure' ([BirdLife International 2004](#)). The mean change across all European countries during the 1990s was a significant decline ([Sanderson et al. 2006](#)). UK clutch sizes appear to have shown a slight decline since the 1960s.

Population changes

Waterways Bird Survey 1974–2007 Common Sandpiper






Table of population changes for Common Sandpiper

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	25	-22	-46	-3		
	25	1981-2006	26	-28	-44	-10	>25	
	10	1996-2006	21	-11	-28	5		
BBS UK	5	2001-2006	18	-1	-16	13		Small sample
	11	1995-2006	62	-19	-35	-4		
	10	1996-2006	63	-19	-34	-5		
BBS Scotland	5	2001-2006	57	-7	-23	7		
	11	1995-2006	32	-17	-32	4		
	10	1996-2006	33	-16	-33	4		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB




Productivity trends

Table of productivity changes for Common Sandpiper

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	11	Curvilinear	3.99 eggs	3.88 eggs	-2.7%	Small sample
Daily failure rate (eggs)	38	1968-2006	13	None				Small sample



Insufficient data on brood size available for this species



Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

REDSHANK *Tringa totanus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 2 (declining)
 UK: amber (>50% population decline but data possibly unrepresentative, >20% of East Atlantic Flyway population in winter)

Long-term trend

UK: moderate decline

UK population size

38,800 (31,400–44,400) pairs in 1985–99 (O'Brien 2005: BiE04, APEP06)

Status summary

UK population decline has recently been added to the criteria by which Redshank qualifies for amber listing. Considerable range contraction had occurred from many areas of the UK by 1988–91, probably as a result of the drainage of farmland (Gibbons *et al.* 1993). WBS results show a decline along waterways that apparently accelerated during the 1990s. BBS shows continuing overall decrease. Surveys in England and Wales revealed a decrease of 29% in breeding birds in wet meadows between 1982 and 2002 (Wilson *et al.* 2005). The substantial section of the British population that nests on saltmarshes decreased by 23% between 1985 and 1996, apparently as a result of increased grazing pressure (Brindley *et al.* 1998, Norris *et al.* 1998). Wintering populations (augmented by many Icelandic and some other northern European breeders) have been stable since the mid 1980s (Austin *et al.* 2008). The failure rate of nests at the egg stage has fallen steeply since the 1960s.



Population changes

Waterways Bird Survey 1974–2007 Redshank






Table of population changes for Redshank

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	17	-49	-85	-16	>25	Small sample
	25	1981-2006	17	-45	-76	-22	>25	Small sample
	10	1996-2006	13	-33	-51	-12	>25	Small sample
	5	2001-2006	10	-2	-20	11		Small sample
BBS UK	11	1995-2006	76	-22	-39	-1		
	10	1996-2006	77	-21	-37	1		
	5	2001-2006	82	-6	-26	7		
BBS England	11	1995-2006	51	-16	-28	6		
	10	1996-2006	52	-20	-32	1		
	5	2001-2006	58	-27	-38	-7	>25	




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB




Productivity trends

Table of productivity changes for Redshank

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	30	None				
Daily failure rate (eggs)	38	1968-2006	33	Linear decline	3.98% nests/day	1.69% nests/day	-57.5%	



Insufficient data on brood size available for this species



Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

FERAL PIGEON

Columba livia

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK: possible increase

UK population size

>100,000 pairs in 1968–72 (1968–72 Atlas: [APEP06](#));
100,000–250,000 pairs in 1988–91 ([BiE04](#))



Status summary

CBC samples for Feral Pigeon were consistently too small for annual monitoring, and there was no trend information before BBS began in 1994. Breeding atlas data show a 39% increase in occupied 10-km squares between 1968–72 and 1988–91 ([Gibbons et al. 1993](#)), suggesting that Feral Pigeons may be on an upward trajectory, like the other *Columba* species in the UK. At the time of the first atlas, however, Feral Pigeons were commonly excluded from bird surveys, and some of the reported subsequent range increase may have been due to greater observer awareness. It is now clear that Feral Pigeons are almost ubiquitous in the UK, nesting in rural as well as urban habitats, and avoiding only the highest ground. No distinction can realistically be drawn between birds of domestic origin and true wild-type Rock Doves, although birds of wild-type plumage may still predominate on remote Scottish islands. In field conditions, it is not usually possible to distinguish between native Rock Doves, wild-nesting Feral Pigeons, semicaptive dovecote breeders, and passing racing pigeons, nor between adults and young of the year, and BBS counts are likely to include birds from all of these groups. BBS indices suggest that a minor decrease has occurred in the last five years.

Population changes




Table of population changes for Feral Pigeon/Rock Dove


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	617	-5	-17	8		
	10	1996-2006	634	-6	-15	5		
	5	2001-2006	676	-9	-16	-1		
BBS England	11	1995-2006	517	-5	-16	6		
	10	1996-2006	531	-5	-15	5		
	5	2001-2006	562	-7	-16	-1		
BBS Scotland	11	1995-2006	55	-12	-43	32		
	10	1996-2006	56	-13	-40	24		
	5	2001-2006	58	-17	-34	5		
BBS Wales	11	1995-2006	31	43	-12	100		
	10	1996-2006	33	29	-12	70		
	5	2001-2006	37	-4	-24	19		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS England 1994—2007

Feral Pigeon/Rock Dove



BBS Scotland 1994—2007

Feral Pigeon/Rock Dove



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

STOCK DOVE
Columba oenas

- Population changes
- Productivity trends
- Additional information



Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
UK: amber (>20% of European breeding population)

Long-term trend

England: rapid increase

UK population size

309,000 territories in 2000 (1988–91 Atlas estimate updated using CBC trend: [BIE04](#), [APEP06](#))

Status summary

Following release from the lethal and sublethal effects of the organochlorine seed-dressings used in the 1950s and early 1960s, Stock Dove populations have increased very substantially ([O'Connor & Mead 1984](#)). Numbers appeared to level off in the early 1980s, and entered a further increasing phase in the early 1990s. Recent indices suggest that numbers have fallen slightly. The increase in nest failure rates at the egg stage, now reversed, was not detectable in farmland habitats alone ([Siriwardena et al. 2000b](#)). Overall, nest failure rates have fallen substantially.

Population changes

CBC/BBS England 1966–2007
Stock Dove






Table of population changes for Stock Dove

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	248	163	79	293		
	25	1981-2006	353	7	-13	32		
	10	1996-2006	673	-2	-12	11		
	5	2001-2006	703	-11	-17	-3		
BBS UK	11	1995-2006	681	2	-7	13		
	10	1996-2006	696	0	-8	10		
	5	2001-2006	764	-7	-14	0		
BBS England	11	1995-2006	628	0	-11	11		
	10	1996-2006	641	-2	-12	9		
	5	2001-2006	703	-10	-16	-2		








The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Table of productivity changes for Stock Dove

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	80	None				
Brood size	38	1968-2006	111	Curvilinear	1.82 chicks	1.84 chicks	1%	
Daily failure rate (eggs)	38	1968-2006	78	Curvilinear	1.2% nests/day	0.36% nests/day	-70%	
Daily failure rate (chicks)	38	1968-2006	57	Linear decline	1.2% nests/day	0.71% nests/day	-40.8%	
Laying date	38	1968-2006	16	None				Small sample



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WOODPIGEON *Columba palumbus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: rapid increase

UK population size

2,570,000–3,160,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

The CBC/BBS trend for this species is of a steady, steep increase since at least the mid 1970s. The spread of intensive arable cultivation, especially of oilseed rape, which has been shown to promote overwinter survival, may explain the rise in numbers ([Gibbons *et al.* 1993](#)). Since 1994, BBS has recorded significant increase in the UK, and in England, Wales and Northern Ireland separately, but stability in Scotland. [O'Connor & Shrubbs \(1986\)](#) found that the breeding season had advanced in response to the switch to autumn sowing, and thus earlier ripening, of cereals, with more pairs nesting in May and June and relatively fewer in July–September. A trend toward earlier nesting could have led CBC, with its fieldwork finishing in early July, to overestimate the rate of increase ([Marchant *et al.* 1990](#)). Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007 Woodpigeon




Table of population changes for Woodpigeon

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	724	146	22	445		
	25	1981-2006	1105	80	55	119		
	10	1996-2006	2280	27	23	32		
	5	2001-2006	2408	11	8	14		
CBC/BBS England	39	1967-2006	581	168	36	581		
	25	1981-2006	886	97	64	136		
	10	1996-2006	1817	33	29	38		
	5	2001-2006	1907	14	11	17		
BBS UK	11	1995-2006	2144	27	21	33		
	10	1996-2006	2204	26	21	31		
	5	2001-2006	2408	11	8	15		
BBS England	11	1995-2006	1709	34	27	40		
	10	1996-2006	1753	31	26	36		
	5	2001-2006	1907	14	11	18		
BBS Scotland	11	1995-2006	174	-5	-23	18		
	10	1996-2006	177	-3	-19	15		
	5	2001-2006	185	-5	-16	7		


BBS W	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
		5	2001-2006	210	17	8	25		
BBS N.Ireland		11	1995-2006	73	64	28	103		
		10	1996-2006	79	49	19	75		
		5	2001-2006	92	8	-2	20		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Woodpigeon




BBS UK 1994—2007

Woodpigeon




BBS England 1994—2007

Woodpigeon




BBS Scotland 1994—2007

Woodpigeon




BBS Wales 1994—2007

Woodpigeon



BBS N. Ireland 1994—2007

Woodpigeon



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

COLLARED DOVE
Streptopelia decaocto

- Population changes
- Productivity trends
- Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK, England: rapid increase

UK population size

298,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Collared Dove abundance has increased rapidly since the species first colonised Britain in 1955. From just four birds known to be present in that year, the population was put conservatively at 15,000–25,000 pairs by 1970 ([Hudson 1972](#)). The CBC index showed an almost exponential rise as colonisation continued during the early 1970s, but had levelled off by about 1980. BBS shows continuing increases, at least in England and Wales. The UK population size now rivals that of [Stock Dove](#). Despite the population increase, breeding productivity has also increased, perhaps as the species has become better adapted to its new environment.

Population changes

CBC/BBS UK 1971–2007
Collared Dove




Table of population changes for Collared Dove


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	34	1972-2006	469	415	236	646		
	25	1981-2006	611	50	23	90		
	10	1996-2006	1251	25	18	32		
	5	2001-2006	1354	8	3	12		
CBC/BBS England	34	1972-2006	413	440	252	666		
	25	1981-2006	539	52	18	95		
	10	1996-2006	1101	24	17	31		
	5	2001-2006	1178	7	2	11		
BBS UK	11	1995-2006	1181	29	23	37		
	10	1996-2006	1217	24	19	31		
	5	2001-2006	1354	7	3	11		
BBS England	11	1995-2006	1041	29	21	39		
	10	1996-2006	1071	24	17	32		
	5	2001-2006	1178	7	2	11		
BBS Scotland	11	1995-2006	42	-2	-28	33		
	10	1996-2006	43	4	-21	37		
	5	2001-2006	50	18	-4	41		
BBS Wales	11	1995-2006	64	49	7	92		
	10	1996-2006	67	40	7	70		
	5	2001-2006	78	15	2	30		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1971–2007

Collared Dove




BBS UK 1994–2007

Collared Dove




BBS England 1994–2007

Collared Dove




BBS Scotland 1994–2007

Collared Dove



BBS Wales 1994–2007

Collared Dove




Productivity trends

Table of productivity changes for Collared Dove

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	43	None				
Brood size	38	1968-2006	70	Linear increase	1.76 chicks	1.85 chicks	4.7%	
Daily failure rate (eggs)	38	1968-2006	62	None				
Daily failure rate (chicks)	38	1968-2006	55	Linear decline	1.82% nests/day	1.06% nests/day	-41.8%	
Laying date	38	1968-2006	44	None				


Clutch size 1966–2007

Collared Dove




Brood size 1966–2007

Collared Dove




Egg stage nest failure rate

Collared Dove




Chick stage nest failure rate

Collared Dove



Laying date 1966–2007

Collared Dove



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

TURTLE DOVE *Streptopelia turtur*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 3 (declining)

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK, England: rapid decline

UK population size

44,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))



Status summary

The CBC/BBS trend is of severe declines in Turtle Dove abundance, beginning in the late 1970s and continuing to the present. Hunting during migration is a possible cause of the UK decline, to add to those related to agricultural intensification that have been postulated for other farmland seed-eaters ([O'Connor & Shrubbs 1986](#), [Krebs et al. 1999](#)). Analysis of nest record cards and ringing data for farmland Turtle Doves suggests, although without statistical significance, that productivity per nesting attempt has increased while annual survival has fallen ([Siriwardena et al. 2000a, 2000b](#), [Browne et al. 2005](#)). [Browne & Aebischer \(2004, 2005\)](#) conclude that Turtle Doves today have a substantially earlier close to the breeding season and consequently produce barely half the number of clutches and young per pair they did in the 1960s. Thus, the recovery of Turtle Doves in Britain would benefit from the provision and sympathetic management of nesting as well as foraging habitats. Turtle Dove is one of the most strongly declining bird species in Europe, having decreased at an annual rate of 4% during 1980–2005 ([PECBMS 2007](#)). Conditions in winter may also be influencing trends: a recent study has demonstrated a positive correlation between survival rate among breeding adults in France and food supply in West Africa, as measured by cereal production ([Eraud et al. 2009](#)).

Population changes

CBC/BBS UK 1966–2007 Turtle Dove




Table of population changes for Turtle Dove


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	106	-85	-90	-76	>50	
	25	1981-2006	123	-82	-88	-75	>50	
	10	1996-2006	199	-58	-62	-51	>50	
	5	2001-2006	157	-40	-47	-31	>25	
CBC/BBS England	39	1967-2006	105	-85	-91	-76	>50	
	25	1981-2006	122	-82	-88	-74	>50	
	10	1996-2006	195	-57	-62	-51	>50	
	5	2001-2006	155	-40	-47	-30	>25	
BBS UK	11	1995-2006	182	-60	-66	-54	>50	
	10	1996-2006	184	-58	-63	-51	>50	
	5	2001-2006	157	-40	-46	-32	>25	
BBS England	11	1995-2006	179	-60	-65	-52	>50	
	10	1996-2006	181	-57	-63	-50	>50	
	5	2001-2006	155	-40	-48	-31	>25	



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Turtle Dove




BBS UK 1994—2007

Turtle Dove



BBS England 1994—2007

Turtle Dove




Productivity trends

Table of productivity changes for Turtle Dove

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	12	None				Small sample
Brood size	38	1968-2006	16	None				Small sample
Daily failure rate (eggs)	38	1968-2006	16	None				Small sample
Daily failure rate (chicks)	38	1968-2006	12	None				Small sample
Laying date	38	1968-2006	13	None				Small sample


Clutch size 1966—2007

Turtle Dove




Brood size 1966—2007

Turtle Dove




Egg stage nest failure rate

Turtle Dove




Chick stage nest failure rate

Turtle Dove



Laying date 1966—2007

Turtle Dove



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

RING-NECKED PARAKEET

Psittacula krameri

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: not evaluated (introduced)

UK: not listed (introduced)

Long-term trend

England: rapid increase

UK population size

4,300 individual adults in winter 2000/01 ([Butler 2002: APEP06](#)); further growth ([Holling & RBBP 2007b](#))



Status summary

Following escapes and releases over many decades, this African and Asian parrot began breeding annually in the UK in 1969. Substantial but highly localised self-sustaining populations of this species have since built up, with the two largest being in the southern part of Greater London and in the Isle of Thanet, east Kent. Population modelling has revealed that populations in Greater London have increased by approximately 30% per year, and those in Thanet by 15% per year, but that the range has expanded by only 0.4 km per year in the Greater London area and so far not at all in Thanet ([Butler 2003](#)). A single roost site used each night by birds from throughout the south London range held 6,818 birds in August 2003 ([Holling & RBBP 2007b](#)). The species has already been reported causing economic damage to crops, as has occurred elsewhere in its native and introduced range ([Butler 2003](#)). A recent study in Belgium has identified negative effects on breeding [Nuthatch](#), but not on other native hole-nesting species, such as [Starling](#) ([Strubbe & Matthysen 2007](#)).

Population changes

BBS England 1994–2007 Ring-necked Parakeet




Table of population changes for Ring-necked Parakeet

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS England	11	1995-2006	34	522	202	1861		
	10	1996-2006	37	424	189	1174		
	5	2001-2006	61	104	21	215		

Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch page](#)

CUCKOO

Cuculus canorus

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
 UK: amber (25–50% population decline)
 UK Biodiversity Action Plan: priority species

Long-term trend

England: rapid decline

UK population size

9,600–20,000 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

The CBC/BBS trend shows Cuckoo abundance to have been in decline since the early 1980s. The species has recently been moved from the green to the amber list, but the data now meet red-list criteria. The sensitivity of CBC to change in this species may have been relatively low, mainly because Cuckoo territories were typically larger than census plots ([Marchant et al. 1990](#)). BBS shows a continuing strong decline in England and Wales, but apparent increase in Scotland. Cuckoo numbers may have fallen because the populations of some key host species, such as [Dunnock](#) and [Meadow Pipit](#), have declined ([Brooke & Davies 1987](#)). Decreases among certain British moths may have reduced food supplies for returning adults, and the species may also be suffering difficulties on migration or in winter ([Glue 2006](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS England 1966–2007 Cuckoo








Table of population changes for Cuckoo

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	260	-61	-71	-48	>50	
	25	1981-2006	352	-61	-67	-54	>50	
	10	1996-2006	614	-46	-49	-43	>25	
	5	2001-2006	525	-20	-23	-14		
BBS UK	11	1995-2006	729	-33	-38	-27	>25	
	10	1996-2006	731	-29	-35	-24	>25	
	5	2001-2006	672	-9	-15	-4		
BBS England	11	1995-2006	584	-50	-53	-46	>25	
	10	1996-2006	582	-46	-49	-42	>25	
	5	2001-2006	525	-20	-24	-16		
BBS Scotland	11	1995-2006	66	16	-2	33		
	10	1996-2006	66	11	-6	29		
	5	2001-2006	62	12	-3	28		
BBS Wales	11	1995-2006	57	-33	-47	-19	>25	
	10	1996-2006	59	-32	-46	-21	>25	
	5	2001-2006	59	-19	-30	-6		



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

BARN OWL *Tyto alba*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (declining)
UK: amber (25–50% distribution decline)

Long-term trend

UK: decline

UK population size

4,000 (3,000–5,000) pairs in 1995–97 (Toms *et al.* 2001; BiE04, APEP06)

Status summary

Distributional data provide good evidence for a decline in this species that lasted throughout the 20th century, although annual monitoring started only very recently. Productivity has tended to improve since the 1950s and 1960s when Barn Owls appear to have been affected by organochlorine pesticides (Percival 1990). A national census during 1995–97, organised jointly by Hawk & Owl Trust and BTO, provided a replicable baseline population estimate (Toms *et al.* 2000, 2001; for more information, click [here](#)). The lack of annual population change data for this species is now being addressed by the BTO's [Barn Owl Monitoring Programme \(BOMP\)](#), which began in 2000; additional nest record, ringing and biometric information is also being collected through this scheme (Leech *et al.* 2005). BOMP already provides evidence that fewer pairs attempt to nest following cold or wet winters (Leech *et al.* 2006a). In earlier decades, the plight of such a charismatic and popular bird led to extensive releasing of captive-bred birds in well-meaning attempts at restocking: by 1992, when licensing became a requirement for such schemes, it was estimated that between 2,000 and 3,000 birds were being released annually by about 600 operators, although many birds died quickly and few would have joined the nesting population (Balmer *et al.* 2000). More recently, the erection of Barn Owl nest boxes, already numbering c. 25,000 by the mid 1990s, has enabled the species to occupy areas (notably the Fens) that were previously devoid of nesting sites, and may have been a factor in improving nesting success. RBBP provide a county breakdown of 2005 nesting totals [here](#) (Holling & RBBP 2008).



Population changes


Annual breeding population changes for this species are not currently monitored by BTO

Productivity trends


Table of productivity changes for Barn Owl

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	22	Linear increase	4.53 eggs	5.15 eggs	13.8%	Small sample
Brood size	38	1968-2006	149	Curvilinear	3.01 chicks	3.12 chicks	3.4%	
Daily failure rate (eggs)	38	1968-2006	18	Linear decline	0.75% nests/day	0.1% nests/day	-86.7%	Small sample
Daily failure rate (chicks)	38	1968-2006	69	Linear decline	0.21% nests/day	0.03% nests/day	-85.7%	


Clutch size 1966–2007
Barn Owl




Brood size 1966–2007
Barn Owl



Egg stage nest failure rate
Barn Owl



Chick stage nest failure rate
Barn Owl



Insufficient data on laying date

Insufficient data on CES

available for this species

available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

LITTLE OWL *Athene noctua*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3, declining
UK: not listed (introduced)

Long-term trend

UK, England: probable decline

UK population size

5,800–11,600 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary



The CBC/BBS trend for Little Owl in the UK shows very wide variation, but a downturn in recent seasons suggests that a moderate long-term decline may lie behind the observed fluctuations. Trends are poorly known, however, because the species has large breeding territories and, being largely inactive during the day, is difficult to detect except by dedicated surveys. A population estimate of c. 7,000 pairs from the BTO/Hawk & Owl Trust's [Project Barn Owl \(Toms et al. 2000\)](#) is the first replicable estimate for Little Owls in the UK. No trends are evident in productivity, but few nest records are available.

Population changes

CBC/BBS UK 1966–2007 Little Owl




Table of population changes for Little Owl


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	54	-46	-68	-12	>25	
	25	1981-2006	69	-46	-63	-29	>25	
	10	1996-2006	113	-20	-33	-7		
	5	2001-2006	101	-25	-39	-13	>25	
CBC/BBS England	39	1967-2006	51	-36	-61	14		
	25	1981-2006	66	-37	-55	-8	>25	
	10	1996-2006	109	-18	-31	-2		
	5	2001-2006	99	-22	-36	-7		
BBS UK	11	1995-2006	98	-17	-31	-1		
	10	1996-2006	101	-20	-33	-7		
	5	2001-2006	101	-24	-36	-9		
BBS England	11	1995-2006	95	-14	-29	2		
	10	1996-2006	97	-17	-32	-4		
	5	2001-2006	99	-20	-34	-7		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Little Owl




BBS UK 1994—2007

Little Owl



BBS England 1994—2007

Little Owl




Productivity trends

Table of productivity changes for Little Owl

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	18	Linear increase	3.39 eggs	3.62 eggs	6.7%	Small sample
Brood size	38	1968-2006	38	None				
Daily failure rate (eggs)	38	1968-2006	16	None				Small sample
Daily failure rate (chicks)	38	1968-2006	19	None				Small sample


Clutch size 1966—2007

Little Owl




Brood size 1966—2007

Little Owl




Egg stage nest failure rate

Little Owl



Chick stage nest failure rate

Little Owl



Insufficient data on laying dates available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

TAWNY OWL

Strix aluco

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: stable

UK population size

19,400 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

As a nocturnal species, Tawny Owl is covered relatively poorly by the BTO's monitoring schemes. The pattern shown by CBC/BBS is a relatively stable one, however, in keeping with the longevity, sedentary behaviour, and slow breeding rate of this species. There has been a shallow downward trend in the index since the early 1970s. It may be relevant to this possible long-term decline that [Gibbons *et al.* \(1993\)](#) found evidence for a contraction of the species' UK range between the two atlas periods. The substantial improvements in nest success during the c.29-day egg stage could be linked to the declining impact of organochlorine pesticides, which were banned in the early 1960s. Special post-breeding surveys of this species were conducted in autumn 2005 ([click here](#)), following methodology established by an earlier survey in 1989 ([Percival 1990](#)).



Population changes

CBC/BBS UK 1966–2007

Tawny Owl




Table of population changes for Tawny Owl


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	75	-2	-40	42		
	25	1981-2006	87	-25	-43	-2		
	10	1996-2006	110	-15	-29	5		
	5	2001-2006	82	-6	-21	16		
CBC/BBS England	39	1967-2006	64	-1	-41	48		
	25	1981-2006	74	-19	-39	12		
	10	1996-2006	95	-9	-24	14		
	5	2001-2006	73	-10	-23	10		
BBS UK	11	1995-2006	82	-12	-27	11		Nocturnal species
	10	1996-2006	84	-12	-25	10		Nocturnal species
	5	2001-2006	82	-7	-22	11		Nocturnal species
BBS England	11	1995-2006	70	-5	-24	24		Nocturnal species
	10	1996-2006	72	-6	-24	19		Nocturnal species
	5	2001-2006	73	-10	-25	8		Nocturnal species



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Tawny Owl




BBS UK 1994—2007

Tawny Owl



BBS England 1994—2007

Tawny Owl




Productivity trends

Table of productivity changes for Tawny Owl

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	78	None				
Brood size	38	1968-2006	139	None				
Daily failure rate (eggs)	38	1968-2006	55	Linear decline	0.95% nests/day	0.18% nests/day	-81.1%	
Daily failure rate (chicks)	38	1968-2006	82	Curvilinear	0.31% nests/day	0.11% nests/day	-64.5%	
Laying date	38	1968-2006	14	Linear decline	Mar 29	Mar 22	-7 days	Small sample


Clutch size 1966—2007

Tawny Owl




Brood size 1966—2007

Tawny Owl




Egg stage nest failure rate

Tawny Owl




Chick stage nest failure rate

Tawny Owl



Laying date 1966—2007

Tawny Owl



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

NIGHTJAR *Caprimulgus europaeus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 2, (declining)
UK: red (>50% distribution decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK: uncertain

UK population size

3,400 males in 1992 ([Morris et al. 1994: BiE04, APEP06](#)); 4,600 males in 2004 ([Conway et al. 2007](#))

Status summary

Following a catastrophic decline in range of more than 50% of 10-km squares between breeding atlases, the 1992 national survey revealed a welcome increase of 50% in population size since 1981, probably due to the increased availability of young forest habitat as plantations were felled and replanted ([Morris et al. 1994](#)). A [National Nightjar Survey](#) in 2004 revealed that a further 36% increase had taken place in the UK population in 12 years, with a 2.6% increase in the number of 10-km squares occupied ([Conway et al. 2007](#)). There was evidence of population declines and range contractions since 1992, however, in North Wales, northwest England, and Scotland. Although annual nest record sample are very small, the increases in nest failure rates and decreases in clutch and brood sizes have resulted in the inclusion of Nightjar on the NRS concern list ([Leech & Barimore 2008](#)). A recent study suggests that nest failure is most likely in areas heavily frequented by walkers and dogs ([Langston et al. 2007](#)).



Population changes


Annual population changes are not monitored for this species

Productivity trends


Table of productivity changes for Nightjar

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	17	Linear decline	2.02 eggs	1.82 eggs	-9.8%	Small sample
Brood size	38	1968-2006	24	Curvilinear	1.82 chicks	1.7 chicks	-6.2%	Small sample
Daily failure rate (eggs)	38	1968-2006	22	Linear increase	1.38% nests/day	3.55% nests/day	157.2%	Small sample
Daily failure rate (chicks)	38	1968-2006	21	Curvilinear	0.05% nests/day	0.67% nests/day	1240%	Small sample
Laying date	38	1968-2006	18	Curvilinear	Jun 17	Jun 16	-1 days	Small sample


Clutch size 1966–2007
Nightjar




Brood size 1966–2007
Nightjar



Egg stage nest failure rate
Nightjar




Chick stage nest failure rate
Nightjar



Insufficient data on CES available for this species

Laying date 1966–2007

Nightjar



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

COMMON SWIFT

Apus apus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK: unknown

UK population size

85,000 pairs in 1990 (1988–91 Atlas: [APEP06](#)); 20,000–100,000 pairs in 2000 ([BiE04](#))

Status summary



Swifts were not monitored before the inception of the BBS. Their monitoring is complicated by the difficulty of finding occupied nests, by the weather-dependent and sometimes extraordinary distances from the nest at which breeding adults may forage, and by the variable midsummer influx of non-breeding individuals. Since Swifts do not normally begin breeding until they are four years old, non-breeding numbers can at times be substantial. BBS results suggest steep declines in England, Scotland and Wales but the relationship between BBS transect counts and nesting numbers is not properly understood so far. [Concern for Swifts](#), a small private organisation, is trying to promote the deliberate provision of nesting sites for this species, as so many suitable cavities are being lost to re-development. It is also gathering information on populations to try to determine whether the species warrants a conservation concern listing.

Population changes

BBS UK 1994–2007

Swift





Table of population changes for Swift

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	961	-26	-33	-18	>25	
	10	1996-2006	980	-23	-29	-15		
	5	2001-2006	1053	-13	-20	-3		
BBS England	11	1995-2006	825	-28	-36	-18	>25	
	10	1996-2006	840	-25	-32	-19	>25	
	5	2001-2006	893	-16	-24	-8		
BBS Scotland	11	1995-2006	46	-42	-59	-18	>25	
	10	1996-2006	48	-33	-53	-13	>25	
	5	2001-2006	54	-2	-20	20		
BBS Wales	11	1995-2006	66	-31	-48	2		
	10	1996-2006	68	-30	-48	-4	>25	
	5	2001-2006	77	-26	-41	-11	>25	




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS England 1994–2007



BBS Scotland 1994–2007



BBS Wales 1994–2007



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

KINGFISHER

Alcedo atthis

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3, depleted
UK: amber (European status)

Long-term trend

UK: fluctuating, with no long-term trend

UK population size

4,800–8,000 pairs in 2000 (1988–91 Atlas estimate updated using WBS trend: [BiE04](#), [APEP06](#))

Status summary

The Kingfisher declined along linear waterways (its principal habitat) until the mid 1980s, since when it seems to have made a complete recovery. The decline was associated with a contraction of range in England ([Gibbons *et al.* 1993](#)). Kingfishers suffer severe mortality during harsh winters but, with up to three broods in a season, and up to six chicks in a brood, their potential for rapid population growth is unusually high. Amber listing of this species in the UK results from its 'depleted' status in Europe as a whole, following declines between 1970 and 1990 ([BirdLife International 2004](#)).



Population changes

Waterways Bird Survey 1974–2007 Kingfisher






Table of population changes for Kingfisher

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	32	7	-34	84		
	25	1981-2006	32	66	18	170		
	10	1996-2006	35	22	1	86		
	5	2001-2006	33	14	-1	44		
BBS UK	11	1995-2006	51	14	-13	44		
	10	1996-2006	52	21	-8	47		
	5	2001-2006	62	2	-22	26		
BBS England	11	1995-2006	44	17	-11	58		
	10	1996-2006	45	24	-5	58		
	5	2001-2006	54	15	-9	40		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GREEN WOODPECKER

Picus viridis

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 2 (depleted)

UK: amber (European status)

Long-term trend

England: rapid increase

UK population size

24,200 pairs in 2000 (1988–91 Atlas estimate updated using

CBC trend: [BiE04](#), [APEP06](#))

Status summary



Green Woodpecker populations have risen steadily in Britain since 1966, except for a period of stability or shallow decline centred around 1980 that was probably the result of a series of harsh winters. There was considerable range expansion in central and eastern Scotland between the 1968–72 and 1988–91 atlas periods. Recent results indicate that the current phase of increase is continuing across most of the UK range. The ecological factors underlying the increase are not yet known but, given the species' susceptibility to cold weather, it may be related to climate change. Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).

Population changes

CBC/BBS England 1966–2007 Green Woodpecker





Table of population changes for Green Woodpecker

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	252	205	133	327		
	25	1981-2006	360	95	61	133		
	10	1996-2006	710	46	37	54		
	5	2001-2006	761	6	1	11		
BBS UK	11	1995-2006	688	43	31	55		
	10	1996-2006	716	37	28	46		
	5	2001-2006	820	8	3	13		
BBS England	11	1995-2006	636	51	42	62		
	10	1996-2006	663	43	35	51		
	5	2001-2006	761	7	3	11		
BBS Wales	11	1995-2006	45	4	-18	30		
	10	1996-2006	47	2	-19	23		
	5	2001-2006	51	11	-5	32		


BBS UK 1994—2007

Green Woodpecker




BBS England 1994—2007

Green Woodpecker



BBS Wales 1994—2007

Green Woodpecker



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GREAT SPOTTED WOODPECKER
Dendrocopos major

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK, England: rapid increase

UK population size

37,000–44,400 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

This species increased rapidly in the 1970s and began a further increase in the early 1990s. Dutch Elm Disease, which greatly increased the amount of standing dead timber and its associated insects, has been linked to the increase that occurred during the 1970s ([Marchant et al. 1990](#)). The ecological factors underlying the current increase are not yet known, but the species may be benefiting from the maturation of new forests and from the increasing provision of winter food in gardens. The decline in [Starling](#) numbers in recent decades has led to increased breeding success of this woodpecker and may have allowed it to expand its breeding distribution into more open, less wooded habitats ([Smith 2005, 2006](#)). Nesting phenology in Hertfordshire woodlands has advanced over the last two decades in response to warmer spring weather ([Smith 2006](#)). Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007
Great Spotted Woodpecker




Table of population changes for Great Spotted Woodpecker

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	322	349	233	603		
	25	1981-2006	460	119	88	156		
	10	1996-2006	904	97	86	109		
	5	2001-2006	1036	44	37	52		
CBC/BBS England	39	1967-2006	289	332	211	556		
	25	1981-2006	412	115	85	157		
	10	1996-2006	805	91	81	102		
	5	2001-2006	916	38	31	44		
BBS UK	11	1995-2006	811	112	100	124		
	10	1996-2006	846	98	87	108		
	5	2001-2006	1036	43	37	49		
BBS England	11	1995-2006	721	105	94	119		
	10	1996-2006	752	92	81	104		
	5	2001-2006	916	38	31	42		
BBS Wales	11	1995-2006	61	98	62	148		
	10	1996-2006	63	89	59	132		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Great Spotted Woodpecker




BBS UK 1994—2007

Great Spotted Woodpecker




BBS England 1994—2007

Great Spotted Woodpecker



BBS Wales 1994—2007

Great Spotted Woodpecker



Productivity trends


Table of productivity changes for Great Spotted Woodpecker

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size	38	1968-2006	24	None				Small sample
Daily failure rate (chicks)	38	1968-2006	31	Linear decline	0.37% nests/day	0.02% nests/day	-94.6%	

>Insufficient data on clutch size available for this species

Brood size 1966—2007


Great Spotted Woodpecker



Insufficient data on egg nest failure available for this species

Chick stage nest failure rate

Great Spotted Woodpecker



Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

LESSER SPOTTED WOODPECKER *Dendrocopos minor*

• Population changes • Productivity trends • Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: red (>50% population decline)
UK Biodiversity Action Plan: priority species

Long-term trend

UK: rapid decline

UK population size

1,400–2,900 pairs in 2000 (1988–91 Atlas estimate updated using CBC trend: [BIE04](#), [APEP06](#))

Status summary

The Lesser Spotted Woodpecker has declined significantly and very rapidly since around 1980, following a shallower increase; it had already contracted in range between the two atlas periods ([Gibbons et al. 1993](#)), and has subsequently disappeared from many more of its former localities. It has become so rare that BBS observers have been unable to continue the annual monitoring that was possible until 2000 through CBC. The species qualifies easily for red listing. Competition with and predation by Great Spotted Woodpeckers, and reductions in small-diameter dead wood suitable for foraging, are the most likely causes of decline, while the species' large home ranges suggest that landscape-scale changes in woodland (loss of mature broadleaved woodland, losses of non-woodland trees such as elms, and woodland fragmentation) may also be important ([Fuller et al. 2005](#)). Lesser Spotted Woodpecker is one of the most strongly declining bird species in Europe, having decreased at an annual rate of 7% during 1980–2005 ([PECBMS 2007](#)).

Population changes

CBC all habitats 1966–2000 Lesser Spotted Woodpecker




Table of population changes for Lesser Spotted Woodpecker

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC all habitats	31	1968-1999	17	-60	-81	40		Small sample
	25	1974-1999	18	-73	-86	-31	>50	Small sample
	10	1989-1999	11	-51	-75	-22	>50	Small sample
	5	1994-1999	9	-33	-56	0		Small sample



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WOODLARK *Lullula arborea*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 2 (depleted)

UK: red (>50% distribution decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK: increase

UK population size

1,426–1,552 pairs in 1997 ([Wotton & Gillings 2000](#): APEP06, rounded to 1,400–1,600 BIE04)

Status summary

This species is too rare and restricted in range for population changes to be monitored annually by BTO observers. A 62% reduction in the number of 10-km squares occupied between 1968–72 and 1988–91 warranted red-listing on grounds of range contraction; the species had ceased to breed in Wales and in several southern English counties over this period ([Gibbons *et al.* 1993](#)). [Sitters *et al.* \(1996\)](#) report that the UK population increased from c.250 pairs in 1986 to c.600 pairs in 1993, probably helped by recent mild winters and increased habitat availability due to storm damage in plantations, forest restocking, and heathland management. A repeat national survey in 1997 showed that the population had increased further, accompanied by expansion of the range into new areas ([Wotton & Gillings 2000](#); for more information, [click here](#)). Farmland setaside, especially close to forest, is valuable additional habitat for the expanding population, although clutch sizes may be lower there than in more traditional habitats ([Wright *et al.* 2007](#)). The small NRS sample suggests that nest failure rates have become less frequent at the egg stage. Human disturbance at heathland sites apparently reduces population density, but the effects are partly offset by higher breeding productivity at lower densities ([Mallord *et al.* 2007](#)). BTO conducted a new national survey in spring 2006 (for more information, [click here](#)).



Population changes

Annual breeding population changes for this species are not currently monitored by BTO


Productivity trends

Table of productivity changes for Woodlark

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	16	None				Small sample
Brood size	38	1968-2006	27	None				Small sample
Daily failure rate (eggs)	38	1968-2006	20	Curvilinear	6.83% nests/day	2.75% nests/day	-59.7%	Small sample
Daily failure rate (chicks)	38	1968-2006	28	None				Small sample
Laying date	38	1968-2006	18	None				Small sample


Clutch size 1966–2007

Woodlark




Brood size 1966–2007

Woodlark




Egg stage nest failure rate

Woodlark



Chick stage nest failure rate


Woodlark



Insufficient data on CES available for this species

Laying date 1966—2007

Woodlark



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

SKYLARK *Alauda arvensis*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 3 (depleted)
 UK: red (>50% population decline)
 UK Biodiversity Action Plan: [click here](#)

Long-term trend

England: rapid decline

UK population size

1,785,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#));
 801,000–1,003,000 pairs in Britain in 1997 ([Browne et al. 2000](#))



Status summary

The Skylark declined rapidly from the mid 1970s until the mid 1980s, when the rate of decline slowed; more recent data show further decline, however, at least in England. Considerable effort by BTO and other researchers in recent years has indicated that the most likely cause of the decline is the change to autumn sowing of cereals: this practice restricts opportunities for late-season nesting attempts, because the crop is by then too tall, and may depress overwinter survival by reducing the area of stubbles ([Wilson et al. 1997](#), [Donald & Vickery 2000, 2001](#); for more information, [click here](#)). [Chamberlain & Siriwardena \(2000\)](#) have provided a general review of the effects of agricultural practice on Skylark population trends. More recently, [Gillings et al. \(2005\)](#) have identified better population performance in areas with extensive winter stubble, presumably because overwinter survival is relatively high. Breeding success per nesting attempt increased during the decline ([Chamberlain & Crick 1999](#), [Siriwardena et al. 2000b](#)) but, since 2000, nest losses have apparently increased and previous gains in clutch and brood sizes have been lost. Leaving small, rectangular patches of bare ground ('[Skylark plots](#)') within autumn-sown cereals appears to provide many of the benefits of spring-sown cereals at very low cost to the farmer ([Donald & Morris 2005](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).

Population changes

CBC/BBS England 1966–2007 Skylark




Table of population changes for Skylark

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	455	-59	-66	-51	>50	
	25	1981-2006	644	-47	-54	-40	>25	
	10	1996-2006	1278	-13	-17	-9		
	5	2001-2006	1307	-1	-5	2		
BBS UK	11	1995-2006	1538	-11	-16	-6		
	10	1996-2006	1568	-8	-14	-4		
	5	2001-2006	1643	-2	-6	2		
BBS England	11	1995-2006	1211	-16	-19	-12		
	10	1996-2006	1236	-12	-16	-8		
	5	2001-2006	1307	-1	-5	2		
BBS Scotland	11	1995-2006	192	1	-11	13		
	10	1996-2006	192	2	-10	14		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Wales	10	1996-2006	103	-5	-16	5		
	5	2001-2006	111	-3	-18	9		
BBS N.Ireland	11	1995-2006	34	-30	-44	-16	>25	
	10	1996-2006	36	-35	-48	-24	>25	
	5	2001-2006	36	-38	-52	-33	>25	



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS UK 1994—2007

Skylark




BBS England 1994—2007

Skylark




BBS Scotland 1994—2007

Skylark



BBS Wales 1994—2007

Skylark




Productivity trends

Table of productivity changes for Skylark

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	38	Linear increase	3.37 eggs	3.68 eggs	9%	
Brood size	38	1968-2006	67	Curvilinear	3.1 chicks	3.29 chicks	6.3%	
Daily failure rate (eggs)	38	1968-2006	47	None				
Daily failure rate (chicks)	38	1968-2006	55	None				
Laying date	38	1968-2006	20	Curvilinear	May 25	Jun 1	7 days	Small sample


Clutch size 1966—2007

Skylark




Brood size 1966—2007

Skylark




Egg stage nest failure rate

Skylark



Chick stage nest failure rate


Skylark



Insufficient data on CES available for this species

Laying date 1966—2007

Skylark



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

SAND MARTIN

Riparia riparia

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (depleted)

UK: amber (European status)

Long-term trend

UK: fluctuating, with no long-term trend

UK population size

85,000–270,000 nests in 1990 (1988–91 Atlas: [APEP06](#)); 66,300–211,000 pairs in 2000 (updated using WBS trend: [BiE04](#))

Status summary

This species is unusually difficult to monitor, because active and inactive nest holes are difficult to distinguish, and because whole colonies frequently disperse or shift to new locations as suitable sand cliffs are created and destroyed. WBS counts, which are of apparently occupied nest holes along riverbanks, suggest a stable or shallowly increasing population, with wide fluctuations, although the ongoing decrease since the late 1990s has been steep enough to raise BTO alerts. BBS counts, which are of birds seen, show clearly that large year-to-year changes occur, but do not yet reveal a clear long-term trend. Nest record samples are small, but indicate that nest success has improved enormously since the 1960s, and that clutch size has also increased. Rainfall in the species' trans-Saharan wintering grounds prior to the birds' arrival promotes annual survival and thus abundance in the following breeding season ([Szép 1995](#)). Annual survival rates from RAS sites in the UK for 1990–2004 were correlated positively with minimum monthly rainfall during the wet season in West Africa ([Robinson *et al.* 2008](#)). More recently, it has been discovered that summer rainfall on the breeding grounds has a negative influence on survival rates through the following winter ([Cowley & Siriwardena 2005](#)).



Population changes

Waterways Bird Survey 1977–2007 Sand Martin






Table of population changes for Sand Martin

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	28	1978-2006	19	-2	-39	55		Small sample
	25	1981-2006	20	-6	-42	76		
	10	1996-2006	22	-44	-59	-25	>25	
	5	2001-2006	18	-16	-27	4		Small sample
BBS UK	11	1995-2006	114	21	-36	122		
	10	1996-2006	117	16	-30	96		
	5	2001-2006	124	30	2	66		
BBS England	11	1995-2006	76	20	-19	53		
	10	1996-2006	77	10	-22	32		
	5	2001-2006	82	12	-9	36		








The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Table of productivity changes for Sand Martin

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	32	Curvilinear	4.68 eggs	5.06 eggs	8.2%	
Brood size	38	1968-2006	35	None				
Daily failure rate (eggs)	38	1968-2006	24	Linear decline	1.27% nests/day	0.01% nests/day	-99.2%	Small sample
Daily failure rate (chicks)	38	1968-2006	36	Linear decline	1.57% nests/day	0.08% nests/day	-94.9%	
Laying date	38	1968-2006	30	None				Small sample



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

SWALLOW

Hirundo rustica

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (depleted)
UK: amber (European status)

Long-term trend

UK, England: fluctuating, with no long-term trend

UK population size

726,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Swallow was originally amber-listed partly on the strength of a perceived CBC decline, but continues to qualify through its widespread decline across the European continent ([BirdLife International 2004](#)). Modern methods of estimating population change from CBC give evidence of fluctuations but not for long-term decline in the UK ([Robinson et al. 2003](#)). BBS data suggest increases throughout the UK since 1994. Analysis has shown that the population fluctuations are most strongly related to variable losses on their wintering grounds ([Baillie & Peach 1992](#)). Population change has been shown to be correlated with rainfall in the western Sahel prior to the birds' spring passage through West Africa, but with neither cattle numbers nor nest-site availability in the UK ([Robinson et al. 2003](#)). Annual survival rates from RAS sites in the UK for 1998–2004 were correlated positively with mean monthly rainfall during the early austral summer in southern Africa ([Robinson et al. 2008](#)). It is likely that, in eastern parts of the UK, the loss of livestock farming and grazed grassland, together with arable intensification, has caused the Swallow population to decline, while an increase in the area of pasture in the west and north has promoted a population increase which apparently has more than compensated for declines elsewhere ([Evans & Robinson 2004](#)). A link between regional changes in the availability of preferred feeding habitats and the regional patterns of UK population change again suggests that habitat change on the breeding grounds may explain population trend, at least partly ([Henderson et al. 2007](#)). Brood sizes increased up to the late 1980s, and may now be falling again. Climatic warming is leading to both an earlier start and later finish to the breeding season for European Swallows, but there has been increased chick mortality in hot, dry summers and reduced post-fledging survival because of poor conditions for birds migrating through North Africa ([Turner 2009](#)).



Population changes

CBC/BBS England 1966–2007 Swallow




Table of population changes for Swallow

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	439	23	-12	75		
	25	1981-2006	649	19	-4	46		
	10	1996-2006	1352	27	19	35		
	5	2001-2006	1444	17	11	24		
BBS UK	11	1995-2006	1679	32	26	39		
	10	1996-2006	1736	28	22	34		
	5	2001-2006	1901	13	9	17		
BBS England	11	1995-2006	1285	28	22	37		
	10	1996-2006	1323	25	19	33		
	5	2001-2006	1444	16	10	21		


BBS Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	160	11	1	27		
BBS Wales	11	1995-2006	158	49	22	86		
	10	1996-2006	166	36	14	63		
	5	2001-2006	189	12	1	24		
BBS N.Ireland	11	1995-2006	75	39	8	74		
	10	1996-2006	81	33	8	56		
	5	2001-2006	93	7	-8	19		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS UK 1994—2007

Swallow




BBS England 1994—2007

Swallow




BBS Scotland 1994—2007

Swallow




BBS Wales 1994—2007

Swallow



BBS N. Ireland 1994—2007

Swallow




Productivity trends

Table of productivity changes for Swallow

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	220	Curvilinear	4.46 eggs	4.51 eggs	1.2%	
Brood size	38	1968-2006	378	Curvilinear	4.07 chicks	4.12 chicks	1.3%	
Daily failure rate (eggs)	38	1968-2006	280	None				
Daily failure rate (chicks)	38	1968-2006	251	Linear increase	0.28% nests/day	0.53% nests/day	89.3%	
Laying date	38	1968-2006	105	Curvilinear	Jun 20	Jun 12	-8 days	


Clutch size 1966—2007

Swallow




Brood size 1966—2007

Swallow




Egg stage nest failure rate

Swallow




Chick stage nest failure rate

Swallow



Laying date 1966—2007

Swallow



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

HOUSE MARTIN

Delichon urbicum

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (declining)
UK: amber (25–50% population decline)

Long-term trend

UK: probable decline

UK population size

273,000–535,000 pairs in 2000 (1988–91 Atlas estimate updated using CBC trend: [BIE04](#), [APEP06](#))

Status summary

The House Martin's loosely colonial habits and strong association with human settlements mean that it is extraordinarily difficult to monitor. Anecdotal evidence of decline is often unreliable, because demise of a colony may be balanced by single nests or small groups becoming established elsewhere. For these reasons, study areas should be large, covered thoroughly, and ideally randomly selected. The available long-term data suggest a rapid decline, although BBS shows overall increase since 1994. The species has been moved from the green to the amber list, because of moderate decline in the CBC trend for 1974–99, and is newly listed as of European concern following declines elsewhere in Europe ([BirdLife International 2004](#)). The mean change across all European countries during the 1990s was a significant decline ([Sanderson *et al.* 2006](#)). Annual survival rates from RAS sites in the UK for 1994–2004 were correlated positively with maximum monthly rainfall in West Africa; some decline in survival rate is apparent over this period but does not correspond to the population decline ([Robinson *et al.* 2008](#)).



Population changes

CBC/BBS England 1966–2007

House Martin







Table of population changes for House Martin

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	218	-56	-86	42		Small CBC sample
	25	1981-2006	329	-60	-88	105		Small CBC sample
	10	1996-2006	701	5	-4	14		
	5	2001-2006	757	3	-4	10		
BBS UK	11	1995-2006	858	12	4	24		
	10	1996-2006	882	9	2	19		
	5	2001-2006	966	-3	-9	3		
BBS England	11	1995-2006	676	4	-4	15		
	10	1996-2006	692	5	-2	15		
	5	2001-2006	757	3	-4	11		
BBS Scotland	11	1995-2006	53	84	21	136		
	10	1996-2006	54	45	-9	73		
	5	2001-2006	60	-21	-42	-2		
BBS Wales	11	1995-2006	86	25	-12	63		
	10	1996-2006	89	14	-15	48		
	5	2001-2006	95	-9	-28	13		

BBS No	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
		5	2001-2006	47	-4	-23	16		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

TREE PIPIT *Anthus trivialis*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
 UK: amber (>50% population decline but data possibly unrepresentative)
 UK Biodiversity Action Plan: priority species

Long-term trend

England: rapid decline

UK population size

74,400 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

The species was moved from the green to the amber list in 2002, on the strength of its population decline. Tree Pipits occur in greatest abundance in Wales, northern England and Scotland, and thus the marked CBC decline between the two atlas periods may reflect the range contraction that occurred then in central and southeast England ([Gibbons et al. 1993](#)). Since 1994, CBC/BBS data have shown further severe decrease, especially in England. The causes of the population decline are unclear, but may be linked to changing forest structure, as new plantations mature, and reduced management of lowland woods ([Fuller et al. 2005](#)). In Thetford Forest, Tree Pipits prefer large blocks of habitat and benefit from targeted management such as the retention of mature trees for use as songposts ([Burton 2007](#)). There has been an increase in brood size and a substantial decline in failure rates over the 17-day egg stage; the species is on the NRS concern list, however, because of an overall decrease in nest survival ([Leech & Barimore 2008](#)). Although the species has no European conservation listing as yet, numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)), and the mean change across all European countries during the 1990s was a significant decline ([Sanderson et al. 2006](#)).



Population changes

CBC/BBS England 1966–2007 Tree Pipit





Table of population changes for Tree Pipit

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	44	-83	-92	-71	>50	
	25	1981-2006	48	-82	-91	-69	>50	
	10	1996-2006	73	-30	-47	-7	>25	
	5	2001-2006	64	-21	-36	-3		
BBS UK	11	1995-2006	124	-14	-34	7		
	10	1996-2006	125	-17	-36	3		
	5	2001-2006	117	-22	-39	-7		
BBS England	11	1995-2006	67	-35	-51	-11	>25	
	10	1996-2006	67	-30	-47	-10	>25	
	5	2001-2006	64	-20	-35	1		
BBS Wales	11	1995-2006	32	-22	-55	17		
	10	1996-2006	33	-27	-56	8		
	5	2001-2006	32	-22	-43	-5		


BBS UK 1994—2007

Tree Pipit




BBS England 1994—2007

Tree Pipit



BBS Wales 1994—2007

Tree Pipit




Productivity trends

Table of productivity changes for Tree Pipit

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	10	None				Small sample
Brood size	38	1968-2006	28	Linear increase	4.38 chicks	4.74 chicks	8.2%	Small sample
Daily failure rate (eggs)	38	1968-2006	12	Curvilinear	4.87% nests/day	3.12% nests/day	-35.9%	Small sample
Daily failure rate (chicks)	38	1968-2006	19	Curvilinear	3.37% nests/day	4.11% nests/day	22%	Small sample
Laying date	38	1968-2006	19	Linear decline	May 25	May 16	-9 days	Small sample


Clutch size 1966—2007

Tree Pipit




Brood size 1966—2007

Tree Pipit




Egg stage nest failure rate

Tree Pipit




Chick stage nest failure rate

Tree Pipit



Laying date 1966—2007

Tree Pipit



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

MEADOW PIPIT *Anthus pratensis*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
UK: amber (25–50% population decline)

Long-term trend

England: moderate decline

UK population size

1,680,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

The CBC/BBS trend has been downward since the mid 1970s, accompanied by a range contraction from lowland England ([Gibbons *et al.* 1993](#)). Meadow Pipits are partial migrants and conditions on the Iberian wintering grounds have been linked to the decline, as have losses of marginal land from parts of the breeding range ([Gibbons *et al.* 1993](#)). Moorland, the key Meadow Pipit habitat, was not covered well by the CBC, leading to some doubt about the significance of the early results for this species, but BBS now provides more representative monitoring and has enabled the species to move from the green to the amber list. Nest failure rates during the 12-day nestling stage have declined markedly, which may reflect the loss of birds from suboptimal habitat. A trend towards earlier laying is probably related to climate change ([Crick & Sparks 1999](#)). Meadow Pipits are estimated to have declined by 57% across Europe during 1980–2005 ([PECBMS 2007](#)).



Population changes

CBC/BBS England 1966–2007 Meadow Pipit




Table of population changes for Meadow Pipit

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	136	-46	-74	-24	>25	
	25	1981-2006	192	-43	-61	-26	>25	
	10	1996-2006	381	-5	-15	6		
	5	2001-2006	411	-8	-16	1		
BBS UK	11	1995-2006	703	-13	-19	-6		
	10	1996-2006	718	-13	-18	-7		
	5	2001-2006	768	-13	-17	-8		
BBS England	11	1995-2006	360	-9	-18	1		
	10	1996-2006	369	-5	-14	5		
	5	2001-2006	411	-6	-13	1		
BBS Scotland	11	1995-2006	197	-23	-30	-13		
	10	1996-2006	195	-23	-30	-14		
	5	2001-2006	185	-17	-23	-11		
BBS Wales	11	1995-2006	84	4	-10	20		
	10	1996-2006	86	4	-12	19		
	5	2001-2006	96	-13	-21	-1		
BBS N.Ireland	11	1995-2006	59	51	21	101		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS UK 1994—2007

Meadow Pipit




BBS England 1994—2007

Meadow Pipit




BBS Scotland 1994—2007

Meadow Pipit




BBS Wales 1994—2007

Meadow Pipit



BBS N. Ireland 1994—2007

Meadow Pipit




Productivity trends

Table of productivity changes for Meadow Pipit

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	36	None				
Brood size	38	1968-2006	68	None				
Daily failure rate (eggs)	38	1968-2006	46	None				
Daily failure rate (chicks)	38	1968-2006	60	Linear decline	2.71% nests/day	1.08% nests/day	-60.1%	
Laying date	38	1968-2006	38	None				


Clutch size 1966—2007

Meadow Pipit




Brood size 1966—2007

Meadow Pipit




Egg stage nest failure rate

Meadow Pipit




Chick stage nest failure rate

Meadow Pipit



Laying date 1966—2007

Meadow Pipit



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

YELLOW WAGTAIL

Motacilla flava

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
 UK: amber (25–50% population decline)
 UK Biodiversity Action Plan: priority species

Long-term trend

UK, England: rapid decline

UK population size

11,500–26,500 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS and WBS trends: [BiE04](#), [APEP06](#))

Status summary

Britain holds almost the entire population of the distinctive race *flavissima*, and so population changes in the UK are of global conservation significance. Yellow Wagtails have been in decline since the early 1980s, according to CBC/BBS and especially WBS, and have now been moved from the green to the amber list. Further losses since 1999 already suggest that red listing is appropriate. [Gibbons et al. \(1993\)](#) identified a range contraction towards a core area in central England, concurrent with the early years of decline. Farmland drainage, the conversion of pasture to arable land, the change from spring to winter cereals, and the loss of insects associated with cattle have been cited as possible causes ([Gibbons et al. 1993](#), [Nelson et al. 2003](#)). Although nest record sample sizes are small, there has been a notable reduction in brood size since the mid 1960s, and the species is listed as of NRS concern ([Leech & Barimore 2008](#)). The European trend, which includes other races of the species, has also been downward since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007

Yellow Wagtail




Table of population changes for Yellow Wagtail


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	71	-74	-88	-46	>50	
	25	1981-2006	91	-70	-82	-55	>50	
	10	1996-2006	164	-42	-53	-31	>25	
	5	2001-2006	149	-25	-35	-14	>25	
CBC/BBS England	39	1967-2006	69	-72	-85	-38	>50	
	25	1981-2006	89	-68	-81	-49	>50	
	10	1996-2006	160	-42	-49	-31	>25	
	5	2001-2006	146	-24	-31	-13		
WBS waterways	31	1975-2006	18	-97	-99	-93	>50	Small sample
	25	1981-2006	16	-97	-99	-95	>50	Small sample
BBS UK	11	1995-2006	158	-44	-53	-34	>25	
	10	1996-2006	157	-43	-52	-33	>25	
	5	2001-2006	149	-25	-31	-16		
BBS England	11	1995-2006	154	-43	-52	-33	>25	
	10	1996-2006	153	-42	-51	-32	>25	

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007
Yellow Wagtail




Waterways Bird Survey 1974—2007
Yellow Wagtail



BBS UK 1994—2007
Yellow Wagtail



BBS England 1994—2007
Yellow Wagtail




Productivity trends

Table of productivity changes for Yellow Wagtail

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size	38	1968-2006	13	Linear decline	4.83 chicks	4.34 chicks	-10.1%	Small sample

Insufficient data on clutch size available for this species

Brood size 1966—2007
Yellow Wagtail



Insufficient data on nest failure available for this species

Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GREY WAGTAIL

Motacilla cinerea

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: amber (25–50% population decline)

Long-term trend

UK: probable shallow decline

UK population size

38,400–46,200 pairs in 2000 (1988–91 Atlas estimate updated using CBC and WBS trends: [BIE04](#), [APEP06](#))

Status summary

Grey Wagtails occur at highest densities along fast-flowing upland streams. WBS shows a fluctuating population size along waterways, with a fall during the late 1970s and early 1980s from an initial high point in 1974, and some increase since the late 1990s. The species has recently been moved from the green to the amber list, because of a 41% decline recorded between 1975 and 1999. BBS figures showed an initial ten-year phase of increase, which has now stabilised. The trends for Grey Wagtail are very similar to those for [Pied Wagtail](#), suggesting that similar factors may be affecting these two species. Clutch and brood size of Grey Wagtails rose as the population fell, and are now getting smaller again, raising NRS concern ([Leech & Barimore 2008](#)). Nest failure rates have dropped substantially. Grey Wagtails are estimated to have declined by 54% across Europe during 1982–2005 ([PECBMS 2007](#)).



Population changes

Waterways Bird Survey 1974–2007

Grey Wagtail






Table of population changes for Grey Wagtail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	55	-23	-44	0		
	25	1981-2006	57	35	9	60		
	10	1996-2006	53	32	14	48		
	5	2001-2006	48	11	-3	22		
BBS UK	11	1995-2006	192	40	18	62		
	10	1996-2006	196	43	20	60		
	5	2001-2006	235	-5	-14	4		
BBS England	11	1995-2006	124	48	22	73		
	10	1996-2006	128	49	27	70		
	5	2001-2006	155	6	-4	14		








The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Table of productivity changes for Grey Wagtail

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	38	Curvilinear	4.68 eggs	4.65 eggs	-0.7%	
Brood size	38	1968-2006	80	Curvilinear	3.95 chicks	3.95 chicks	0.1%	
Daily failure rate (eggs)	38	1968-2006	59	None				
Daily failure rate (chicks)	38	1968-2006	58	Linear decline	2.06% nests/day	0.98% nests/day	-52.4%	
Laying date	38	1968-2006	60	None				



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

PIED WAGTAIL

Motacilla alba

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK: uncertain

UK population size

272,000–352,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS and WBS trends: [BIE04](#), [APEP06](#))

Status summary

Britain and Ireland together hold almost the entire population of the distinctive dark-backed race *yarrellii*, and for this reason population changes in the UK are of global conservation significance. The CBC shows that a strong increase occurred up to the mid 1970s, such that populations have shown moderate increase overall since 1966. Since 1974, however, the results of monitoring are somewhat conflicting: CBC/BBS and WBS trends fluctuate in parallel but, whereas little overall change is evident in the CBC/BBS index, WBS has shown a moderate decline, perhaps suggesting the influence of factors specific to linear waterways. The long-term trend in abundance is similar to those shown by [Wren](#) and [Long-tailed Tit](#), two other resident insectivores ([Siriwardena et al. 1998a](#)). Average clutch and brood sizes have declined a little, raising NRS concern ([Leech & Barimore 2008](#)), but this has been counteracted by a fall in nest failure rates.



Population changes

CBC/BBS UK 1966–2007

Pied Wagtail




Table of population changes for Pied Wagtail

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	409	80	36	147		
	25	1981-2006	588	2	-17	31		
	10	1996-2006	1206	3	-3	9		
	5	2001-2006	1288	-7	-12	-3		
CBC/BBS England	39	1967-2006	315	86	32	163		
	25	1981-2006	449	5	-18	41		
	10	1996-2006	916	6	0	12		
	5	2001-2006	979	-7	-11	-3		
WBS waterways	31	1975-2006	64	-53	-66	-43	>50	
	25	1981-2006	64	-39	-53	-25	>25	
	10	1996-2006	58	-18	-32	-4		
	5	2001-2006	52	-10	-23	2		
BBS UK	11	1995-2006	1139	6	-1	12		
	10	1996-2006	1171	4	-3	10		
	5	2001-2006	1288	-6	-10	-2		
BBS England	11	1995-2006	865	9	1	16		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Scotland	11	1995-2006	123	-1	-17	17		
	10	1996-2006	124	-3	-17	13		
	5	2001-2006	123	-9	-20	-1		
BBS Wales	11	1995-2006	110	9	-10	32		
	10	1996-2006	114	8	-10	30		
	5	2001-2006	131	3	-8	14		
BBS N.Ireland	11	1995-2006	37	33	.	.		
	10	1996-2006	41	28	.	.		
	5	2001-2006	50	11	.	.		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Pied Wagtail




Waterways Bird Survey 1974—2007

Pied Wagtail




BBS UK 1994—2007

Pied Wagtail




BBS England 1994—2007

Pied Wagtail




BBS Scotland 1994—2007

Pied Wagtail



BBS Wales 1994—2007

Pied Wagtail




Productivity trends

Table of productivity changes for Pied Wagtail

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	59	Linear decline	5.12 eggs	4.92 eggs	-3.9%	
Brood size	38	1968-2006	114	Linear decline	4.53 chicks	4.34 chicks	-4.3%	
Daily failure rate (eggs)	38	1968-2006	83	Curvilinear	1.52% nests/day	0.68% nests/day	-55.3%	
Daily failure rate (chicks)	38	1968-2006	91	Linear decline	1.27% nests/day	0.84% nests/day	-33.9%	
Laying date	38	1968-2006	79	None				


Clutch size 1966—2007

Pied Wagtail




Brood size 1966—2007

Pied Wagtail




Egg stage nest failure rate

Pied Wagtail




Chick stage nest failure rate

Pied Wagtail



Laying date 1966—2007

Pied Wagtail



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

DIPPER *Cinclus cinclus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK: fluctuating, with no long-term trend

UK population size

6,800–20,000 pairs in 2000 (1988–91 Atlas estimate updated using WBS trend: [BiE04](#), [APEP06](#))

Status summary

The WBS trend shows that Dipper populations have fluctuated over the last thirty years, but shown little overall trend. The species is unusually sensitive to acidity and other water pollution ([Ormerod & Tyler 1989, 1990](#)), with lower breeding densities and productivity on acidic than on more neutral streams ([Ormerod et al. 1991](#), [Vickery 1991, 1992](#)). Breeding performance has improved strongly over time, and laying dates have shifted earlier, perhaps because of climate change ([Crick & Sparks 1999](#)). Broods now average larger than in the late 1960s and 1970s, and there has been substantial reduction in failure rates of nests at the egg stage.



Population changes

Waterways Bird Survey 1974–2007 Dipper





Table of population changes for Dipper

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
WBS waterways	31	1975-2006	35	-11	-38	28		
	25	1981-2006	37	4	-28	49		
	10	1996-2006	31	3	-16	32		
	5	2001-2006	29	-3	-16	15		
BBS UK	11	1995-2006	51	0	-26	36		
	10	1996-2006	52	-2	-27	27		
	5	2001-2006	53	-7	-23	7		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

BBS UK 1994–2007




Productivity trends


Table of productivity changes for Dipper

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	72	Curvilinear	4.48 eggs	4.5 eggs	0.4%	
Brood size	38	1968-2006	135	Curvilinear	3.4 chicks	3.69 chicks	8.6%	
Daily failure rate (eggs)	38	1968-2006	100	Curvilinear	2.57% nests/day	0.33% nests/day	-87.2%	
Daily failure rate (chicks)	38	1968-2006	78	Curvilinear	0.59% nests/day	0.62% nests/day	5.1%	
Laying date	38	1968-2006	59	Linear decline	Apr 18	Apr 9	-9 days	


Clutch size 1966–2007




Brood size 1966–2007




Egg stage nest failure rate



Chick stage nest failure rate



Laying date 1966–2007



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WREN

Troglodytes troglodytes

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

UK Biodiversity Action Plan: priority species (Fair Isle & St Kilda races only)

Long-term trend

UK, England: moderate increase

UK population size

8,512,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

The Wren's current UK population estimate is the highest for any species. Abundance can vary sharply from year to year in this species, however, although this is not evident from the smoothed trends presented here. Annual numbers are influenced by mortality rates that may be very high in severe winters and by the species' high breeding potential ([Peach et al. 1995b](#)). Wren numbers in the UK were greatly depleted by the cold winter of 1962/63 ([Marchant et al. 1990](#)). Following a rapid recovery up to the mid 1970s, abundance fell again in response to a further series of cold winters only to return to its previous high level. BBS results suggest that increase since 1994 has been much stronger in Scotland and Northern Ireland than in Wales and England. Rather fewer nests are now failing at the egg stage. Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007

Wren




Table of population changes for Wren

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	818	98	74	120		
	25	1981-2006	1155	51	36	60		
	10	1996-2006	2265	28	22	30		
	5	2001-2006	2392	1	-2	3		
CBC/BBS England	39	1967-2006	651	90	65	118		
	25	1981-2006	910	42	28	55		
	10	1996-2006	1764	22	17	24		
	5	2001-2006	1847	-2	-4	0		
CES adults	22	1984-2006	95	58	35	87		
	10	1996-2006	107	-1	-9	7		
	5	2001-2006	98	-11	-18	-4		
CES juveniles	22	1984-2006	94	30	6	61		
	10	1996-2006	106	-5	-15	5		
	5	2001-2006	99	-16	-23	-11		
BBS UK	11	1995-2006	2107	24	18	27		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS England	11	1995-2006	1634	16	11	18		
	10	1996-2006	1676	20	15	22		
	5	2001-2006	1847	2	-1	3		
BBS Scotland	11	1995-2006	194	66	43	80		
	10	1996-2006	195	56	34	66		
	5	2001-2006	205	11	2	21		
BBS Wales	11	1995-2006	185	17	4	25		
	10	1996-2006	192	21	8	28		
	5	2001-2006	223	0	-5	6		
BBS N.Ireland	11	1995-2006	82	62	21	95		
	10	1996-2006	88	40	13	56		
	5	2001-2006	102	3	-5	12		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Wren




CES adult abundance 1983—2007

Wren




CES juvenile abundance 1983—2007

Wren




BBS UK 1994—2007

Wren




BBS England 1994—2007

Wren




BBS Scotland 1994—2007

Wren




BBS Wales 1994—2007

Wren



BBS N. Ireland 1994—2007

Wren




Productivity trends


Table of productivity changes for Wren

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	95	None				
Brood size	38	1968-2006	96	None				
Daily failure rate (eggs)	38	1968-2006	141	Linear decline	1.83% nests/day	1.43% nests/day	-21.9%	
Daily failure rate (chicks)	38	1968-2006	98	None				
Laying date	38	1968-2006	87	Linear decline	May 14	May 9	-5 days	
Juvenile to Adult ratio (CES)	22	1984-2006	98	Smoothed trend	114 Index value	100 Index value	-12%	
Juvenile to Adult ratio (CES)	10	1996-2006	109	Smoothed trend	105 Index value	100 Index value	-5%	
Juvenile to Adult ratio (CES)	5	2001-2006	100	Smoothed trend	106 Index value	100 Index value	-6%	


Clutch size 1966–2007




Brood size 1966–2007




Egg stage nest failure rate




Chick stage nest failure rate



Laying date 1966–2007



CES productivity 1983–2007



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

DUNNOCK

Prunella modularis

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: amber (25–50% population decline)

UK Biodiversity Action Plan: priority species

Long-term trend

UK, England: moderate decline

UK population size

2,163,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Dunnock abundance fell substantially between the mid 1970s and mid 1980s, after a period of population stability. Some recovery has occurred throughout the UK since the late 1990s, but the species is still amber listed. The cause of the decline remains unknown. In many lowland woods, canopy closure in the absence of forest management and increasing browsing pressure from deer are likely to have reduced the suitability of the habitat for this species ([Fuller et al. 2005](#)). There has been little variation in survival rates over time ([Siriwardena et al. 1998a](#)). Clutch and brood sizes increased as the population fell. Egg-stage nest failure rates are currently increasing, and are of NRS concern ([Leech & Barimore 2008](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007

Dunnock




Table of population changes for Dunnock

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	703	-33	-42	-22	>25	
	25	1981-2006	978	-6	-17	5		
	10	1996-2006	1910	23	18	28		
	5	2001-2006	2024	9	6	12		
CBC/BBS England	39	1967-2006	582	-37	-46	-26	>25	
	25	1981-2006	805	-10	-22	0		
	10	1996-2006	1555	20	16	24		
	5	2001-2006	1632	8	6	12		
CES adults	22	1984-2006	94	-1	-16	14		
	10	1996-2006	105	2	-6	10		
	5	2001-2006	95	-2	-9	7		
CES juveniles	22	1984-2006	91	-19	-37	4		
	10	1996-2006	103	-8	-17	0		
	5	2001-2006	95	-12	-20	-5		
BBS UK	11	1995-2006	1769	23	18	28		
	10	1996-2006	1822	22	18	27		
	5	2001-2006	2024	10	7	13		


BBS E _n Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	1632	10	7	12		
BBS Scotland	11	1995-2006	115	39	14	64		
	10	1996-2006	119	33	10	53		
	5	2001-2006	128	10	-7	24		
BBS Wales	11	1995-2006	140	40	19	68		
	10	1996-2006	146	36	19	59		
	5	2001-2006	170	16	6	26		
BBS N.Ireland	11	1995-2006	62	90	37	136		
	10	1996-2006	68	51	19	76		
	5	2001-2006	82	0	-9	13		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Dunnock




CES adult abundance 1983—2007

Dunnock




CES juvenile abundance 1983—2007

Dunnock




BBS UK 1994—2007

Dunnock




BBS England 1994—2007

Dunnock




BBS Scotland 1994—2007

Dunnock




BBS Wales 1994—2007

Dunnock



BBS N. Ireland 1994—2007







Dunnock



Productivity trends

Table of productivity changes for Dunnock

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	99	Linear increase	3.94 eggs	4.2 eggs	6.6%	
Brood size	38	1968-2006	106	Linear increase	3.48 chicks	3.63 chicks	4.5%	
Daily failure rate (eggs)	38	1968-2006	143	Curvilinear	2.51% nests/day	2.48% nests/day	-1.2%	
Daily failure rate (chicks)	38	1968-2006	117	None				
Laying date	38	1968-2006	79	Linear decline	May 3	Apr 29	-4 days	
Juvenile to Adult ratio (CES)	22	1984-2006	98	Smoothed trend	115 Index value	100 Index value	-13%	
Juvenile to Adult ratio (CES)	10	1996-2006	109	Smoothed trend	114 Index value	100 Index value	-13%	
Juvenile to Adult ratio (CES)	5	2001-2006	99	Smoothed trend	122 Index value	100 Index value	-18%	



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

ROBIN

Erithacus rubecula

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: shallow increase

UK population size

5,895,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Robins have increased markedly since the mid 1980s, according to both CBC/BBS and CES results, having been set back earlier by a succession of cold winters. Significant improvements have occurred concurrently in breeding performance, as measured by nest record data, due to reductions in nest failure rates at both egg and chick stages, although CES productivity measures have declined. The CES and BBS data show that marked and significant annual fluctuations occur, perhaps in response to winter weather, although these are not evident in the smoothed trends that are presented. Laying dates have advanced by almost a week since the 1960s. Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007

Robin





Table of population changes for Robin

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	794	39	25	56		
	25	1981-2006	1118	44	34	55		
	10	1996-2006	2195	17	14	20		
	5	2001-2006	2310	1	-1	3		
CBC/BBS England	39	1967-2006	637	49	32	65		
	25	1981-2006	891	52	39	62		
	10	1996-2006	1730	20	17	23		
	5	2001-2006	1808	3	0	4		
CES adults	22	1984-2006	89	63	40	87		
	10	1996-2006	101	9	0	18		
	5	2001-2006	94	-3	-13	7		
CES juveniles	22	1984-2006	94	26	9	51		
	10	1996-2006	107	2	-5	10		
	5	2001-2006	98	-5	-11	1		
BBS UK	11	1995-2006	2037	17	13	20		
	10	1996-2006	2096	16	13	20		
	5	2001-2006	2310	3	0	5		
BBS England	11	1995-2006	1600	20	18	24		

Clutch	Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size									
	Daily failure rate (eggs)	38	1968-2006	188	Curvilinear	2.47% nests/day	1.03% nests/day	-58.3%	
	Daily failure rate (chicks)	38	1968-2006	162	Curvilinear	2.49% nests/day	1.77% nests/day	-28.9%	
	Laying date	38	1968-2006	120	Linear decline	Apr 28	Apr 22	-6 days	
	Juvenile to Adult ratio (CES)	22	1984-2006	98	Smoothed trend	113 Index value	100 Index value	-12%	
	Juvenile to Adult ratio (CES)	10	1996-2006	110	Smoothed trend	105 Index value	100 Index value	-5%	
	Juvenile to Adult ratio (CES)	5	2001-2006	100	Smoothed trend	97 Index value	100 Index value	3%	


Clutch size 1966–2007

Robin




Brood size 1966–2007

Robin




Egg stage nest failure rate

Robin




Chick stage nest failure rate

Robin




Laying date 1966–2007

Robin



CES productivity 1983–2007

Robin



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

NIGHTINGALE

Luscinia megarhynchos



- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
UK: amber (25–50% distribution decline)

Long-term trend

UK: probable shallow decline


UK population size

6,700 (5,600–9,350) males in 1999 ([Wilson et al. 2002](#): BIE04, APEP06)

Status summary

In 1999, the BTO organised a national survey of Nightingales, which showed a marked range contraction since the previous survey in 1980, but only an 8% overall population decline ([Wilson et al. 2002](#); for more details [click here](#)). Nightingales are scarce birds, and CBC and BBS data are correspondingly meagre. Nevertheless, analysis of the available CBC data shows continuous decline (G.M. Siriwardena, unpubl.) and CES suggests a fluctuating pattern, or possible decline. [Fuller et al. \(2005\)](#) suggest the likely causes of Nightingale decline relate to pressures on migration and in winter, perhaps compounded by habitat loss in Britain. The increasing intensity of browsing by deer is known to reduce habitat quality for this species ([Gill & Fuller 2007](#)). CES indicates a sharp decline in productivity during the 1980s, perhaps because Nightingale nesting success may be adversely affected by cold and wet springs. Nightingale is one of the most strongly declining bird species in Europe, having decreased at an annual rate of 4% during 1980–2005; this overall figure masks a contrast between severe decreases in southern and western Europe and strong increases in the east of the range ([PECBMS 2007](#)).

Population changes



My BTO



VOLUNTEER

JOIN


DONATE

MENU



The page you are looking for could not be found.

- Try searching using the search bar (top right)
- Navigate to the area you are looking for using the main menu (above)
- [Return to our homepage](#)
- Email us about what you are trying to find, or to tell us about a broken link - info@bto.org




See live NBC results and watch the breeding season unfold



BTO science contributes to our understanding of future scenarios, and informing policies and conservation management strategies to help species adapt.



The NI Seabird Report is unique in the UK, presenting the most up-to-date seabird monitoring data collected by volunteers, NGOs and NIEA from around Northern Ireland. Each March we present freshly produced hard copies...



✉ info@bto.org

[Press releases](#)
[Terms & Conditions](#)
[Privacy](#)
[Cookies](#)



Registered Charity Number 216652 (England & Wales), SC039193 (Scotland)
 © British Trust for Ornithology, BTO, The Nunnery, Thetford, Norfolk IP24 2PU, Tel: +44 (0)1842 750050 Fax: +44 (0)1842 750030


REPORT IT TO US ►

E-NEWSLETTER SIGN UP






 The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Table of productivity changes for Nightingale

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Juvenile to Adult ratio (CES)	22	1984-2006	11	Smoothed trend	684 Index value	100 Index value	-85% >50	Small sample
Juvenile to Adult ratio (CES)	10	1996-2006	10	Smoothed trend	112 Index value	100 Index value	-11%	Small sample


Insufficient data on clutch size available for this species

Insufficient data on brood size available for this species

Insufficient data on nest failure available for this species

Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

These optional cookies are used for collecting anonymous visit data only.



REDSTART *Phoenicurus phoenicurus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 2 (depleted)
UK: amber (European status)

Long-term trend

UK, England: shallow decline

UK population size

At least 101,000 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

The decline in the late 1960s and early 1970s was thought to be due to severe drought conditions in the Sahel wintering area in Africa ([Marchant *et al.* 1990](#)). There was a loss of range of 20% in Britain between 1968–72 and 1988–91, in terms of the numbers of occupied 10-km squares ([Gibbons *et al.* 1993](#)). A recovery in population size began in the mid 1970s and appears to have continued, at least in England, into the late 1990s. This increase has been associated with improving breeding performance and progressively earlier laying dates. The trend towards earlier laying can be partly explained by recent climate change ([Crick & Sparks 1999](#)).



Population changes

CBC/BBS UK 1966–2007 Redstart





Table of population changes for Redstart

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	63	-19	-44	23		
	25	1981-2006	86	27	-8	60		
	10	1996-2006	156	-7	-22	4		
	5	2001-2006	144	-5	-15	5		
CBC/BBS England	39	1967-2006	38	-18	-42	37		Small CBC sample
	25	1981-2006	51	27	-8	62		Small CBC sample
	10	1996-2006	87	-11	-23	4		
	5	2001-2006	77	-9	-20	2		
BBS UK	11	1995-2006	142	0	-14	12		
	10	1996-2006	145	-7	-18	4		
	5	2001-2006	144	-6	-14	2		
BBS England	11	1995-2006	76	-1	-18	15		
	10	1996-2006	77	-10	-23	4		
	5	2001-2006	77	-10	-18	1		
BBS Wales	11	1995-2006	54	-8	-27	15		
	10	1996-2006	55	-10	-28	12		
	5	2001-2006	56	-8	-22	5		


CBC/BBS England 1966—2007

Redstart




BBS UK 1994—2007

Redstart




BBS England 1994—2007

Redstart



BBS Wales 1994—2007

Redstart




Productivity trends

Table of productivity changes for Redstart

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	48	Curvilinear	5.89 eggs	6.23 eggs	5.8%	
Brood size	38	1968-2006	85	Curvilinear	5.1 chicks	5.48 chicks	7.4%	
Daily failure rate (eggs)	38	1968-2006	73	Linear decline	1.16% nests/day	0.31% nests/day	-73.3%	
Daily failure rate (chicks)	38	1968-2006	52	Linear decline	1.23% nests/day	0.39% nests/day	-68.3%	
Laying date	38	1968-2006	61	Curvilinear	May 21	May 9	-12 days	


Clutch size 1966—2007

Redstart




Brood size 1966—2007

Redstart




Egg stage nest failure rate

Redstart



Chick stage nest failure rate


Redstart



Insufficient data on CES available for this species

Laying date 1966–2007

Redstart



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WHINCHAT *Saxicola rubetra*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
UK: green

Long-term trend

UK: decline

UK population size

14,000–28,000 pairs in 1990 (1988–91 Atlas: [APEP06](#));
11,000–22,100 pairs in 2000 (updated using BBS trend: [BIE04](#))

Status summary

Whinchats were not monitored until the BBS began in 1994. By then, however, [Gibbons *et al.* \(1993\)](#) had already identified a major range contraction, mainly from lowland England, that was probably at least partly due to the loss of marginal farmland habitats ([Marchant *et al.* 1990](#)). Further extinctions have occurred since then among the remaining pockets of lowland breeders. BBS data indicate that further strong population decline took place during the 1990s, raising BTO alerts for the UK as a whole as well as for England. Nest record samples are small, but indicate substantial recent rises in nest losses at the egg and chick stages, which are of NRS concern ([Leech & Barimore 2008](#)). Whinchats are estimated to have declined by 55% across Europe during 1980–2005 ([PECBMS 2007](#)). Despite these changes, the species currently has no conservation listings.



Population changes

BBS UK 1994–2007 Whinchat




Table of population changes for Whinchat

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	74	-38	-53	-18	>25	
	10	1996-2006	73	-38	-53	-18	>25	
	5	2001-2006	62	-14	-29	5		
BBS England	11	1995-2006	30	-34	-60	-9	>25	
	10	1996-2006	30	-32	-58	-9	>25	








The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

Productivity trends

Table of productivity changes for Whinchat

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	12	None				Small sample
Brood size	38	1968-	37	None				

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Daily failure rate (eggs)								
Daily failure rate (chicks)		2006			nests/day	nests/day		sample
Laying date	38	1968-2006	27	Curvilinear	May 28	May 23	-5 days	Small sample



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

STONECHAT *Saxicola torquatus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: amber (European status)

Long-term trend

UK: uncertain, possible decline

UK population size

9,000–23,000 pairs in 1990 (1988–91 Atlas: [APEP06](#));
19,300–49,400 pairs in 2000 (updated using BBS trend: [BIE04](#))

Status summary

Numerical trends were not measured before the start of the BBS, but a long-term decline is suspected: severe winter weather, and loss and fragmentation of suitable breeding habitat in many inland regions, are believed to have reduced the population from the 1940s onward ([Marchant et al. 1990](#)). Breeding atlas data showed a substantial contraction in the Stonechat's range between the early 1970s and late 1980s ([Gibbons et al. 1993](#)). Nest failure rates have fallen markedly over the long term. Against this background, the current, strongly increasing BBS trend represents substantial recovery. Following similar increases widely across Europe, the species is now provisionally categorised as 'secure' ([BirdLife International 2004](#)). The UK amber listing rests on the earlier European decline, so a change to green may be warranted at the next review.



Population changes

BBS UK 1994–2007 Stonechat




Table of population changes for Stonechat


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	122	200	140	265		
	10	1996-2006	128	188	136	228		
	5	2001-2006	172	48	29	61		
BBS England	11	1995-2006	48	167	83	278		
	10	1996-2006	50	156	75	253		
	5	2001-2006	69	37	11	61		
BBS Scotland	5	2001-2006	38	79	40	109		
BBS Wales	11	1995-2006	33	207	117	360		
	10	1996-2006	35	158	81	247		
	5	2001-2006	47	24	-5	54		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

BBS England 1994–2007

Stonechat




Productivity trends

Table of productivity changes for Stonechat

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	30	None				
Brood size	38	1968-2006	59	None				
Daily failure rate (eggs)	38	1968-2006	35	Curvilinear	0.52% nests/day	0.27% nests/day	-48.1%	
Daily failure rate (chicks)	38	1968-2006	54	Curvilinear	1.16% nests/day	0.4% nests/day	-65.5%	
Laying date	38	1968-2006	35	Curvilinear	May 3	Apr 27	-6 days	


Clutch size 1966–2007

Stonechat




Brood size 1966–2007

Stonechat




Egg stage nest failure rate

Stonechat




Chick stage nest failure rate

Stonechat



Laying date 1966–2007

Stonechat



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WHEATEAR

Oenanthe oenanthe

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (declining)

UK: green

Long-term trend

UK: possible decline

UK population size

56,000 pairs in 1990 (1988–91 Atlas: [APEP06](#)); 52,500

pairs in 2000 (updated using BBS trend: [BiE04](#));

100,000–200,000 pairs in Britain ([Sellers 2006](#))



Status summary

Although it is a common breeding species in many upland areas, the Wheatear was not monitored at the UK scale until the BBS began in 1994. [Gibbons *et al.* \(1993\)](#) had by then identified range contractions from lowland Britain since 1968–72, perhaps due to losses of suitable grassland and declines in rabbit abundance. BBS shows wide fluctuations, with further decrease in England and Wales, but as yet no clear trend in abundance since 1994 in Scotland. BBS data indicate that the estimates of UK population made for the 1988–91 Atlas may have been far too low, possibly by an order of magnitude ([Gillings *et al.* 2007](#)). Failure rates at the egg stage (18 days, comprising 14 days incubation and 4 days laying) have fallen substantially. Wheatear is one of the most strongly declining bird species in Europe, having decreased at an annual rate of 5% during 1980–2005 ([PECBMS 2007](#)). Following widespread declines across Europe during the 1990s, the European status of this species is no longer considered 'secure' ([BirdLife International 2004](#)).

Population changes

BBS UK 1994–2007

Wheatear




Table of population changes for Wheatear


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	269	-13	-28	4		
	10	1996-2006	273	-17	-31	1		
	5	2001-2006	293	-1	-16	14		
BBS England	11	1995-2006	133	-21	-35	1		
	10	1996-2006	135	-22	-35	-4		
	5	2001-2006	149	-13	-27	4		
BBS Scotland	11	1995-2006	75	-10	-34	27		
	10	1996-2006	74	-14	-38	21		
	5	2001-2006	71	4	-18	31		
BBS Wales	11	1995-2006	50	-25	-43	-8	>25	
	10	1996-2006	52	-24	-42	-9		
	5	2001-2006	59	-8	-20	6		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS England 1994–2007

Wheatear




BBS Scotland 1994–2007

Wheatear



BBS Wales 1994–2007

Wheatear




Productivity trends

Table of productivity changes for Wheatear

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	None				Small sample
Brood size	38	1968-2006	59	None				
Daily failure rate (eggs)	38	1968-2006	18	Curvilinear	0.82% nests/day	0.09% nests/day	-89%	Small sample
Daily failure rate (chicks)	38	1968-2006	40	None				
Laying date	38	1968-2006	13	None				Small sample


Clutch size 1966–2007

Wheatear




Brood size 1966–2007

Wheatear




Egg stage nest failure rate

Wheatear




Chick stage nest failure rate

Wheatear



Laying date 1966–2007

Wheatear



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

RING OUZEL *Turdus torquatus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
 UK: red (>50% population decline)
 UK Biodiversity Action Plan: priority species

Long-term trend

UK: probable decline

UK population size

6,157–7,549 pairs in 1999 (Wotton *et al.* 2002: BIE04, APEP06)

Status summary

The first breeding atlases showed a decline of 27% in the number of 10-km squares occupied between 1968–72 and 1988–91 (Gibbons *et al.* 1993), and the extent of population decline has since been established by a special survey: a 58% population decline was estimated for the period between 1988–91 and 1999, warranting red listing for this species (Gregory *et al.* 2002). British & Irish bird observatory data show a decline in spring passage Ring Ouzels at western locations during 1970–98 that matches the estimated UK breeding decline, but no decline at eastern observatories where most birds are of Fennoscandian origin (Burfield & Brooke 2005). These authors infer that, since these populations winter together, the reasons for decline among UK breeders must lie on the breeding grounds or on passage: they also point out that UK birds are more exposed to hunting pressures, particularly in southwest France. It has proved difficult to establish any reasons for decline that are linked to the breeding grounds (Buchanan *et al.* 2003). In southeast Scotland, however, the breeding sites that are still occupied tend to be those at higher altitude and that have retained an extensive cover of heather (Sim *et al.* 2007b). In the same study, it was shown that declines were greatest in years following warm summers on the breeding grounds and also greater two years after high spring rainfall in Morocco: these results suggest that the population decline could be linked to reduced food supplies, and consequently higher rates of natural mortality, in autumn and winter (Beale *et al.* 2006).



Population changes

Annual breeding population changes for this species are not currently monitored by BTO


Productivity trends

Table of productivity changes for Ring Ouzel


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size	38	1968-2006	22	None				Small sample
Daily failure rate (eggs)	38	1968-2006	11	None				Small sample
Daily failure rate (chicks)	38	1968-2006	14	None				Small sample
Laying date	38	1968-2006	24	Linear decline	May 14	May 8	-6 days	Small sample

Insufficient data on clutch size available for this species


Brood size 1966–2007
Ring Ouzel



Egg stage nest failure rate
Ring Ouzel




Chick stage nest failure rate
Ring Ouzel



Insufficient data on CES available for this species

Laying date 1966–2007

Ring Ouzel



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

BLACKBIRD *Turdus merula*

• Population changes

• Productivity trends

• Additional information



Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: shallow decline

UK population size

4,935,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Both CBC/BBS and CES data show long-term declines in Blackbird abundance, but recent increases suggest that the population has begun to recover. The moderate-decline criteria for amber listing and for BTO alerts are no longer met, and the species is now listed in the green category. CBC results indicate that the decline began in the mid 1970s. Nest success has improved over this period, and it is likely that reduced survival drove the decline ([Siriwardena et al. 1998a](#)). Agricultural intensification is likely to have contributed ([Fuller et al. 1995](#)), but, since numbers fell in woodland as well as farmland, additional factors probably operated.

Population changes

CBC/BBS UK 1966–2007 Blackbird




Table of population changes for Blackbird


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	828	-16	-23	-7		
	25	1981-2006	1166	2	-5	9		
	10	1996-2006	2292	21	18	24		
	5	2001-2006	2406	4	2	6		
CBC/BBS England	39	1967-2006	671	-18	-25	-10		
	25	1981-2006	938	-1	-9	6		
	10	1996-2006	1823	19	16	22		
	5	2001-2006	1893	3	2	5		
CES adults	22	1984-2006	96	-13	-25	0		
	10	1996-2006	107	-9	-15	0		
	5	2001-2006	97	-17	-23	-10		
CES juveniles	22	1984-2006	86	-25	-45	0	>25	
	10	1996-2006	97	8	-4	24		
	5	2001-2006	91	12	-7	35		
BBS UK	11	1995-2006	2128	22	19	26		
	10	1996-2006	2190	20	17	23		
	5	2001-2006	2406	5	4	7		
BBS England	11	1995-2006	1691	20	17	24		
	10	1996-2006	1735	18	16	21		

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Scotland	10	1996-2006	168	28	14	46		
	5	2001-2006	180	12	2	21		
BBS Wales	11	1995-2006	183	40	29	50		
	10	1996-2006	191	37	27	45		
BBS N.Ireland	5	2001-2006	223	17	12	21		
	11	1995-2006	77	40	10	55		
	10	1996-2006	83	20	-1	27		
	5	2001-2006	95	-10	-17	-4		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007 Blackbird




CES adult abundance 1983–2007 Blackbird




CES juvenile abundance 1983–2007 Blackbird




BBS UK 1994–2007 Blackbird




BBS England 1994–2007 Blackbird




BBS Scotland 1994–2007 Blackbird



BBS Wales 1994–2007 Blackbird



BBS N. Ireland 1994–2007 Blackbird



Productivity trends


Table of productivity changes for Blackbird

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-	118	None				

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size								
Daily failure rate (eggs)	38	1968-2006	119	Curvilinear	2.57% nests/day	3.63% nests/day	-42%	
Daily failure rate (chicks)	38	1968-2006	154	Linear decline	3.03% nests/day	1.92% nests/day	-36.6%	
Laying date	38	1968-2006	143	None				
Juvenile to Adult ratio (CES)	22	1984-2006	98	Smoothed trend	122 Index value	100 Index value	-18%	
Juvenile to Adult ratio (CES)	10	1996-2006	109	Smoothed trend	89 Index value	100 Index value	12%	
Juvenile to Adult ratio (CES)	5	2001-2006	100	Smoothed trend	77 Index value	100 Index value	30%	


Clutch size 1966–2007

Blackbird




Brood size 1966–2007

Blackbird




Egg stage nest failure rate

Blackbird




Chick stage nest failure rate

Blackbird




Laying date 1966–2007

Blackbird



CES productivity 1983–2007

Blackbird



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

SONG THRUSH *Turdus philomelos*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK, England: rapid decline

UK population size

1,144,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

CBC/BBS shows a rapid decline in Song Thrush abundance that began in the mid 1970s. The second half of this decline can also be seen in the CES index. CES productivity showed an initial decrease, followed by some partial recovery, and NRS data indicate that nest success has improved since 1981. Changes in survival in the first winter, and perhaps also the post-fledging period, are sufficient to have caused the population decline ([Thomson et al. 1997](#), [Siriwardena et al. 1998a](#), [Robinson et al. 2004](#)). The environmental causes of these changes are not known, but changes in farming practices, land drainage, pesticides and predators are all possible contributors ([Fuller et al. 1995](#), [Robinson et al. 2004](#)). In woodland, drainage of damp ground and the depletion of woodland shrub layers through canopy closure and deer browsing may also be implicated ([Fuller et al. 2005](#)). Recent CBC/BBS data show a general increase, but population levels remain relatively low. Recovery of rural Song Thrush populations requires challenging new policy initiatives that should aim to restore nesting cover in scrub and woodland understorey, grazed grassland in arable-dominated areas, and damper soils in summer ([Peach et al. 2004](#)).



Population changes

CBC/BBS UK 1966–2007 Song Thrush




Table of population changes for Song Thrush

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	678	-51	-58	-43	>50	
	25	1981-2006	939	-16	-27	-5		
	10	1996-2006	1830	23	17	29		
	5	2001-2006	1975	4	0	6		
CBC/BBS England	39	1967-2006	540	-51	-60	-42	>50	
	25	1981-2006	740	-18	-30	-6		
	10	1996-2006	1422	24	19	29		
	5	2001-2006	1528	8	4	11		
CES adults	22	1984-2006	81	-17	-35	1		
	10	1996-2006	88	-2	-17	10		
	5	2001-2006	83	1	-12	12		
CES juveniles	22	1984-2006	66	-45	-59	-22	>25	
	10	1996-2006	75	20	0	38		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	10	1996-2006	1743	22	17	27		
	5	2001-2006	1975	5	2	9		
BBS England	11	1995-2006	1310	22	17	28		
	10	1996-2006	1347	23	19	29		
	5	2001-2006	1528	10	7	12		
BBS Scotland	11	1995-2006	148	20	2	41		
	10	1996-2006	151	19	3	39		
	5	2001-2006	161	0	-11	11		
BBS Wales	11	1995-2006	156	32	18	50		
	10	1996-2006	162	30	17	45		
	5	2001-2006	191	2	-5	9		
BBS N.Ireland	11	1995-2006	67	47	16	76		
	10	1996-2006	73	37	15	54		
	5	2001-2006	85	0	-13	13		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Song Thrush




CES adult abundance 1983—2007

Song Thrush




CES juvenile abundance 1983—2007

Song Thrush




BBS UK 1994—2007

Song Thrush




BBS England 1994—2007

Song Thrush




BBS Scotland 1994—2007

Song Thrush



BBS Wales 1994—2007

Song Thrush



BBS N. Ireland 1994—2007

Song Thrush






Table of productivity changes for Song Thrush

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	166	None				
Brood size	38	1968-2006	182	None				
Daily failure rate (eggs)	25	1981-2006	320	Curvilinear	4.37% nests/day	4.78% nests/day	9.4%	
Daily failure rate (chicks)	25	1981-2006	240	None				
Laying date	38	1968-2006	191	None				
Juvenile to Adult ratio (CES)	22	1984-2006	88	Smoothed trend	159 Index value	100 Index value	-37% >25	
Juvenile to Adult ratio (CES)	10	1996-2006	98	Smoothed trend	88 Index value	100 Index value	14%	
Juvenile to Adult ratio (CES)	5	2001-2006	93	Smoothed trend	99 Index value	100 Index value	1%	


Clutch size 1966–2007
Song Thrush




Brood size 1966–2007
Song Thrush




Egg stage nest failure rate
Song Thrush




Chick stage nest failure rate
Song Thrush



Laying date 1966–2007
Song Thrush



CES productivity 1983–2007
Song Thrush



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

MISTLE THRUSH

Turdus viscivorus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: amber (25–50% population decline)

Long-term trend

UK, England: moderate decline

UK population size

222,500 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

Like those of [Song Thrush](#) and [Blackbird](#), Mistle Thrush populations have declined significantly since the mid 1970s, especially on farmland. The species was recently moved from the green to the amber list because of population decline, and recent BBS data suggest that this decline is continuing. The Scottish BBS trend, in contrast to those elsewhere in the UK, is of strong increase since the late 1990s. There have been no trends in breeding performance, other than a minor increase in clutch size, and the decline is likely to have been driven by reduced annual survival ([Siriwardena et al. 1998](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007

Mistle Thrush




Table of population changes for Mistle Thrush

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	447	-43	-51	-31	>25	
	25	1981-2006	615	-38	-45	-31	>25	
	10	1996-2006	1175	-6	-11	2		
	5	2001-2006	1203	-12	-16	-7		
CBC/BBS England	39	1967-2006	367	-49	-57	-39	>25	
	25	1981-2006	500	-46	-52	-38	>25	
	10	1996-2006	939	-18	-22	-13		
	5	2001-2006	944	-18	-21	-14		
BBS UK	11	1995-2006	1088	-6	-13	2		
	10	1996-2006	1119	-6	-12	1		
	5	2001-2006	1203	-11	-14	-5		
BBS England	11	1995-2006	869	-19	-24	-13		
	10	1996-2006	890	-18	-22	-13		
	5	2001-2006	944	-17	-20	-13		
BBS Scotland	11	1995-2006	66	51	21	101		
	10	1996-2006	67	40	16	81		
	5	2001-2006	73	0	-12	18		
BBS Wales	11	1995-2006	95	2	-22	18		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS N.Ireland	11	1995-2006	54	23	-45	113		
	10	1996-2006	59	22	-25	60		
	5	2001-2006	70	-2	-19	18		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007

Mistle Thrush




BBS UK 1994–2007

Mistle Thrush




BBS England 1994–2007

Mistle Thrush




BBS Scotland 1994–2007

Mistle Thrush




BBS Wales 1994–2007

Mistle Thrush



BBS N. Ireland 1994–2007

Mistle Thrush




Productivity trends

Table of productivity changes for Mistle Thrush

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	35	Linear increase	3.88 eggs	4.09 eggs	5.4%	
Brood size	38	1968-2006	67	None				
Daily failure rate (eggs)	38	1968-2006	57	None				
Daily failure rate (chicks)	38	1968-2006	60	None				
Laying date	38	1968-2006	29	None				Small sample


Clutch size 1966–2007

Mistle Thrush




Brood size 1966–2007

Mistle Thrush




Egg stage nest failure rate

Mistle Thrush




Chick stage nest failure rate

Mistle Thrush



Laying date 1966—2007

Mistle Thrush



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

CETTI'S WARBLER *Cettia cetti*

- Population changes
- Productivity trends
- Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

England, Wales: rapid increase

UK population size

534 pairs in 1997–2001 (RBBP data: [BiE04](#)); mean of 645 pairs in 1998–2002 (RBBP data: [APEP06](#)); at least 1,331 singing males in 2005 ([Holling & RBBP 2008](#))

Status summary

Cetti's Warblers were first recorded in Britain as recently as 1961. Colonisation, which began in Kent in 1972 or 1973, continues to be monitored annually by [RBBP](#). Numbers and breeding range increased spectacularly during the first 12 years, with Norfolk and Dorset gradually overtaking Kent as the main host counties ([Gibbons *et al.* 1993](#), [Wotton *et al.* 1998](#)). Severe winters after 1978 led to the temporary extinction of the Kent population in 1988. Populations in milder regions continued to grow, but overall the UK population fell by over a third between 1984 and 1986. In the absence of severe winters since 1986, increase and range expansion have continued. For 2005, RBBP received reports of 1,331 singing males in 29 counties as far north as Anglesey and Norfolk ([Holling & RBBP 2008](#)). Much constant-effort ringing takes place in prime Cetti's Warbler habitat; despite the comparative rarity of this species, therefore, CES population and productivity indices are already available ([Robinson *et al.* 2007](#)). CES data confirm the species' sensitivity to cold winters, which appears to have become more evident as the breeding range has expanded into more testing climates. Numbers have risen widely in Europe since 1990 ([PECBMS 2007](#)).

Population changes

CES adult abundance 1986–2007 Cetti's Warbler





Table of population changes for Cetti's Warbler

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CES adults	5	2001-2006	10	46	-18	192		Small sample
CES juveniles	5	2001-2006	11	56	-3	231		Small sample

CES juvenile abundance 1986–2007 Cetti's Warbler



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GRASSHOPPER WARBLER

Locustella naevia

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: red (>50% population decline)

UK Biodiversity Action Plan: priority species

Long-term trend

UK: rapid decline

UK population size

11,750 pairs in 1990 (1988–91 Atlas: [APEP06](#)); 12,300 pairs in 2000 (updated using BBS trend: [BiE04](#))

Status summary

Grasshopper Warbler was previously amber-listed because of a contraction in range during the period preceding the 1988–91 Atlas, reportedly due to habitat loss ([Gibbons *et al.* 1993](#)). The CBC index suffered from small and severely dwindling sample sizes, but the available data indicate a rapid population decline between the mid 1960s and mid 1980s, when numbers became too small for annual monitoring ([Marchant *et al.* 1990](#)). On this basis, the species is now red-listed. The BBS shows wide fluctuations in abundance since 1994, and currently an overall moderate increase. Given suitable habitat and conditions, the species has high reproductive potential, as demonstrated by analysis of nest record data ([Glue 1990](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

BBS UK 1994–2007 Grasshopper Warbler




Table of population changes for Grasshopper Warbler

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	66	9	-23	35		
	10	1996-2006	68	-6	-32	13		
	5	2001-2006	68	8	-14	25		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

Productivity trends

No productivity information available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

SEDGE WARBLER *Acrocephalus schoenobaenus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK: moderate decline

UK population size

321,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary



The trend in England is apparently of moderate decline, but this is uncertain because the long-term changes are partly obscured by shorter fluctuations in numbers. Detailed analysis of BTO data sets has shown that much of the year-to-year variation in population size is driven by changes in adult survival rates which, in turn, are related to changes in rainfall on their wintering grounds, just south of the Sahara Desert, in the West African Sahel ([Peach et al. 1991](#)). The smoothed CBC/BBS and WBS trends show four troughs in population, related to years of poor West African rainfall, with a low point in 1984–85. The CES, which provides the biggest Sedge Warbler sample, shows the most recent three of the same troughs. Daily nest failure rates at the egg stage have halved. CES productivity data show a sustained decrease since the late 1980s. Sedge Warblers are estimated to have declined by 65% across Europe during 1980–2005 ([PECBMS 2007](#)).

Population changes




Table of population changes for Sedge Warbler

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	118	-28	-58	18		
	25	1981-2006	158	2	-20	38		
	10	1996-2006	297	-3	-18	11		
	5	2001-2006	283	-16	-29	-6		
CBC/BBS England	39	1967-2006	81	-39	-69	-13	>25	
	25	1981-2006	104	-10	-31	13		
	10	1996-2006	190	-12	-24	1		
	5	2001-2006	182	-19	-28	-11		
WBS waterways	31	1975-2006	43	-28	-56	24		
	25	1981-2006	46	-25	-47	10		
	10	1996-2006	47	-18	-37	8		
	5	2001-2006	36	-11	-24	4		
CES adults	22	1984-2006	64	-33	-49	-11	>25	
	10	1996-2006	74	-46	-54	-37	>25	
	5	2001-2006	64	-26	-34	-19	>25	


CES juv	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
		5	2001-2006	63	-27	-40	-11	>25	
BBS UK		11	1995-2006	270	-1	-17	15		
		10	1996-2006	278	-3	-18	10		
		5	2001-2006	283	-14	-26	-5		
BBS England		11	1995-2006	172	-12	-28	5		
		10	1996-2006	176	-13	-27	4		
		5	2001-2006	182	-16	-25	-6		
BBS Scotland		11	1995-2006	49	15	-19	57		
		10	1996-2006	49	9	-20	44		
		5	2001-2006	46	-14	-35	5		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Sedge Warbler




Waterways Bird Survey 1974—2007

Sedge Warbler




CES adult abundance 1983—2007

Sedge Warbler




CES juvenile abundance 1983—2007

Sedge Warbler




BBS UK 1994—2007

Sedge Warbler




BBS England 1994—2007

Sedge Warbler



BBS Scotland 1994—2007

Sedge Warbler




Productivity trends

Table of productivity changes for Sedge Warbler

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	37	Curvilinear	4.95 eggs	4.85 eggs	-2%	
Brood size	38	1968-2006	57	None				
Daily failure rate (eggs)	38	1968-2006	44	Linear decline	1.3% nests/day	0.61% nests/day	-53.1%	
Daily failure rate (chicks)	38	1968-2006	49	None				
Laying date	38	1968-2006	49	Curvilinear	May 29	May 23	-6 days	
Juvenile to Adult ratio (CES)	22	1984-2006	69	Smoothed trend	221 Index value	100 Index value	-55% >50	
Juvenile to Adult ratio (CES)	10	1996-2006	80	Smoothed trend	121 Index value	100 Index value	-17%	
Juvenile to Adult ratio (CES)	5	2001-2006	70	Smoothed trend	101 Index value	100 Index value	-1%	


Clutch size 1966–2007

Sedge Warbler




Brood size 1966–2007

Sedge Warbler




Egg stage nest failure rate

Sedge Warbler




Chick stage nest failure rate

Sedge Warbler




Laying date 1966–2007

Sedge Warbler



CES productivity 1983–2007

Sedge Warbler



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

REED WARBLER *Acrocephalus scirpaceus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK: uncertain

UK population size

60,800–122,000 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))



Status summary

This species has an unusually clumped distribution, with very high breeding concentrations in *Phragmites* reedbeds, where numbers are very hard to census. Because of this, CES, which has many sites in reedbeds, should perhaps be a better measure of population change than either CBC/BBS or WBS, where the species is encountered mainly at low density or in linear habitats. CES shows a decline from 1983 until the early 1990s, followed by a partial recovery, and another more recent decline. Both CBC/BBS and WBS show progressive moderate increases, however, perhaps linked to increasingly sensitive management of small and linear wetland sites. Population increase, as indicated by the census work, accords with the remarkable range expansion the species has achieved since the 1960s. West Wales, northwest and northeast England were colonised, as was the east coast of Ireland, between 1968–72 and 1988–91 ([Gibbons et al. 1993](#)), and the species probably now breeds sporadically as far north as the Tay reedbeds ([Robertson 2003](#)). Breeding performance as measured by brood size and failure rates has improved slightly, and a small improvement is apparent in CES productivity. The trend towards earlier laying can be partly explained by recent climate change ([Crick & Sparks 1999](#)).

Population changes

CES adult abundance 1983–2007 Reed Warbler




Table of population changes for Reed Warbler


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	52	106	39	263		
	25	1981-2006	70	84	31	201		
	10	1996-2006	125	27	7	51		
	5	2001-2006	125	7	-6	20		
CBC/BBS England	39	1967-2006	50	80	22	202		
	25	1981-2006	67	60	20	150		
	10	1996-2006	120	24	7	42		
	5	2001-2006	119	7	-3	20		
WBS waterways	25	1981-2006	22	55	5	151		
	10	1996-2006	26	23	-11	83		
	5	2001-2006	22	12	-9	45		
CES adults	22	1984-2006	52	-33	-46	-10	>25	
	10	1996-2006	60	-26	-36	-13	>25	
	5	2001-2006	52	-22	-31	-11		
CES juveniles	22	1984-2006	54	-28	-45	-4	>25	

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	107	28	9	50		
	10	1996-2006	111	23	7	43		
	5	2001-2006	125	9	0	19		
BBS England	11	1995-2006	103	25	6	50		
	10	1996-2006	106	20	5	42		
	5	2001-2006	119	9	-1	20		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS UK 1966—2007




CBC/BBS England 1966—2007




Waterways Bird Survey 1980—2007




CES juvenile abundance 1983—2007



BBS UK 1994—2007



BBS England 1994—2007




Productivity trends


Table of productivity changes for Reed Warbler

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	110	None				
Brood size	38	1968-2006	127	Linear increase	3.45 chicks	3.58 chicks	4%	
Daily failure rate (eggs)	38	1968-2006	144	None				
Daily failure rate (chicks)	38	1968-2006	110	Curvilinear	1.74% nests/day	0.5% nests/day	-71.3%	
Laying date	38	1968-2006	159	Curvilinear	Jun 16	Jun 10	-6 days	
Juvenile to Adult ratio (CES)	22	1984-2006	59	Smoothed trend	81 Index value	100 Index value	24%	
Juvenile to Adult ratio (CES)	10	1996-2006	67	Smoothed trend	108 Index value	100 Index value	-7%	
Juvenile to Adult ratio (CES)	5	2001-2006	58	Smoothed trend	109 Index value	100 Index value	-8%	


Clutch size 1966—2007




Brood size 1966—2007




Egg stage nest failure rate




Chick stage nest failure rate



Laying date 1966—2007



CES productivity 1983—2007



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

BLACKCAP

Sylvia atricapilla

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
UK: green

Long-term trend

UK, England: rapid increase

UK population size

932,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))



Status summary

Blackcap abundance has increased consistently since the late 1970s, a trend common to all habitats and evident from both the CBC/BBS and the CES indices, although the causes remain unknown. There have been no clear accompanying trends in productivity. The trend towards earlier laying may be a response to recent climate change ([Crick & Sparks 1999](#)). The more rapid increase in Scotland indicated by BBS suggests that climatic warming may be allowing this species to spread its range northwards. Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).

Population changes

CBC/BBS UK 1966–2007

Blackcap




Table of population changes for Blackcap

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	518	149	104	213		
	25	1981-2006	734	108	86	130		
	10	1996-2006	1431	48	41	54		
	5	2001-2006	1532	10	6	14		
CBC/BBS England	39	1967-2006	454	139	97	205		
	25	1981-2006	640	102	81	128		
	10	1996-2006	1236	44	37	51		
	5	2001-2006	1306	9	6	13		
CES adults	22	1984-2006	86	69	39	112		
	10	1996-2006	98	33	17	54		
	5	2001-2006	90	22	10	33		
CES juveniles	22	1984-2006	88	26	4	54		
	10	1996-2006	100	28	13	44		
	5	2001-2006	92	16	6	30		
BBS UK	11	1995-2006	1293	55	48	64		
	10	1996-2006	1343	45	37	51		
	5	2001-2006	1532	11	7	15		
BBS England	11	1995-2006	1118	48	41	55		
	10	1996-2006	1157	40	33	45		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Scotland	10	1996-2006	38	107	55	176		
	5	2001-2006	46	14	-3	42		
BBS Wales	11	1995-2006	109	56	30	83		
	10	1996-2006	115	41	21	63		
	5	2001-2006	134	11	2	22		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007

Blackcap




CES adult abundance 1983–2007

Blackcap




CES juvenile abundance 1983–2007

Blackcap




BBS UK 1994–2007

Blackcap




BBS England 1994–2007

Blackcap



BBS Wales 1994–2007

Blackcap




Productivity trends

Table of productivity changes for Blackcap

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	36	None				
Brood size	38	1968-2006	42	None				
Daily failure rate (eggs)	38	1968-2006	47	None				
Daily failure rate (chicks)	38	1968-2006	37	None				
Laying date	38	1968-2006	37	Curvilinear	May 20	May 11	-9 days	
Juvenile to Adult ratio (CES)	22	1984-2006	94	Smoothed trend	134 Index value	100 Index value	-25% >25	
Juvenile to Adult ratio (CES)	10	1996-2006	105	Smoothed trend	104 Index value	100 Index value	-3%	
Juvenile to Adult ratio (CES)	5	2001-2006	97	Smoothed trend	92 Index value	100 Index value	8%	


Clutch size 1966—2007

Blackcap




Brood size 1966—2007

Blackcap




Egg stage nest failure rate

Blackcap




Chick stage nest failure rate

Blackcap




Laying date 1966—2007

Blackcap



CES productivity 1983—2007

Blackcap



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

GARDEN WARBLER

Sylvia borin

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: shallow decline

UK population size

190,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Garden Warbler abundance has varied alongside that of other trans-Saharan migrant warblers ([Siriwardena et al. 1998b](#)), probably reflecting the influence of changes in their winter environment. Despite large short-term fluctuations in abundance, the CBC/BBS and CES now both suggest that the population is in long-term decline. There has been a substantial increase in nest losses at the chick stage, and post-fledging productivity, as measured by the CES, has declined sharply since 1983. Numbers have shown widespread moderate decline in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007

Garden Warbler




Table of population changes for Garden Warbler

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	195	-19	-47	25		
	25	1981-2006	264	16	-9	55		
	10	1996-2006	463	-18	-26	-8		
	5	2001-2006	415	-5	-13	5		
CBC/BBS England	39	1967-2006	165	-21	-46	13		
	25	1981-2006	221	6	-19	39		
	10	1996-2006	382	-20	-28	-11		
	5	2001-2006	339	-7	-16	2		
CES adults	22	1984-2006	63	-12	-39	20		
	10	1996-2006	68	-22	-36	-9		
	5	2001-2006	57	4	-10	19		
CES juveniles	22	1984-2006	63	-45	-59	-19	>25	
	10	1996-2006	67	-35	-47	-21	>25	
	5	2001-2006	60	1	-16	19		
BBS UK	11	1995-2006	406	-17	-27	-7		
	10	1996-2006	415	-19	-28	-9		
	5	2001-2006	415	-5	-14	3		
BBS England	11	1995-2006	333	-19	-28	-11		
	10	1996-2006	340	-21	-29	-15		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Wales	10	1996-2006	55	-28	-45	-7	>25	
	5	2001-2006	55	-8	-27	6		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Garden Warbler




CES adult abundance 1983—2007

Garden Warbler




CES juvenile abundance 1983—2007

Garden Warbler




BBS UK 1994—2007

Garden Warbler




BBS England 1994—2007

Garden Warbler



BBS Wales 1994—2007

Garden Warbler




Productivity trends


Table of productivity changes for Garden Warbler

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	16	None				Small sample
Brood size	38	1968-2006	24	None				Small sample
Daily failure rate (eggs)	38	1968-2006	22	None				Small sample
Daily failure rate (chicks)	38	1968-2006	19	Linear increase	1.05% nests/day	2.57% nests/day	144.8%	Small sample
Laying date	38	1968-2006	21	Curvilinear	May 25	May 17	-8 days	Small sample
Juvenile to Adult ratio (CES)	22	1984-2006	76	Smoothed trend	168 Index value	100 Index value	-41%	
Juvenile to Adult ratio (CES)	10	1996-2006	81	Smoothed trend	107 Index value	100 Index value	-7%	
Juvenile to Adult ratio (CES)	5	2001-2006	72	Smoothed trend	88 Index value	100 Index value	14%	


Clutch size 1966—2007




Brood size 1966—2007




Egg stage nest failure rate



Chick stage nest failure rate



Laying date 1966—2007



CES productivity 1983—2007



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

LESSER WHITETHROAT

Sylvia curruca

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK: fluctuating, with no long-term trend
England: shallow decline

UK population size

64,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Lesser Whitethroat abundance was roughly stable (albeit with short-term fluctuations) from the 1960s until the late 1980s, but the CBC/BBS and CES trends provide evidence for a subsequent moderate decline that lasted into the late 1990s. These changes were statistically significant, and large enough over the relevant periods to trigger BTO alerts. BBS has subsequently shown a significant sharp upturn, but this contrasts strongly with the continued decrease recorded by CES ringers. Wide fluctuations in productivity have been recorded by CES ringers, and may be influencing population change, but pressures during migration and in winter are the most likely causes of decline ([Fuller et al. 2005](#)).



Population changes

CBC/BBS UK 1966–2007

Lesser Whitethroat




Table of population changes for Lesser Whitethroat

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	117	-8	-37	37		
	25	1981-2006	155	-18	-35	4		
	10	1996-2006	256	8	-6	16		
	5	2001-2006	259	12	1	24		
CBC/BBS England	39	1967-2006	112	-14	-40	30		
	25	1981-2006	148	-22	-36	1		
	10	1996-2006	245	6	-8	15		
	5	2001-2006	247	11	1	23		
CES adults	22	1984-2006	40	-59	-78	-40	>50	
	10	1996-2006	37	-49	-61	-32	>25	
	5	2001-2006	33	-17	-35	1		
CES juveniles	22	1984-2006	44	-54	-78	-27	>50	
	10	1996-2006	42	-51	-62	-38	>50	
	5	2001-2006	40	-35	-46	-18	>25	
BBS UK	11	1995-2006	229	-8	-24	3		
	10	1996-2006	233	5	-12	15		
	5	2001-2006	259	19	8	33		


BBS Ex	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
			5 2001-2006	247	17	8	29		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Lesser Whitethroat




CES adult abundance 1983—2007

Lesser Whitethroat




CES juvenile abundance 1983—2007

Lesser Whitethroat




BBS UK 1994—2007

Lesser Whitethroat



BBS England 1994—2007

Lesser Whitethroat



Productivity trends

Table of productivity changes for Lesser Whitethroat

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Juvenile to Adult ratio (CES)	22	1984-2006	55	Smoothed trend	74 Index value	100 Index value	36%	
Juvenile to Adult ratio (CES)	10	1996-2006	53	Smoothed trend	79 Index value	100 Index value	27%	
Juvenile to Adult ratio (CES)	5	2001-2006	49	Smoothed trend	113 Index value	100 Index value	-12%	

Insufficient data on clutch size available for this species

Insufficient data on brood size available for this species


Insufficient data on nest failure available for this species

Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

CES productivity 1983—2007

Lesser Whitethroat



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WHITETHROAT *Sylvia communis*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: rapid decline, followed by shallow increase

UK population size

945,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Whitethroat populations had been stable for a few years up to 1968 but, despite a normal departure for their wintering grounds in West Africa, crashed by around 70% between the 1968 and 1969 breeding seasons. They fluctuated around their lower level until the mid 1980s, since when they have sustained a consistent shallow recovery. Recovery has been most apparent along linear waterways. In a pioneering study, [Winstanley *et al.* \(1974\)](#) linked the 1969 crash to droughts in the Whitethroat's wintering grounds in the western Sahel, just south of the Sahara Desert. Annual fluctuations in abundance, which are not shown in the smoothed trends, correlate to those in overwinter survival ([Baillie & Peach 1992](#)). Other trans-Saharan migrant warblers have shared similarly timed changes in abundance ([Siriwardena *et al.* 1998b](#)). Productivity, as measured by CES, rose during the 1980s and has since fluctuated and fallen back. It seems likely that habitat loss since the 1960s, particularly on farmland, will eventually limit the degree of recovery. A shallow upturn has been detected widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007 Whitethroat




Table of population changes for Whitethroat

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	447	-62	-73	-50	>50	
	25	1981-2006	627	65	30	110		
	10	1996-2006	1253	14	7	20		
	5	2001-2006	1304	7	2	10		
CBC/BBS England	39	1967-2006	388	-63	-73	-50	>50	
	25	1981-2006	543	66	32	113		
	10	1996-2006	1082	13	8	17		
	5	2001-2006	1120	4	1	7		
WBS waterways	31	1975-2006	41	171	15	367		
	25	1981-2006	45	375	169	613		
	10	1996-2006	53	60	17	96		
	5	2001-2006	43	32	11	50		
CES adults	22	1984-2006	59	-36	-57	-22	>25	
	10	1996-2006	68	-37	-46	-27	>25	
	5	2001-2006	61	-11	-22	4		


CES juv Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	65	-14	-28	0		
BBS UK	11	1995-2006	1158	19	11	26		
	10	1996-2006	1195	15	8	22		
	5	2001-2006	1304	7	2	10		
BBS England	11	1995-2006	1000	17	11	23		
	10	1996-2006	1032	14	8	19		
	5	2001-2006	1120	5	1	8		
BBS Scotland	11	1995-2006	66	87	41	155		
	10	1996-2006	68	74	32	122		
	5	2001-2006	76	26	7	51		
BBS Wales	11	1995-2006	74	-22	-38	-5		
	10	1996-2006	77	-26	-39	-12	>25	
	5	2001-2006	86	-5	-17	7		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Whitethroat




Waterways Bird Survey 1974—2007

Whitethroat




CES adult abundance 1983—2007

Whitethroat




CES juvenile abundance 1983—2007

Whitethroat




BBS UK 1994—2007

Whitethroat




BBS England 1994—2007

Whitethroat




BBS Scotland 1994—2007

Whitethroat



BBS Wales 1994—2007

Whitethroat




Productivity trends


Table of productivity changes for Whitethroat

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	28	None				Small sample
Brood size	38	1968-2006	62	None				
Daily failure rate (eggs)	38	1968-2006	41	None				
Daily failure rate (chicks)	38	1968-2006	48	None				
Laying date	38	1968-2006	18	Curvilinear	May 26	May 19	-7 days	Small sample
Juvenile to Adult ratio (CES)	22	1984-2006	73	Smoothed trend	93 Index value	100 Index value	8%	
Juvenile to Adult ratio (CES)	10	1996-2006	83	Smoothed trend	133 Index value	100 Index value	-25%	
Juvenile to Adult ratio (CES)	5	2001-2006	73	Smoothed trend	139 Index value	100 Index value	-28%	


**Clutch size 1966—2007
Whitethroat**




**Brood size 1966—2007
Whitethroat**




**Egg stage nest failure rate
Whitethroat**




**Chick stage nest failure rate
Whitethroat**



**Laying date 1966—2007
Whitethroat**



**CES productivity 1983—2007
Whitethroat**



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WOOD WARBLER

Phylloscopus sibilatrix

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 2 (declining)
 UK: amber (25–50% population decline)
 UK Biodiversity Action Plan: priority species

Long-term trend

UK: decline

UK population size

17,200 (15,830–18,570) males in 1984–85 (Bibby 1989: APEP06); 9,000–10,500 pairs in 2000 (updated using BBS trend: BiE04)

Status summary

Wood Warblers, which have a westerly distribution in Britain, were covered relatively poorly until BBS began. Little change was apparent at the few CBC plots on which the species occurred (Marchant *et al.* 1990, Crick *et al.* 1998). The species' breeding range varied little between the two atlas periods (Gibbons *et al.* 1993), but has subsequently retreated heavily from lowland England. BBS shows a rapid and significant decline since 1994, and accordingly the species has been moved from the green to the amber list. Nest success has apparently improved considerably at the egg stage, although nest record samples are small. Numbers have fallen widely in Europe since 1980 (PECBMS 2007). With declines evident across northern and western Europe, this previously 'secure' species is now provisionally categorised as 'declining' (BirdLife International 2004).



Population changes

BBS UK 1994–2007

Wood Warbler




Table of population changes for Wood Warbler

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	53	-58	-70	-44	>50	
	10	1996-2006	53	-53	-67	-36	>50	
	5	2001-2006	47	-16	-44	13		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

Productivity trends


Table of productivity changes for Wood Warbler

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	16	None				Small sample
Brood size	38	1968-	36	None				

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Daily failure rate (eggs)								
Daily failure rate (chicks)		2006						sample
Laying date	38	1968-2006	31	Curvilinear	May 23	May 18	-5 days	


Clutch size 1966—2007

Wood Warbler




Brood size 1966—2007

Wood Warbler




Egg stage nest failure rate

Wood Warbler




Chick stage nest failure rate

Wood Warbler



Laying date 1966—2007

Wood Warbler



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

CHIFFCHAFF *Phylloscopus collybita*

• Population changes

• Productivity trends

• Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK, England: shallow increase

UK population size

807,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Chiffchaff abundance crashed in the late 1960s/early 1970s in common with that of other trans-Saharan warblers ([Siriwardena et al. 1998a](#)). After remaining stable for a decade, the population recovered strongly, and has continued to increase. This recovery is evident from both CBC/BBS and CES data. Climate change may partly explain the strong trend towards earlier laying ([Crick & Sparks 1999](#)). Overwinter survival may be the critical factor responsible for changes in abundance, as it is for [Whitethroat](#) and [Sedge Warbler](#). Productivity as measured by CES has decreased as the population has risen. Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).

Population changes

CBC/BBS UK 1966–2007 Chiffchaff




Table of population changes for Chiffchaff

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	464	25	4	65		
	25	1981-2006	659	112	86	149		
	10	1996-2006	1305	24	19	31		
	5	2001-2006	1425	7	3	11		
CBC/BBS England	39	1967-2006	397	31	4	72		
	25	1981-2006	560	120	95	160		
	10	1996-2006	1102	27	22	35		
	5	2001-2006	1192	6	3	10		
CES adults	22	1984-2006	67	161	75	304		
	10	1996-2006	81	25	2	64		
	5	2001-2006	79	21	5	39		
CES juveniles	22	1984-2006	77	137	59	292		
	10	1996-2006	91	19	-3	47		
	5	2001-2006	85	4	-7	17		
BBS UK	11	1995-2006	1182	29	24	37		
	10	1996-2006	1226	25	20	32		
	5	2001-2006	1425	9	4	12		
BBS England	11	1995-2006	997	32	26	40		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Scotland	10	1996-2006	31	130	50	257		
	5	2001-2006	39	23	-13	58		
BBS Wales	11	1995-2006	123	13	0	28		
	10	1996-2006	128	12	-1	24		
	5	2001-2006	154	13	4	20		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007

Chiffchaff




CES adult abundance 1983–2007

Chiffchaff




CES juvenile abundance 1983–2007

Chiffchaff




BBS UK 1994–2007

Chiffchaff




BBS England 1994–2007

Chiffchaff



BBS Wales 1994–2007

Chiffchaff




Productivity trends

Table of productivity changes for Chiffchaff

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	30	None				Small sample
Brood size	38	1968-2006	33	Linear decline	5.12 chicks	4.74 chicks	-7.4%	
Daily failure rate (eggs)	38	1968-2006	39	None				
Daily failure rate (chicks)	38	1968-2006	35	None				
Laying date	38	1968-2006	44	Linear decline	May 17	May 3	-14 days	
Juvenile to Adult ratio (CES)	22	1984-2006	85	Smoothed trend	131 Index value	100 Index value	-24%	
Juvenile to Adult ratio (CES)	10	1996-2006	98	Smoothed trend	140 Index value	100 Index value	-29% >25	
Juvenile to Adult ratio (CES)	5	2001-2006	90	Smoothed trend	130 Index value	100 Index value	-23%	


Clutch size 1966–2007

Chiffchaff




Brood size 1966–2007

Chiffchaff




Egg stage nest failure rate

Chiffchaff




Chick stage nest failure rate

Chiffchaff




Laying date 1966–2007

Chiffchaff



CES productivity 1983–2007

Chiffchaff



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

WILLOW WARBLER *Phylloscopus trochilus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: amber (25–50% population decline)

Long-term trend

England: rapid decline

UK population size

2,125,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Willow Warbler abundance has apparently shown different trends at different UK latitudes. The overall CBC/BBS trend shows a rapid decline during the 1980s and early 1990s, after 20 years of relative stability, and, on the strength of a 31% decline on CBC plots between 1974 and 1999, the species was moved from the green to the amber list. This decline occurred mainly in southern Britain, however, accompanied by a fall in survival rates there ([Peach *et al.* 1995a](#)), with Scottish populations remaining unaffected. BBS figures since 1994 indicate a stark contrast between an initially upward trend in Scotland and in Northern Ireland, and continued severe decreases in England and in Wales. Pressures on migration and in the winter are likely to be affecting the population, as is a reduction in habitat quality on the breeding grounds ([Fuller *et al.* 2005](#)). The recent population decline is associated with a moderate decline in productivity as measured by CES and with a substantial increase in failure rates at the egg stage, which raises NRS concern ([Leech & Barimore 2008](#)). Average laying dates have become a week earlier, perhaps in response to recent climatic warming ([Crick & Sparks 1999](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS England 1966–2007 Willow Warbler




Table of population changes for Willow Warbler


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	399	-58	-70	-46	>50	
	25	1981-2006	532	-58	-67	-49	>50	
	10	1996-2006	938	-30	-36	-24	>25	
	5	2001-2006	820	-11	-18	-7		
CES adults	22	1984-2006	88	-61	-70	-52	>50	
	10	1996-2006	91	-48	-54	-42	>25	
	5	2001-2006	77	-7	-14	1		
CES juveniles	22	1984-2006	89	-72	-78	-63	>50	
	10	1996-2006	98	-53	-62	-45	>50	
	5	2001-2006	86	-10	-19	1		
BBS UK	11	1995-2006	1275	-14	-20	-7		
	10	1996-2006	1293	-17	-23	-10		
	5	2001-2006	1260	-8	-13	-5		
BBS England	11	1995-2006	866	-32	-37	-27	>25	
	10	1996-2006	873	-31	-36	-26	>25	

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Scotland	10	1996-2006	183	-5	-18	5		
	5	2001-2006	180	-7	-14	-1		
BBS Wales	11	1995-2006	152	-27	-37	-14	>25	
	10	1996-2006	157	-26	-36	-14	>25	
	5	2001-2006	170	-11	-19	-5		
BBS N.Ireland	11	1995-2006	70	36	5	56		
	10	1996-2006	76	12	-9	25		
	5	2001-2006	86	-4	-16	8		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CES adult abundance 1983–2007
Willow Warbler




CES juvenile abundance 1983–2007
Willow Warbler




BBS UK 1994–2007
Willow Warbler




BBS England 1994–2007
Willow Warbler




BBS Scotland 1994–2007
Willow Warbler



BBS Wales 1994–2007
Willow Warbler



BBS N. Ireland 1994–2007
Willow Warbler



Productivity trends


Table of productivity changes for Willow Warbler

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	49	None				
Brood size	38	1968-2006	131	None				
Daily failure rate (eggs)	38	1968-2006	69	Linear increase	0.93% nests/day	1.66% nests/day	78.5%	
Daily failure rate (chicks)	38	1968-2006	120	None				
Laying date	38	1968-2006	84	Linear decline	May 20	May 14	-6 days	
Juvenile to Adult ratio (CES)	22	1984-2006	95	Smoothed trend	144 Index value	100 Index value	-31% >25	
Juvenile to Adult ratio (CES)	10	1996-2006	103	Smoothed trend	112 Index value	100 Index value	-11%	
Juvenile to Adult ratio	5	2001-	90	Smoothed	109 Index	100 Index	-8%	

(CES)	Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
-------	----------	--------------	-------	--------------------	-------	------------------------	------------------	--------	---------


Clutch size 1966–2007

Willow Warbler




Brood size 1966–2007

Willow Warbler




Egg stage nest failure rate

Willow Warbler




Chick stage nest failure rate

Willow Warbler




Laying date 1966–2007

Willow Warbler



CES productivity 1983–2007

Willow Warbler



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

GOLDCREST *Regulus regulus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)
UK: amber (>50% population decline, but data possibly unrepresentative)

Long-term trend

England: fluctuating, with no long-term trend

UK population size

842,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Goldcrest abundance is unusually severely affected by winter weather, and the strong increase in the species' CBC/BBS index up to the mid 1970s can be interpreted as recovery from the cold winters of the early 1960s. The subsequent decline has resulted in the recent addition of the species to the amber list, although it meets the criterion only because 1975, at the start of the relevant 25-year period, was the peak year of the population index. Trends over longer and shorter periods all suggest population increase, and the long-term trend looks very much like a series of damped oscillations following the 1962/63 winter. The high amplitude of year-to-year change reflects the species high breeding potential, and its sensitivity to cold winter weather. BBS has recorded substantial increases in all UK countries except Wales, where a significant decline has been registered. CBC had relatively poor coverage of conifer plantations, in which Goldcrests occur at increasing densities as the trees mature. The increase in area of prime habitat has therefore been poorly reflected in the long-term trend.



Population changes

CBC/BBS England 1966–2007 Goldcrest




Table of population changes for Goldcrest

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	206	42	-20	197		
	25	1981-2006	278	-18	-38	14		
	10	1996-2006	528	22	11	35		
	5	2001-2006	588	-2	-11	6		
BBS UK	11	1995-2006	678	25	7	40		
	10	1996-2006	701	16	1	27		
	5	2001-2006	817	-8	-12	0		
BBS England	11	1995-2006	475	27	15	42		
	10	1996-2006	491	22	9	35		
	5	2001-2006	588	0	-10	8		
BBS Scotland	11	1995-2006	79	57	24	82		
	10	1996-2006	79	37	9	53		
	5	2001-2006	85	-8	-17	4		
BBS Wales	11	1995-2006	80	-38	-58	3		
	10	1996-2006	83	-38	-56	-7	>25	


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS N.Ireland	10	1996-2006	42	45	3	64		
	5	2001-2006	45	7	-10	27		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS UK 1994—2007

Goldcrest




BBS England 1994—2007

Goldcrest




BBS Scotland 1994—2007

Goldcrest



BBS Wales 1994—2007

Goldcrest



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

SPOTTED FLYCATCHER

Muscicapa striata

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3, declining

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK, England: rapid decline

UK population size

63,700 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary



Spotted Flycatchers have declined rapidly and consistently since the 1960s according to census data, and the decline is also reflected in the trend revealed by CES. Productivity measures indicate lower clutch and brood sizes and greater nest losses at the egg and chick stages, which raise NRS concern ([Leech & Barimore 2008](#)), and a decrease overall in the ratio of juveniles to adults. Demographic modelling shows that decreases in the annual survival rates of birds in their first year of life are most likely to have driven the decline ([Freeman & Crick 2003](#)). Decreasing survival rates may have been caused by deteriorations in woodland quality, particularly leading to declines in the large flying insects that are food to the flycatcher, or by conditions either on the wintering grounds or along migration routes ([Fuller et al. 2005](#)). Since trends have been similar across UK regions and habitats, however, it is more likely that the decline has been driven by factors operating outside the UK. Spotted Flycatchers are estimated to have declined by 59% across Europe during 1980–2005 ([PECBMS 2007](#)). A predator 'control' experiment has indicated that the abundance of nest predators may be determining the breeding success of Spotted Flycatchers, especially in woodland, where nest success was lower overall than in gardens ([Stoate & Szczur 2006](#)). Another study using nest cameras has identified avian predators, especially [Jays](#), as responsible for most nest losses ([Stevens et al. 2008](#)).

Population changes

CBC/BBS UK 1966–2007 Spotted Flycatcher




Table of population changes for Spotted Flycatcher

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	121	-86	-90	-80	>50	
	25	1981-2006	142	-81	-86	-75	>50	
	10	1996-2006	222	-32	-38	-16	>25	
	5	2001-2006	196	-18	-27	-3		
CBC/BBS England	39	1967-2006	93	-87	-92	-82	>50	
	25	1981-2006	105	-82	-87	-75	>50	
	10	1996-2006	159	-32	-44	-16	>25	
	5	2001-2006	136	-23	-32	-10		
CES adults	22	1984-2006	15	-30	-75	32		Small sample
	10	1996-2006	13	13	-51	79		Small sample
	5	2001-2006	13	31	-24	94		Small sample
CES juveniles	22	1984-2006	11	-56	-86	-15	>50	Small sample


BBS UK Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	196	-20	-29	-2		
BBS England	11	1995-2006	144	-37	-48	-21	>25	
	10	1996-2006	146	-34	-44	-20	>25	
	5	2001-2006	136	-22	-32	-11		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Spotted Flycatcher




CES adult abundance 1983—2007

Spotted Flycatcher




CES juvenile abundance 1983—2007

Spotted Flycatcher




BBS UK 1994—2007

Spotted Flycatcher



BBS England 1994—2007

Spotted Flycatcher




Productivity trends

Table of productivity changes for Spotted Flycatcher

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	81	Curvilinear	4.22 eggs	4.15 eggs	-1.7%	
Brood size	38	1968-2006	130	Curvilinear	3.61 chicks	3.6 chicks	-0.3%	
Daily failure rate (eggs)	38	1968-2006	122	Curvilinear	1.77% nests/day	1.88% nests/day	6.2%	
Daily failure rate (chicks)	38	1968-2006	109	Linear increase	0.98% nests/day	1.43% nests/day	45.9%	
Laying date	38	1968-2006	72	None				
Juvenile to Adult ratio (CES)	22	1984-2006	21	Smoothed trend	133 Index value	100 Index value	-25%	
Juvenile to Adult ratio (CES)	10	1996-2006	18	Smoothed trend	119 Index value	100 Index value	-16%	Small sample
Juvenile to Adult ratio (CES)	5	2001-2006	16	Smoothed trend	149 Index value	100 Index value	-33%	Small sample


Clutch size 1966—2007

Spotted Flycatcher




Brood size 1966—2007

Spotted Flycatcher




Egg stage nest failure rate

Spotted Flycatcher




Chick stage nest failure rate

Spotted Flycatcher




Laying date 1966—2007

Spotted Flycatcher



CES productivity 1983—2007

Spotted Flycatcher



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

PIED FLYCATCHER

Ficedula hypoleuca

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK: possible decline

UK population size

35,000–40,000 pairs in 1990 (1988–91 Atlas: [APEP06](#)); 29,500–33,800 pairs in 2000 (updated using BBS trend: [BI04](#))

Status summary

Pied Flycatchers are restricted to upland deciduous woods in parts of western and northern Britain. The proportions of CBC plots occupied rose during the 1980s, but the species was never numerous enough for trends to be estimated ([Marchant *et al.* 1990](#)). The 1988–91 breeding atlas revealed a small expansion in range since 1968–72, aided by the provision of nest boxes in new areas ([Gibbons *et al.* 1993](#)). BBS suggests that abundance has decreased steeply since 1994, raising a BTO alert. Percentage nestbox occupancy has also fallen over a similar period at a number of sites monitored as [RAS](#) projects. Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)). The reasons for this decline are unknown. In the Netherlands, climate change may have brought about decline in Pied Flycatchers by advancing the peak period of food availability for this species in deciduous forests – the birds being unable so far to compensate for the change in food supply by breeding earlier ([Both 2002](#), [Both *et al.* 2006](#)).



Population changes

BBS UK 1994–2007

Pied Flycatcher




Table of population changes for Pied Flycatcher

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	41	-50	-63	-38	>25	
	10	1996-2006	41	-49	-61	-37	>25	



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

Productivity trends

Information on productivity not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

LONG-TAILED TIT *Aegithalos caudatus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

England: moderate increase

UK population size

273,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

This species undergoes wide fluctuations in numbers between breeding seasons, suffering heavy mortality when winters are severe, but is able to recover quickly by virtue of its high breeding potential. Numbers were low after the severe winters of the early 1960s and again during a series of relatively cold winters beginning in the late 1970s. The starting years of the 25-year and longest monitoring periods coincided with troughs in population, thus exaggerating the long-term trend. Both CBC/BBS and CES index trends show progressive increases in Long-tailed Tit abundance beginning in the mid 1980s, but tailing off in recent years. Clutch and brood sizes have become smaller since the 1960s and, curiously, nest losses have switched from the egg to the chick stage. The marked trend towards earlier laying may be explained by recent climatic changes ([Crick & Sparks 1999](#)).



Population changes

CBC/BBS England 1966–2007 Long-tailed Tit




Table of population changes for Long-tailed Tit

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	302	77	28	168		
	25	1981-2006	414	67	36	103		
	10	1996-2006	762	3	-4	10		
	5	2001-2006	795	-3	-8	3		
CES adults	22	1984-2006	76	26	1	75		
	10	1996-2006	89	-12	-23	-1		
	5	2001-2006	82	-14	-27	-2		
CES juveniles	22	1984-2006	69	11	-24	71		
	10	1996-2006	83	-2	-13	13		
	5	2001-2006	76	-17	-28	-4		
BBS UK	11	1995-2006	772	1	-7	9		
	10	1996-2006	795	1	-7	8		
	5	2001-2006	903	-4	-11	1		
BBS England	11	1995-2006	678	0	-9	6		
	10	1996-2006	698	3	-6	9		
	5	2001-2006	795	-1	-8	4		
BBS Wales	11	1995-2006	54	4	-19	28		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CES adult abundance 1983–2007

Long-tailed Tit




CES juvenile abundance 1983–2007

Long-tailed Tit




BBS UK 1994–2007

Long-tailed Tit




BBS England 1994–2007

Long-tailed Tit



BBS Wales 1994–2007

Long-tailed Tit




Productivity trends

Table of productivity changes for Long-tailed Tit

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	34	Linear decline	7.63 eggs	6.46 eggs	-15.3%	
Brood size	38	1968-2006	28	Curvilinear	6.68 chicks	6.17 chicks	-7.6%	Small sample
Daily failure rate (eggs)	38	1968-2006	52	Linear decline	3.58% nests/day	0.84% nests/day	-76.5%	
Daily failure rate (chicks)	38	1968-2006	37	Linear increase	0.8% nests/day	1.71% nests/day	113.8%	
Laying date	38	1968-2006	44	Linear decline	Apr 21	Apr 6	-15 days	
Juvenile to Adult ratio (CES)	22	1984-2006	83	Smoothed trend	105 Index value	100 Index value	-5%	
Juvenile to Adult ratio (CES)	10	1996-2006	97	Smoothed trend	86 Index value	100 Index value	16%	
Juvenile to Adult ratio (CES)	5	2001-2006	90	Smoothed trend	99 Index value	100 Index value	1%	


Clutch size 1966–2007

Long-tailed Tit




Brood size 1966–2007

Long-tailed Tit




Egg stage nest failure rate

Long-tailed Tit




Chick stage nest failure rate

Long-tailed Tit




Laying date 1966–2007

Long-tailed Tit



CES productivity 1983–2007

Long-tailed Tit



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

BLUE TIT *Cyanistes caeruleus*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: shallow increase

UK population size

3,535,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))



Status summary

Blue Tit populations have increased in abundance, in parallel with those of [Great Tits](#), with brief pauses in the long-term upward trend. The recent years of the CBC/BBS index show fluctuations but the trend still appears to be upward. Food provision in gardens during winter and availability of nest boxes, which may reduce egg and nestling predation, have both increased and may have contributed to the rise in population. Decreasing clutch and brood sizes, and a substantial decline in the proportion of young birds in early autumn, have accompanied the population increase. Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).

Population changes

CBC/BBS UK 1966–2007 Blue Tit





Table of population changes for Blue Tit


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	780	42	27	60		
	25	1981-2006	1098	23	15	33		
	10	1996-2006	2157	7	4	11		
	5	2001-2006	2264	8	5	11		
CBC/BBS England	39	1967-2006	642	40	25	57		
	25	1981-2006	898	19	10	29		
	10	1996-2006	1746	6	2	8		
	5	2001-2006	1814	6	4	9		
CES adults	22	1984-2006	95	16	0	40		
	10	1996-2006	105	-2	-12	7		
	5	2001-2006	96	13	3	25		
CES juveniles	22	1984-2006	94	-42	-53	-23	>25	
	10	1996-2006	106	-35	-43	-25	>25	
	5	2001-2006	97	-15	-23	-2		
BBS UK	11	1995-2006	1995	11	7	14		
	10	1996-2006	2056	8	5	11		
	5	2001-2006	2264	6	4	9		
BBS England	11	1995-2006	1614	8	4	12		

Clutch	Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size									
	Daily failure rate (eggs)	38	1968-2006	259	Linear decline	0.46% nests/day	0.25% nests/day	-45.7%	
	Daily failure rate (chicks)	38	1968-2006	194	Linear increase	0.65% nests/day	0.87% nests/day	33.8%	
	Laying date	38	1968-2006	197	Linear decline	May 3	Apr 26	-7 days	
	Juvenile to Adult ratio (CES)	22	1984-2006	99	Smoothed trend	218 Index value	100 Index value	-54% >50	
	Juvenile to Adult ratio (CES)	10	1996-2006	109	Smoothed trend	164 Index value	100 Index value	-39% >25	
	Juvenile to Adult ratio (CES)	5	2001-2006	100	Smoothed trend	121 Index value	100 Index value	-18%	


Clutch size 1966–2007
Blue Tit




Brood size 1966–2007
Blue Tit




Egg stage nest failure rate
Blue Tit




Chick stage nest failure rate
Blue Tit



Laying date 1966–2007
Blue Tit



CES productivity 1983–2007
Blue Tit



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

GREAT TIT

Parus major

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK, England: rapid increase

UK population size

2,074,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Great Tits have increased steadily since the 1960s, with the exception of two brief periods of stability or shallow decline during the mid 1970s and late 1980s. Recent CBC/BBS and BBS results suggest that this increase is continuing, in all UK countries. More widespread food provision in gardens during winter is one possible explanation for the increase. Changes in different aspects of breeding performance are contradictory: CES productivity has fluctuated, brood size has decreased, and nest success has improved at the egg stage but decreased at the chick stage. Laying date has advanced by about a week in the UK, in line with climatic change. In a Dutch study population, however, the breeding period did not advance during 1973–95 and became increasingly mistimed with respect to the peak of insect abundance ([Visser et al. 1998](#)).



Population changes

CBC/BBS UK 1966–2007

Great Tit





Table of population changes for Great Tit


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	733	110	90	142		
	25	1981-2006	1031	53	40	67		
	10	1996-2006	2016	38	33	44		
	5	2001-2006	2146	23	21	27		
CBC/BBS England	39	1967-2006	605	104	76	130		
	25	1981-2006	845	49	34	61		
	10	1996-2006	1635	38	34	43		
	5	2001-2006	1724	22	20	26		
CES adults	22	1984-2006	89	40	11	72		
	10	1996-2006	101	34	16	51		
	5	2001-2006	94	43	26	61		
CES juveniles	22	1984-2006	91	4	-17	39		
	10	1996-2006	103	15	-2	36		
	5	2001-2006	96	30	18	44		
BBS UK	11	1995-2006	1858	43	38	48		
	10	1996-2006	1917	38	33	43		
	5	2001-2006	2146	22	19	25		
BBS England	11	1995-2006	1505	41	36	45		

Clutch	Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
	Daily failure rate (eggs)	38	1968-2006	265	Curvilinear	0.68% nests/day	0.43% nests/day	-36.8%	
	Daily failure rate (chicks)	38	1968-2006	197	Linear increase	0.57% nests/day	0.83% nests/day	45.6%	
	Laying date	38	1968-2006	176	Linear decline	May 4	Apr 27	-7 days	
	Juvenile to Adult ratio (CES)	22	1984-2006	97	Smoothed trend	146 Index value	100 Index value	-32% >25	
	Juvenile to Adult ratio (CES)	10	1996-2006	109	Smoothed trend	119 Index value	100 Index value	-16%	
	Juvenile to Adult ratio (CES)	5	2001-2006	100	Smoothed trend	102 Index value	100 Index value	-2%	


Clutch size 1966–2007
Great Tit




Brood size 1966–2007
Great Tit




Egg stage nest failure rate
Great Tit




Chick stage nest failure rate
Great Tit



Laying date 1966–2007
Great Tit



CES productivity 1983–2007
Great Tit



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

COAL TIT *Pariparus ater*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

England: probable moderate increase

UK population size

653,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary



While other common tit species have increased, the UK Coal Tit population has been rather stable since the mid 1970s, following earlier rapid increase. The ratios of Coal Tit to [Blue](#) and [Great Tits](#) caught for ringing have both shown a sustained increase since 1960 ([Perrins 2003](#)), however, although in these figures population change may be confounded to some degree with changes in behaviour among birds and bird ringers. Confidence intervals are wide, but BBS shows large changes in population sizes that have varied geographically across the UK. This pattern suggests that Coal Tit abundance in the UK may be controlled by a complex range of factors.

Population changes

CBC/BBS England 1966–2007 Coal Tit




Table of population changes for Coal Tit

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	207	50	-17	188		
	25	1981-2006	274	6	-20	44		
	10	1996-2006	491	9	-6	32		
	5	2001-2006	527	0	-10	11		
BBS UK	11	1995-2006	663	6	-5	18		
	10	1996-2006	687	2	-8	12		
	5	2001-2006	791	2	-5	8		
BBS England	11	1995-2006	435	12	-6	39		
	10	1996-2006	451	9	-8	31		
	5	2001-2006	527	1	-10	11		
BBS Scotland	11	1995-2006	105	-1	-18	15		
	10	1996-2006	105	-6	-22	10		
	5	2001-2006	112	3	-10	18		
BBS Wales	11	1995-2006	68	-26	-46	-1	>25	
	10	1996-2006	71	-24	-44	-3		
	5	2001-2006	82	-11	-29	5		
BBS N.Ireland	11	1995-2006	53	87	40	137		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS UK 1994—2007

Coal Tit




BBS England 1994—2007

Coal Tit




BBS Scotland 1994—2007

Coal Tit




BBS Wales 1994—2007

Coal Tit



BBS N. Ireland 1994—2007

Coal Tit



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

WILLOW TIT *Poecile montana*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: red (>50% population decline)
UK Biodiversity Action Plan: priority species

Long-term trend

UK, England: rapid decline

UK population size

8,500 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Willow Tits have been in decline since the mid 1970s, and have become extinct in an ever-growing number of former haunts. The continuing decline in the CBC/BBS index through the 1990s, following a brief period of stability during the 1980s, is replicated in the CES abundance trend. The UK conservation listing has recently been upgraded from amber to red. Numbers have changed least in the wet woodlands that the species prefers ([Siriwardena 2004](#)). Farmland is now only rarely occupied. The most likely causes of decline are competition with other tit species, increasing nest predation by [Great Spotted Woodpeckers](#), and deterioration in the quality of woodland as feeding habitat for Willow Tits through canopy closure and increased browsing by deer ([Perrins 2003](#), [Siriwardena 2004](#), [Fuller et al. 2005](#)). A study of former CBC sites and other woods that were known to have held the species in the past found that the sites still holding Willow Tits tended to be wetter but did not differ in the density of potential nest predators or avian competitors ([Lewis et al. 2007](#)). Willow Tit is one of the most strongly declining bird species in Europe, having decreased at an annual rate of 4% during 1980–2005, but has declined to a lesser extent in central and east Europe than in the north, west and south ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007 Willow Tit




Table of population changes for Willow Tit

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	43	-85	-93	-72	>50	
	25	1981-2006	44	-82	-90	-69	>50	
	10	1996-2006	60	-64	-72	-51	>50	
	5	2001-2006	47	-33	-47	-14	>25	
CBC/BBS England	39	1967-2006	40	-83	-92	-70	>50	
	25	1981-2006	40	-82	-91	-72	>50	
	10	1996-2006	53	-62	-71	-51	>50	
	5	2001-2006	40	-34	-51	-17	>25	
CES adults	22	1984-2006	20	-58	-89	-25	>50	
	10	1996-2006	15	-49	-82	-4	>25	Small sample
CES juveniles	22	1984-2006	28	-58	-78	-35	>50	
	10	1996-2006	21	-55	-72	-38	>50	
	5	2001-2006	14	-20	-53	17		Small sample
BBS UK	11	1995-2006	54	-66	-74	-54	>50	


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS England	11	1995-2006	47	-66	-75	-57	>50	
	10	1996-2006	46	-63	-72	-54	>50	
	5	2001-2006	40	-36	-52	-19	>25	



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Willow Tit




CES adult abundance 1983—2007

Willow Tit




CES juvenile abundance 1983—2007

Willow Tit




BBS UK 1994—2007

Willow Tit



BBS England 1994—2007

Willow Tit



Productivity trends

Table of productivity changes for Willow Tit

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Juvenile to Adult ratio (CES)	22	1984-2006	31	Smoothed trend	118 Index value	100 Index value	-15%	
Juvenile to Adult ratio (CES)	10	1996-2006	24	Smoothed trend	87 Index value	100 Index value	15%	
Juvenile to Adult ratio (CES)	5	2001-2006	16	Smoothed trend	88 Index value	100 Index value	13%	Small sample

Insufficient data on clutch size available for this species

Insufficient data on brood size available for this species

Insufficient data on nest failure available for this species

Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)



MARSH TIT *Poecile palustris*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3, declining
UK: red (>50% population decline)
UK Biodiversity Action Plan: priority species

Long-term trend

UK, England: rapid decline

UK population size

52,800 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP2](#))

Status summary

Marsh Tit abundance has declined almost continuously since BTO monitoring began. The species' UK conservation listing has recently been upgraded from amber to red. Detailed demographic work suggests that the decline may have been driven by low annual survival, and that neither increased predation nor interspecific competition is responsible ([Siriwardena 2006](#)). Nest failure rates have fallen during the period of decline. Marsh Tits nest in woods as small as half a hectare ([Hinsley et al. 1995](#)), but there is evidence from CBC that declines are steeper on smaller plots (G.M. Siriwardena, unpubl.). Reductions in the structural and floristic diversity of woodland, resulting partly from increased browsing by deer, are likely to have caused the decline ([Perrins 2003](#), [Fuller et al. 2005](#)). Marsh Tits appear to select breeding territories on the quality of the shrub layer rather than the tree canopy, and may be adversely affected by factors that damage the shrub layer, such as overgrazing and canopy closure ([Hinsley et al. 2007](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)). Following declines elsewhere in western Europe during the 1990s, the European status of this species is no longer considered 'secure' ([BirdLife International 2004](#)).



Population changes

CBC/BBS UK 1966–2007 Marsh Tit




Table of population changes for Marsh Tit

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	89	-68	-77	-56	>50	
	25	1981-2006	107	-38	-51	-16	>25	
	10	1996-2006	164	-16	-28	-2		
	5	2001-2006	154	-16	-27	-4		
CBC/BBS England	39	1967-2006	82	-68	-78	-55	>50	
	25	1981-2006	98	-37	-52	-15	>25	
	10	1996-2006	148	-16	-28	-1		
	5	2001-2006	139	-15	-27	-2		
BBS UK	11	1995-2006	138	-10	-25	10		
	10	1996-2006	141	-12	-27	4		
	5	2001-2006	154	-18	-28	-5		
BBS England	11	1995-2006	124	-13	-30	10		
	10	1996-2006	126	-13	-27	7		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Marsh Tit




BBS UK 1994—2007

Marsh Tit



BBS England 1994—2007

Marsh Tit




Productivity trends

Table of productivity changes for Marsh Tit

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	None				Small sample
Brood size	38	1968-2006	22	None				Small sample
Daily failure rate (eggs)	38	1968-2006	20	Linear decline	0.76% nests/day	0.14% nests/day	-81.6%	Small sample
Daily failure rate (chicks)	38	1968-2006	19	None				Small sample
Laying date	38	1968-2006	14	Linear decline	Apr 28	Apr 19	-9 days	Small sample


Clutch size 1966—2007

Marsh Tit




Brood size 1966—2007

Marsh Tit




Egg stage nest failure rate

Marsh Tit




Chick stage nest failure rate

Marsh Tit



Laying date 1966—2007

Marsh Tit



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

NUTHATCH *Sitta europaea*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK, England: rapid increase

UK population size

144,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Nuthatch abundance has increased rapidly since the mid 1970s. Despite minor setbacks during the 1990s, there is no indication yet of a halt to the upward trend. This increase has been accompanied by a range expansion into northern England ([Gibbons et al. 1993](#)) and has been associated with a large increase in brood size. The reasons for these changes are unknown. A trend towards earlier laying, perhaps as a result of climate change ([Crick et al. 1997](#)), has also been identified. Numbers have risen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007 Nuthatch





Table of population changes for Nuthatch

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	169	177	98	282		
	25	1981-2006	235	92	58	142		
	10	1996-2006	435	40	27	51		
	5	2001-2006	480	25	18	34		
CBC/BBS England	39	1967-2006	145	182	97	310		
	25	1981-2006	200	88	45	143		
	10	1996-2006	364	43	28	56		
	5	2001-2006	400	28	19	35		
BBS UK	11	1995-2006	381	49	36	60		
	10	1996-2006	398	41	29	52		
	5	2001-2006	480	24	16	30		
BBS England	11	1995-2006	316	50	36	67		
	10	1996-2006	330	43	30	58		
	5	2001-2006	400	26	17	35		
BBS Wales	11	1995-2006	65	38	14	67		
	10	1996-2006	67	25	5	49		
	5	2001-2006	79	10	-3	27		


CBC/BBS England 1966—2007

Nuthatch




BBS UK 1994—2007

Nuthatch




BBS England 1994—2007

Nuthatch



BBS Wales 1994—2007

Nuthatch




Productivity trends

Table of productivity changes for Nuthatch

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	26	None				Small sample
Brood size	38	1968-2006	62	Curvilinear	4.05 chicks	4.87 chicks	20.1%	
Daily failure rate (eggs)	38	1968-2006	47	None				
Daily failure rate (chicks)	38	1968-2006	53	None				
Laying date	38	1968-2006	26	Linear decline	May 2	Apr 21	-11 days	Small sample


Clutch size 1966—2007

Nuthatch




Brood size 1966—2007

Nuthatch




Egg stage nest failure rate

Nuthatch



Chick stage nest failure rate


Nuthatch



Insufficient data on CES available for this species

Laying date 1966–2007

Nuthatch



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

TREECREEPER

Certhia familiaris

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK, England: fluctuating, with no long-term trend

UK population size

214,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary



The UK Tree Creeper population peaked in the mid 1970s, but has been roughly stable since about 1980. Intensive study has shown that Tree Creeper numbers and survival rates are reduced by wet winter weather ([Peach et al. 1995b](#)). The influence of cold weather is also evident in the low start to the index, following the severe winter of 1962/63, and the trough around 1980. Census data suggest a minor decline has occurred since the early 1980s, but CES adult captures have increased for much of this period. Productivity, calculated using CES data, shows fluctuations around a long-term shallow increase but a sharp downturn in recent years. There has been a significant fall in nest failure rates at the egg stage (18 days, comprising 14 days incubation and 4 days laying). The trend towards earlier laying can be partly explained by recent climate change ([Crick & Sparks 1999](#)).

Population changes

CBC/BBS UK 1966–2007

Tree Creeper




Table of population changes for Tree Creeper

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	177	14	-15	51		
	25	1981-2006	225	4	-18	27		
	10	1996-2006	361	-1	-14	12		
	5	2001-2006	338	3	-11	19		
CBC/BBS England	39	1967-2006	140	4	-22	44		
	25	1981-2006	176	-6	-24	18		
	10	1996-2006	273	-10	-21	2		
	5	2001-2006	251	-4	-11	6		
CES adults	22	1984-2006	37	60	9	122		
	10	1996-2006	43	8	-13	40		
	5	2001-2006	39	30	7	58		
CES juveniles	22	1984-2006	58	16	-14	65		
	10	1996-2006	67	-10	-23	5		
	5	2001-2006	64	-10	-24	5		
BBS UK	11	1995-2006	305	3	-10	17		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS England	11	1995-2006	226	-8	-19	5		
	10	1996-2006	233	-10	-21	1		
	5	2001-2006	251	-5	-13	2		
BBS Scotland	5	2001-2006	31	38	-10	75		
BBS Wales	11	1995-2006	40	13	-20	45		
	10	1996-2006	41	-1	-26	22		
	5	2001-2006	43	-21	-37	-6		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Treecreeper




CES adult abundance 1983—2007

Treecreeper




CES juvenile abundance 1983—2007

Treecreeper




BBS UK 1994—2007

Treecreeper




BBS England 1994—2007

Treecreeper



BBS Wales 1994—2007

Treecreeper




Productivity trends


Table of productivity changes for Treecreeper

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	None				Small sample
Brood size	38	1968-2006	27	None				Small sample
Daily failure rate (eggs)	38	1968-2006	22	Linear decline	1.91% nests/day	0.62% nests/day	-67.5%	Small sample
Daily failure rate (chicks)	38	1968-2006	23	None				Small sample
Laying date	38	1968-2006	13	Linear decline	May 7	Apr 28	-9 days	Small sample
Juvenile to Adult ratio (CES)	22	1984-2006	66	Smoothed trend	145 Index value	100 Index value	-31%	
Juvenile to Adult ratio (CES)	10	1996-2006	75	Smoothed trend	135 Index value	100 Index value	-26%	
Juvenile to Adult ratio (CES)	5	2001-2006	72	Smoothed trend	146 Index value	100 Index value	-31% >25	


Clutch size 1966—2007




Brood size 1966—2007




Egg stage nest failure rate




Chick stage nest failure rate



Laying date 1966—2007



CES productivity 1983—2007



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

JAY *Garrulus glandarius*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK, England: fluctuating, with no long-term trend

UK population size

160,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

The UK Jay population remained stable in the species' preferred woodland habitat until the late 1980s, after which the population began to decline. This decrease followed an earlier decline on farmland CBC plots ([Gregory & Marchant 1996](#)). Long-term trends are stable overall, and the CBC/BBS index has recorded some increase in the recent ten-year period. No trends are known in breeding performance.



Population changes

CBC/BBS UK 1966–2007

Jay






Table of population changes for Jay

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	284	1	-19	25		
	25	1981-2006	386	-2	-15	13		
	10	1996-2006	704	11	5	18		
	5	2001-2006	756	4	-4	10		
CBC/BBS England	39	1967-2006	252	-6	-26	20		
	25	1981-2006	339	-8	-21	7		
	10	1996-2006	610	5	-3	14		
	5	2001-2006	648	5	-1	13		
BBS UK	11	1995-2006	628	9	1	18		
	10	1996-2006	652	11	4	19		
	5	2001-2006	756	5	-2	11		
BBS England	11	1995-2006	543	1	-7	9		
	10	1996-2006	561	5	-3	11		
	5	2001-2006	648	6	-1	11		
BBS Wales	11	1995-2006	65	24	-4	48		
	10	1996-2006	69	17	-6	36		
	5	2001-2006	79	0	-11	16		


CBC/BBS England 1966—2007




BBS UK 1994—2007



BBS England 1994—2007



BBS Wales 1994—2007




Productivity trends

Table of productivity changes for Jay

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size	38	1968-2006	10	None				Small sample

Insufficient data on clutch size available for this species

Brood size 1966—2007



Insufficient data on egg stage failure available for this species

Insufficient data on nestling failure available for this species

Insufficient data on laying date available for this species

Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

MAGPIE
Pica pica

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK, England: rapid increase

UK population size

650,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary



The remarkable adaptability of Magpies has enabled them to colonise many new urban and suburban localities since the 1960s. Magpies increased steadily until the late 1980s, when abundance stabilised ([Gregory & Marchant 1996](#)). Minor decrease has been recorded in the UK during the last five years. The declines in nest failure rates, during both the egg and the chick stages, have been substantial, perhaps as human persecution of nests has diminished. Larsen traps, introduced to the UK in the late 1980s, are now widely used by gamekeepers as a control measure. Clutch sizes, however, have decreased. A strong trend towards earlier laying has also been identified and may be partly explained by recent climate change ([Crick & Sparks 1999](#)).

Population changes

CBC/BBS UK 1966–2007
Magpie




Table of population changes for Magpie

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	625	101	65	152		
	25	1981-2006	897	27	16	41		
	10	1996-2006	1774	-4	-7	-1		
	5	2001-2006	1854	-6	-9	-4		
CBC/BBS England	39	1967-2006	530	105	61	156		
	25	1981-2006	757	33	18	47		
	10	1996-2006	1476	-3	-6	0		
	5	2001-2006	1527	-5	-7	-2		
BBS UK	11	1995-2006	1648	-2	-5	2		
	10	1996-2006	1698	-4	-7	-1		
	5	2001-2006	1854	-6	-8	-4		
BBS England	11	1995-2006	1371	-2	-6	2		
	10	1996-2006	1406	-3	-7	1		
	5	2001-2006	1527	-4	-7	-2		
BBS Scotland	11	1995-2006	38	11	-15	53		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Wales	11	1995-2006	154	-17	-27	-7		
	10	1996-2006	160	-19	-28	-9		
	5	2001-2006	178	-12	-22	-4		
BBS N.Ireland	11	1995-2006	74	29	-5	56		
	10	1996-2006	80	18	-5	32		
	5	2001-2006	92	-5	-13	5		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Magpie




BBS UK 1994—2007

Magpie




BBS England 1994—2007

Magpie




BBS Scotland 1994—2007

Magpie




BBS Wales 1994—2007

Magpie



BBS N. Ireland 1994—2007

Magpie




Productivity trends

Table of productivity changes for Magpie

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	45	Curvilinear	5.62 eggs	4.56 eggs	-18.8%	
Brood size	38	1968-2006	79	Curvilinear	3.09 chicks	2.84 chicks	-8.1%	
Daily failure rate (eggs)	38	1968-2006	52	Linear decline	2.73% nests/day	0.28% nests/day	-89.7%	
Daily failure rate (chicks)	38	1968-2006	50	Linear decline	1.73% nests/day	0.13% nests/day	-92.5%	
Laying date	38	1968-2006	35	Curvilinear	Apr 21	Mar 21	-31 days	


Clutch size 1966—2007

Magpie




Brood size 1966—2007

Magpie




Egg stage nest failure rate

Magpie




Chick stage nest failure rate

Magpie



Laying date 1966—2007

Magpie



Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

JACKDAW *Corvus monedula*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: moderate increase

UK population size

555,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

Jackdaws have increased in abundance since the 1960s ([Gregory & Marchant 1996](#)), and more recent BBS data suggest that the increase is continuing in all UK countries. As with [Magpie](#), [Rook](#) and [Carrion Crow](#), the increase has been associated with improvements in breeding performance and probably reflects the species' generalist feeding habits, which allow it to exploit diverse and ephemeral food resources. A minor decrease in average brood size has been countered by substantial declines in nest failure rates during the egg and chick stages. Overall, from egg-laying to fledging, the proportion of nests that fail has fallen by about two-thirds. Typically in this species, the younger chicks of a brood perish quickly if food becomes limited. Increases in fledging success are therefore likely to be due to improved parental provisioning success ([Henderson & Hart 1993](#)).



Population changes

CBC/BBS UK 1966–2007

Jackdaw




Table of population changes for Jackdaw

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	489	92	30	206		
	25	1981-2006	728	62	20	124		
	10	1996-2006	1514	25	14	39		
	5	2001-2006	1648	9	4	16		
CBC/BBS England	39	1967-2006	389	79	15	184		
	25	1981-2006	577	56	18	110		
	10	1996-2006	1195	27	19	35		
	5	2001-2006	1299	12	6	17		
BBS UK	11	1995-2006	1427	28	17	41		
	10	1996-2006	1474	24	14	36		
	5	2001-2006	1648	9	3	15		
BBS England	11	1995-2006	1127	32	23	43		
	10	1996-2006	1162	27	19	36		
	5	2001-2006	1299	12	7	18		
BBS Scotland	11	1995-2006	99	8	-9	34		
	10	1996-2006	101	4	-11	29		
	5	2001-2006	107	-2	-16	19		


BBS W Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	155	8	-9	28		
BBS N.Ireland	11	1995-2006	66	58	11	94		
	10	1996-2006	71	50	16	72		
	5	2001-2006	83	8	-2	26		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Jackdaw




BBS UK 1994—2007

Jackdaw




BBS England 1994—2007

Jackdaw




BBS Scotland 1994—2007

Jackdaw




BBS Wales 1994—2007

Jackdaw



BBS N. Ireland 1994—2007

Jackdaw




Productivity trends

Table of productivity changes for Jackdaw

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	43	Linear increase	4.35 eggs	4.57 eggs	4.9%	
Brood size	38	1968-2006	91	Curvilinear	2.69 chicks	2.6 chicks	-3.2%	
Daily failure rate (eggs)	38	1968-2006	54	Linear decline	0.76% nests/day	0.2% nests/day	-73.7%	
Daily failure rate (chicks)	38	1968-2006	52	Linear decline	1.26% nests/day	0.24% nests/day	-81%	
Laying date	38	1968-2006	23	Curvilinear	Apr 23	Apr 18	-5 days	Small sample


Clutch size 1966—2007

Jackdaw




Brood size 1966—2007

Jackdaw




Egg stage nest failure rate

Jackdaw




Chick stage nest failure rate

Jackdaw



Laying date 1966—2007

Jackdaw



Insufficient data for CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

ROOK *Corvus frugilegus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

UK: increase

UK population size

1,120,000–1,430,000 pairs in 1996 (Marchant & Gregory 1999: BiE04); 1,130,000–1,440,000 pairs in 2000 (1996 estimate updated using BBS trend: APEP06)

Status summary

Relatively few rookeries fell within CBC plots, but an index calculated from the available nest counts showed a shallow, long-term increase (Wilson *et al.* 1998). The trend is confirmed by the results of the most recent BTO rookeries survey, which identified a 40% increase in abundance between 1975 and 1996 (Marchant & Gregory 1999). This increase probably reflects the species' considerable adaptability in the face of agricultural change. BBS indices, which are drawn from sightings during transect walks and not from the BBS nest counts, suggest possible decrease in Scotland, Wales and Northern Ireland since around 2000. There have been no clear trends in breeding productivity since the 1960s.



Population changes

BBS UK 1994–2007

Rook






Table of population changes for Rook

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	1141	-6	-15	5		
	10	1996-2006	1169	-7	-15	3		
	5	2001-2006	1241	-9	-16	-2		
BBS England	11	1995-2006	895	2	-9	14		
	10	1996-2006	914	1	-8	11		
	5	2001-2006	968	2	-6	11		
BBS Scotland	11	1995-2006	101	-20	-39	14		
	10	1996-2006	102	-18	-38	13		
	5	2001-2006	100	-21	-35	-4		
BBS Wales	11	1995-2006	75	-22	-44	7		
	10	1996-2006	78	-23	-45	-1		
	5	2001-2006	85	-14	-31	3		
BBS N.Ireland	11	1995-2006	67	14	-15	51		
	10	1996-2006	72	3	-23	22		
	5	2001-2006	85	-27	-39	-13	>25	


BES England 1994—2007




BES Scotland 1994—2007



BES Wales 1994—2007



BES N. Ireland 1994—2007




Productivity trends


Table of productivity changes for Rook

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	Curvilinear	4.15 eggs	4.12 eggs	-0.7%	Small sample
Brood size	38	1968-2006	85	Curvilinear	2.22 chicks	2.34 chicks	5.2%	
Daily failure rate (eggs)	38	1968-2006	32	None				
Daily failure rate (chicks)	38	1968-2006	51	None				
Laying date	38	1968-2006	12	None				Small sample


Clutch size 1966—2007




Brood size 1966—2007




Egg stage nest failure rate



Chick stage nest failure rate



Laying date 1966—2007



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

CARRION CROW *Corvus corone*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe (*C. corone/cornix*): no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK (*C. corone/cornix*): green

Long-term trend

England: rapid increase

UK population size

790,000 territories in 1990 (1988–91 Atlas: [APEP06](#)); 987,500 pairs in 2000 (updated using CBC/BBS trend)

Status summary

Carrion Crows have increased steadily since the 1960s ([Gregory & Marchant 1996](#)) and only now are there any signs of the population size stabilising. This trend has been associated with increases in nesting success and with earlier laying (perhaps an effect of climate change: [Crick et al. 1997](#)) and probably reflects the species' adaptability to changing habitats and the exploitation of ephemeral food resources in intensive agriculture. Reduced control activities by gamekeepers may also have contributed ([Marchant et al. 1990](#)), as may an increase in roadside carrion.



Population changes

CBC/BBS England 1966–2007 Carrion Crow








Table of population changes for Carrion Crow






Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	614	119	77	179		Includes Hooded Crow
	25	1981-2006	884	49	27	77		Includes Hooded Crow
	10	1996-2006	1764	14	7	21		Includes Hooded Crow
	5	2001-2006	1858	4	-1	9		Includes Hooded Crow
BBS UK	11	1995-2006	2013	14	7	21		
	10	1996-2006	2069	9	4	15		
	5	2001-2006	2272	-1	-5	5		
BBS England	11	1995-2006	1647	19	11	25		
	10	1996-2006	1693	15	8	21		
	5	2001-2006	1858	4	0	8		
BBS Scotland	11	1995-2006	164	1	-17	21		
	10	1996-2006	167	-3	-18	15		
	5	2001-2006	173	-11	-26	5		
BBS Wales	11	1995-2006	189	13	-3	33		
	10	1996-2006	197	7	-8	26		
	5	2001-2006	227	-2	-14	14		



Productivity trends

Table of productivity changes for Carrion Crow

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	33	Curvilinear	4.08 eggs	4.09 eggs	0.1%	
Brood size	38	1968-2006	79	Curvilinear	2.87 chicks	2.48 chicks	-13.9%	
Daily failure rate (eggs)	38	1968-2006	50	Linear decline	1.59% nests/day	0.22% nests/day	-86.2%	
Daily failure rate (chicks)	38	1968-2006	42	Linear decline	0.74% nests/day	0.14% nests/day	-81.1%	
Laying date	38	1968-2006	32	Curvilinear	Apr 16	Apr 5	-11 days	



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

HOODED CROW

Corvus cornix

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe (*C. corone/cornix*): no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK (*C. corone/cornix*): green

Long-term trend

UK: uncertain

UK population size

213,900 territories in 1990 (1988–91 Atlas: [APEP06](#))

Status summary

The BOU Records Committee took the decision in 2002 to treat Hooded Crow and [Carrion Crow](#) as separate species ([Parkin et al. 2003](#)). This split is not yet recognised in conservation listings. In the UK, Hooded Crows occur in Northern Ireland, the Isle of Man, and in Scotland, mainly west and north of the Great Glen. Retrospective analysis of BBS trends is simple because observers record Hooded Crows (coded HC) separately from Carrion Crows and from intermediates (coded HB). Intermediate forms between Carrion and Hooded, which predominate in a band across western Scotland and occur less frequently elsewhere in the UK, are not included in either BBS index. BBS data suggest that some decrease in Hooded Crows may have occurred in Scotland, but that this has been countered by increase in Northern Ireland. Hooded Crows have increased markedly in Ireland since 1924 ([Hutchinson 1989](#)).



Population changes

BBS UK 1994–2007

Hooded Crow





Table of population changes for Hooded Crow

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	126	-1	-20	25		
	10	1996-2006	131	3	-14	24		
	5	2001-2006	142	1	-14	17		
BBS Scotland	11	1995-2006	50	-27	-45	0		
	10	1996-2006	49	-19	-39	8		
	5	2001-2006	46	9	-14	34		
BBS N.Ireland	11	1995-2006	71	111	59	160		
	10	1996-2006	77	76	37	100		
	5	2001-2006	92	-4	-15	11		


BBS Scotland 1994—2007

Hooded Crow



BBS N. Ireland 1994—2007

Hooded Crow



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

RAVEN

Corvus corax

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: green

Long-term trend

UK: probable increase

UK population size

12,900 pairs in 2000 (1988–91 Atlas estimate updated using BBS trend: [BiE04](#), [APEP06](#))



Status summary

Between the two atlas periods, the Raven's range contracted from some areas of Scotland and northern England. Declines in southern Scotland and northern England were associated with large-scale afforestation ([Marquiss et al. 1978](#)), while closer sheep husbandry and conversion of pasture to arable were also implicated ([Mearns 1983](#)). A thorough survey of northwest Wales during 1998 to 2005 found at least 69% more nesting pairs than a previous survey of the same area during 1978–85 and evidence of an increase of 173% since around 1950, at a rate that accelerated after 1990 ([Driver 2006](#)). Ravens have also increased along the English–Welsh border and in parts of lowland England, helping to balance the local declines in northern Britain ([Cross 2002](#)). BBS indicates steep increase in England, Scotland and Wales since 1994. Nesting success appears to have improved, but brood size has fallen. Ravens are estimated to have increased by 118% across Europe during 1980–2005 ([PECBMS 2007](#)).

Population changes

BBS UK 1994–2007

Raven





Table of population changes for Raven

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	214	80	40	133		
	10	1996-2006	224	76	36	124		
	5	2001-2006	266	34	8	70		
BBS England	11	1995-2006	71	172	68	303		
	10	1996-2006	75	152	63	270		
	5	2001-2006	100	61	28	99		
BBS Scotland	11	1995-2006	39	63	3	138		
	10	1996-2006	39	67	2	160		
	5	2001-2006	38	51	-11	143		
BBS Wales	11	1995-2006	82	46	-5	138		
	10	1996-2006	86	39	-6	109		
	5	2001-2006	101	4	-14	25		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS England 1994—2007



BBS Scotland 1994—2007



BBS Wales 1994—2007




Productivity trends


Table of productivity changes for Raven

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	13	None				Small sample
Brood size	38	1968-2006	67	Linear decline	3.23 chicks	2.83 chicks	-12.4%	
Daily failure rate (eggs)	38	1968-2006	22	Curvilinear	0.21% nests/day	0.09% nests/day	-57.1%	Small sample
Daily failure rate (chicks)	38	1968-2006	29	Curvilinear	0.03% nests/day	0.02% nests/day	-33.3%	Small sample
Laying date	38	1968-2006	11	None				Small sample


Clutch size 1966—2007




Brood size 1966—2007




Egg stage nest failure rate



Chick stage nest failure rate



Laying date 1966—2007



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

STARLING *Sturnus vulgaris*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: SPEC category 3 (declining)
 UK: red (>50% population decline)
 UK Biodiversity Action Plan: priority species

Long-term trend

England: rapid decline

UK population size

804,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#));
 8,500,000 birds in Britain in 1994–2000 ([Robinson et al. 2005a](#))



Status summary

The abundance of breeding Starlings in the UK has fallen rapidly, particularly since the early 1980s, and especially in woodland ([Robinson et al. 2002, 2005a](#)) and continues to be strongly downward. The declines have been greatest in the south and west of Britain; recent BBS data suggest that populations are also decreasing in Scotland and Northern Ireland, where the trends were initially upward. The species' UK conservation listing has been upgraded from amber to red as the decline has become more severe. Strong improvements have occurred in breeding performance, suggesting that decreasing survival rates, particularly of young birds, may be responsible for the observed decline ([Freeman et al. 2002, 2007b](#)). Loss of permanent pasture, which is the species' preferred feeding habitat, and general intensification of livestock rearing are likely to be having adverse effects on rural populations, but other causes should be sought in urban areas ([Robinson et al. 2002, 2005a](#)). Widespread declines in northern Europe during the 1990s outweighed increases in the south, and the European status of this species is no longer considered 'secure' ([BirdLife International 2004](#)).

Population changes

CBC/BBS England 1966–2007 Starling




Table of population changes for Starling

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	492	-83	-88	-77	>50	
	25	1981-2006	702	-78	-83	-73	>50	
	10	1996-2006	1394	-36	-39	-32	>25	
	5	2001-2006	1406	-20	-25	-16		
BBS UK	11	1995-2006	1631	-29	-34	-24	>25	
	10	1996-2006	1665	-28	-32	-23	>25	
	5	2001-2006	1739	-17	-22	-14		
BBS England	11	1995-2006	1335	-38	-41	-34	>25	
	10	1996-2006	1359	-36	-39	-32	>25	
	5	2001-2006	1406	-20	-24	-16		
BBS Scotland	11	1995-2006	134	-9	-26	6		
	10	1996-2006	136	-12	-27	2		
	5	2001-2006	145	-16	-27	-4		
BBS Wales	11	1995-2006	83	-46	-64	-23	>25	


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS N.Ireland	11	1995-2006	69	86	35	169		
	10	1996-2006	75	64	23	108		
	5	2001-2006	89	-8	-21	7		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS UK 1994—2007

Starling




BBS England 1994—2007

Starling




BBS Scotland 1994—2007

Starling




BBS Wales 1994—2007

Starling



BBS N. Ireland 1994—2007

Starling




Productivity trends

Table of productivity changes for Starling

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	76	Linear increase	4.42 eggs	4.97 eggs	12.6%	
Brood size	38	1968-2006	205	None				
Daily failure rate (eggs)	38	1968-2006	118	Linear decline	1.12% nests/day	0.3% nests/day	-73.2%	
Daily failure rate (chicks)	38	1968-2006	136	Linear decline	0.63% nests/day	0.19% nests/day	-69.8%	
Laying date	38	1968-2006	82	None				


Clutch size 1966—2007




Starling



Brood size 1966—2007

Starling





Insufficient data on CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

HOUSE SPARROW

Passer domesticus

• Population changes

• Productivity trends

• Additional information



Conservation listings

Europe: SPEC category 3, declining
 UK: red (>50% population decline)
 UK Biodiversity Action Plan: priority species

Long-term trend

England: rapid decline

UK population size

2,100,000–3,675,000 pairs in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#)); about 6 million pairs in Britain ([Robinson et al. 2005b](#))

Status summary

CBC sample sizes did not allow monitoring of House Sparrows until 1976; previously, there had been many farmland plots with high populations that could not be properly quantified without better access to farm buildings and housing. CBC/BBS data indicate a rapid decline in abundance over the last 25 years, as does the BTO's Garden Bird Feeding Survey ([Siriwardena et al. 2002](#), [Robinson et al. 2005b](#)). These results are supported by many other studies and anecdotal reports, and have generated great conservation concern (see [Summers-Smith 2003](#)). A change in the listing criteria resulted in the admission of the species, green-listed until 2002, to the red list. A temporary drop in first-year survival coincided with the steepest decline, but changes in breeding performance, especially nest failure rates at the chick stage, have also helped drive population change ([Freeman & Crick 2002](#)). Possible explanations include a general reduction in food supply, less grain being spilt during agricultural operations, tighter hygiene regulations, increases in predation, and toxic additives to unleaded petrol ([Siriwardena et al. 2002](#), [Robinson et al. 2005b](#), [Vincent 2005](#), [Summers-Smith 2007](#), [Peach et al. 2008](#)).

The overall national decline since the 1970s masks much heterogeneity by region and habitat, and population processes may be relatively fine-grained: overall, populations in rural areas had declined by 47% by 2000, and those in urban and suburban areas by about 60% (CBC data: [Robinson et al. 2005b](#)). Within urban areas, House Sparrows may have disappeared predominantly from more affluent areas, where changes are more likely to have occurred to habitat structure ([Shaw et al. 2008](#)). The continued availability of allotments, gardens and other green spaces in urban areas is crucial to preventing further decline ([Chamberlain et al. 2007](#)). BBS suggests increases recently in Scotland and Wales. Overall, brood size has decreased, raising NRS concern ([Leech & Barimore 2008](#)), but nest success has improved markedly. Following widespread declines across Europe since the 1980s, the European status of this species is no longer considered 'secure' ([BirdLife International 2004](#)).

Population changes

CBC/BBS England 1976–2007 House Sparrow




Table of population changes for House Sparrow

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	29	1977-2006	511	-70	-79	-60	>50	
	25	1981-2006	587	-59	-71	-49	>50	


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	1416	-6	-12	1		
	10	1996-2006	1453	-4	-10	2		
	5	2001-2006	1567	1	-2	5		
BBS England	11	1995-2006	1169	-15	-20	-10		
	10	1996-2006	1195	-13	-17	-9		
	5	2001-2006	1272	-3	-6	1		
BBS Scotland	11	1995-2006	80	37	2	75		
	10	1996-2006	81	39	7	73		
	5	2001-2006	85	13	-5	26		
BBS Wales	11	1995-2006	113	68	45	100		
	10	1996-2006	117	53	33	76		
	5	2001-2006	138	14	5	26		
BBS N.Ireland	11	1995-2006	45	43	-18	91		
	10	1996-2006	49	47	4	65		
	5	2001-2006	58	18	-4	36		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


BBS UK 1994—2007

House Sparrow




BBS England 1994—2007

House Sparrow




BBS Scotland 1994—2007

House Sparrow




BBS Wales 1994—2007

House Sparrow



BBS N. Ireland 1994—2007

House Sparrow




Productivity trends

Table of productivity changes for House Sparrow

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	66	None				
Brood size	38	1968-2006	111	Curvilinear	3.39 chicks	2.92 chicks	-13.9%	
Daily failure rate (eggs)	38	1968-2006	92	Linear decline	1.15% nests/day	0.41% nests/day	-64.3%	
Daily failure rate (chicks)	38	1968-2006	88	Linear decline	1.3% nests/day	0.36% nests/day	-72.3%	
Laying date	38	1968-2006	51	Linear decline	May 25	May 18	-7 days	


Clutch size 1966—2007

House Sparrow




Brood size 1966—2007

House Sparrow




Egg stage nest failure rate

House Sparrow




Chick stage nest failure rate

House Sparrow



Laying date 1966—2007

House Sparrow



Insufficient data for CES
available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

TREE SPARROW

Passer montanus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 3 (declining)

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

England: rapid decline

UK population size

68,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Tree Sparrow abundance crashed spectacularly in the UK between the late 1970s and the early 1990s. BBS data indicate significant increase since 1994, but it should be remembered that, for every Tree Sparrow today there were around 30 in the 1970s, and any recovery therefore has a very long way to go. Clear range contractions occurred between the two breeding atlas periods ([Gibbons *et al.* 1993](#)), and have continued subsequently, with many local extinctions occurring during the 1990s. Components of agricultural intensification, such as reductions in winter stubble, are likely to be implicated in the decline. Breeding performance has improved substantially as population sizes have decreased, suggesting that decreases in productivity were not responsible for the decline. It is more likely that survival was the critical demographic measure, although ring-recovery analyses have produced equivocal results because of small sample sizes ([Siriwardena *et al.* 1998b, 2000b](#)). Following declines across western and northwestern Europe during the 1990s, the European status of this species is no longer considered 'secure' ([BirdLife International 2004](#)).



Population changes

CBC/BBS England 1966–2007

Tree Sparrow






Table of population changes for Tree Sparrow

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	83	-97	-99	-94	>50	
	25	1981-2006	80	-93	-97	-88	>50	
	10	1996-2006	126	24	7	54		
	5	2001-2006	125	6	-8	24		
BBS UK	11	1995-2006	147	46	12	100		
	10	1996-2006	149	48	13	99		
	5	2001-2006	156	20	-3	46		
BBS England	11	1995-2006	121	20	-3	44		
	10	1996-2006	121	20	-1	41		
	5	2001-2006	125	6	-8	24		








The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB



Productivity trends

Table of productivity changes for Tree Sparrow

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	155	Curvilinear	4.71 eggs	5.11 eggs	8.5%	
Brood size	38	1968-2006	197	Curvilinear	3.78 chicks	4.21 chicks	11.2%	
Daily failure rate (eggs)	38	1968-2006	207	Curvilinear	0.74% nests/day	0.32% nests/day	-56.8%	
Daily failure rate (chicks)	38	1968-2006	151	Curvilinear	1.25% nests/day	0.64% nests/day	-48.8%	
Laying date	38	1968-2006	167	Linear decline	May 29	May 24	-5 days	



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

CHAFFINCH *Fringilla coelebs*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: shallow increase

UK population size

5,974,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))



Status summary

Chaffinch abundance has increased rapidly since the early 1970s, according to CBC/BBS and CES, but numbers seemed to stabilise for a period during the 1990s. This relative stability was associated with a reduction in annual survival, which could be density-dependent ([Siriwardena et al. 1999](#)). There was also some evidence of improved breeding performance during the early years of population increase, with larger brood sizes and fewer egg-stage nest failures, but these trends are now reversed. The trend towards earlier laying may be partly explained by recent climate change ([Crick & Sparks 1999](#)). Chaffinches are well adapted to suburban and garden habitats, as well as to highly fragmented woodland and hedgerows, occurring less in the open-field, arable habitats that have been affected most by agricultural intensification, so it is possible that they have benefited by environmental changes from which other seed-eating passerines have suffered.

Population changes

CBC/BBS UK 1966–2007 Chaffinch




Table of population changes for Chaffinch


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	821	30	14	45		
	25	1981-2006	1164	19	11	30		
	10	1996-2006	2291	12	7	16		
	5	2001-2006	2406	7	4	10		
CBC/BBS England	39	1967-2006	651	38	22	56		
	25	1981-2006	915	27	18	39		
	10	1996-2006	1781	17	13	21		
	5	2001-2006	1860	8	5	11		
CES adults	22	1984-2006	78	27	-27	106		
	10	1996-2006	88	-9	-28	13		
	5	2001-2006	84	-1	-11	11		
CES juveniles	22	1984-2006	59	75	-21	251		
	10	1996-2006	69	21	-31	86		
	5	2001-2006	69	43	10	70		
BBS UK	11	1995-2006	2129	13	9	18		
	10	1996-2006	2190	11	7	16		
	5	2001-2006	2406	6	4	9		

BBS E ¹ Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	1860	8	5	10		
BBS Scotland	11	1995-2006	205	6	-3	20		
	10	1996-2006	207	5	-4	17		
	5	2001-2006	211	7	-1	15		
BBS Wales	11	1995-2006	186	-3	-17	11		
	10	1996-2006	193	0	-11	11		
	5	2001-2006	223	6	-1	14		
BBS N.Ireland	11	1995-2006	79	37	4	57		
	10	1996-2006	86	20	-3	33		
	5	2001-2006	98	-5	-14	6		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007 Chaffinch




CES adult abundance 1983–2007 Chaffinch




CES juvenile abundance 1983–2007 Chaffinch




BBS UK 1994–2007 Chaffinch




BBS England 1994–2007 Chaffinch




BBS Scotland 1994–2007 Chaffinch



BBS Wales 1994–2007 Chaffinch



BBS N. Ireland 1994–2007 Chaffinch




Productivity trends


Table of productivity changes for Chaffinch

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	86	Curvilinear	4.22 eggs	4.03 eggs	-4.4%	
Brood size	38	1968-2006	136	Curvilinear	3.57 chicks	3.49 chicks	-2.5%	
Daily failure rate (eggs)	38	1968-2006	165	Curvilinear	2.97% nests/day	3.68% nests/day	23.9%	
Daily failure rate (chicks)	38	1968-2006	116	None				
Laying date	38	1968-2006	107	Linear decline	May 11	May 3	-8 days	
Juvenile to Adult ratio (CES)	22	1984-2006	84	Smoothed trend	62 Index value	100 Index value	61%	
Juvenile to Adult ratio (CES)	10	1996-2006	95	Smoothed trend	103 Index value	100 Index value	-3%	
Juvenile to Adult ratio (CES)	5	2001-2006	92	Smoothed trend	86 Index value	100 Index value	16%	


Clutch size 1966—2007
Chaffinch




Brood size 1966—2007
Chaffinch




Egg stage nest failure rate
Chaffinch




Chick stage nest failure rate
Chaffinch



Laying date 1966—2007
Chaffinch



CES productivity 1983—2007
Chaffinch



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

GREENFINCH *Carduelis chloris*

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK, England: shallow increase

UK population size

734,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary



Greenfinch abundance varied little up to the mid 1990s, and there was little change in either survival or breeding performance during this period ([Siriwardena et al. 1998b, 2000b](#)). More recent CBC/BBS data indicate population increases widely across the UK. Productivity data have become more complex, with a substantial reduction in brood size and increased nest survival at the egg stage. Possibly these recent changes are linked to the species' regular year-round use of gardens for feeding. The trend towards earlier laying may be explained by recent climate change ([Crick & Sparks 1999](#)).

Population changes

CBC/BBS UK 1966–2007 Greenfinch




Table of population changes for Greenfinch

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	596	26	-1	59		
	25	1981-2006	846	46	20	68		
	10	1996-2006	1709	26	21	33		
	5	2001-2006	1871	6	3	10		
CBC/BBS England	39	1967-2006	505	38	6	67		
	25	1981-2006	714	50	22	77		
	10	1996-2006	1435	30	25	36		
	5	2001-2006	1559	7	4	11		
CES adults	22	1984-2006	43	59	-21	278		
	10	1996-2006	48	-15	-35	9		
	5	2001-2006	48	-13	-30	10		
CES juveniles	22	1984-2006	27	-35	-67	53		
	10	1996-2006	35	13	-30	63		
	5	2001-2006	38	43	-12	99		
BBS UK	11	1995-2006	1595	31	25	39		
	10	1996-2006	1648	26	21	32		
	5	2001-2006	1871	6	3	9		


BBS Ex Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	1559	7	5	11		
BBS Scotland	11	1995-2006	93	8	-12	32		
	10	1996-2006	96	4	-14	24		
	5	2001-2006	103	-1	-17	13		
BBS Wales	11	1995-2006	106	28	5	57		
	10	1996-2006	111	20	-2	45		
	5	2001-2006	132	0	-8	13		
BBS N.Ireland	11	1995-2006	46	115	34	204		
	10	1996-2006	50	82	25	122		
	5	2001-2006	63	18	-3	37		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Greenfinch




CES adult abundance 1983—2007

Greenfinch




CES juvenile abundance 1983—2007

Greenfinch




BBS UK 1994—2007

Greenfinch




BBS England 1994—2007

Greenfinch




BBS Scotland 1994—2007

Greenfinch



BBS Wales 1994—2007

Greenfinch




Productivity trends

Table of productivity changes for Greenfinch


Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	93	Curvilinear	4.73 eggs	4.54 eggs	-4%	
Brood size	38	1968-2006	115	Curvilinear	4.03 chicks	3.61 chicks	-10.5%	
Daily failure rate (eggs)	38	1968-2006	132	Linear decline	2.48% nests/day	1.8% nests/day	-27.4%	
Daily failure rate (chicks)	38	1968-2006	98	None				
Laying date	38	1968-2006	95	Linear decline	May 25	May 10	-15 days	
Juvenile to Adult ratio (CES)	22	1984-2006	47	Smoothed trend	148 Index value	100 Index value	-32%	

Juvenile to Adult ratio (CES)	Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment


Clutch size 1966—2007
Greenfinch




Brood size 1966—2007
Greenfinch




Egg stage nest failure rate
Greenfinch




Chick stage nest failure rate
Greenfinch



Laying date 1966—2007
Greenfinch



CES productivity 1983—2007
Greenfinch



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

GOLDFINCH *Carduelis carduelis*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)
UK: green

Long-term trend

England: fluctuating, with no long-term trend

UK population size

313,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Goldfinch abundance fell sharply from the mid 1970s until the mid 1980s, but the decline was both preceded and followed by significant population increases. The recent upturn has lifted the species from the amber list of conservation concern into the green category, and has been accompanied by an increase in its use of gardens for winter feeding. These population changes can be explained almost entirely by changes in annual survival rates, which may have resulted from a reduction in the availability of weed seeds, due to agricultural intensification, and subsequent increased use of other food sources such as garden bird tables. Alternatively, the effects of environmental change or increased hunting pressure in France and Iberia, where the migrant majority of the population wintered, may have temporarily reduced survival rates ([Siriwardena et al. 1999](#)). There has been some long-term reduction in productivity as measured by CES.



Population changes

CBC/BBS England 1966–2007 Goldfinch




Table of population changes for Goldfinch


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	382	18	-11	53		
	25	1981-2006	543	28	8	48		
	10	1996-2006	1103	24	17	30		
	5	2001-2006	1213	13	7	18		
CES adults	22	1984-2006	33	14	-44	77		
	10	1996-2006	40	-19	-40	10		
	5	2001-2006	38	-7	-34	36		
CES juveniles	22	1984-2006	19	-45	-74	16		Small sample
	10	1996-2006	23	-11	-46	24		
	5	2001-2006	22	12	-34	75		
BBS UK	11	1995-2006	1266	35	25	47		
	10	1996-2006	1309	34	25	45		
	5	2001-2006	1501	16	10	22		
BBS England	11	1995-2006	1036	24	17	32		
	10	1996-2006	1067	25	18	32		
	5	2001-2006	1213	14	10	19		

BBS Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	84	18	-7	50		
BBS Wales	11	1995-2006	115	56	21	113		
	10	1996-2006	120	45	14	93		
	5	2001-2006	142	-1	-9	11		
BBS N.Ireland	11	1995-2006	34	543	.	.		
	10	1996-2006	37	442	.	.		
	5	2001-2006	52	101	.	.		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CES adult abundance 1983—2007
Goldfinch




CES juvenile abundance 1983—2007
Goldfinch




BBS UK 1994—2007
Goldfinch




BBS England 1994—2007
Goldfinch



BBS Scotland 1994—2007
Goldfinch



BBS Wales 1994—2007
Goldfinch




Productivity trends

Table of productivity changes for Goldfinch

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	19	None				Small sample
Brood size	38	1968-2006	32	None				
Daily failure rate (eggs)	38	1968-2006	35	None				
Daily failure rate (chicks)	38	1968-2006	29	None				Small sample
Laying date	38	1968-2006	22	Linear decline	Jun 6	May 30	-7 days	Small sample
Juvenile to Adult ratio (CES)	22	1984-2006	38	Smoothed trend	267 Index value	100 Index value	-63% >50	
Juvenile to Adult ratio (CES)	10	1996-2006	46	Smoothed trend	148 Index value	100 Index value	-32%	
Juvenile to Adult ratio (CES)	5	2001-2006	45	Smoothed trend	164 Index value	100 Index value	-39%	


Clutch size 1966—2007

Goldfinch




Brood size 1966—2007

Goldfinch




Egg stage nest failure rate

Goldfinch




Chick stage nest failure rate

Goldfinch




Laying date 1966—2007

Goldfinch



CES productivity 1983—2007

Goldfinch



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

SISKIN

Carduelis spinus

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: green

Long-term trend

UK: increase

UK population size

369,000 pairs in 2000 (1988–91 Atlas estimate updated using BBS trend: [BiE04](#), [APEP06](#))

Status summary

The maturing of new conifer plantations has aided the spread of breeding Siskins throughout the UK, from their previous stronghold in the Scottish Highlands, since about 1950. Its habit of using garden feeders, especially in late winter, has developed since the 1960s and, despite many of the birds involved migrating to the Baltic region to breed, may also have helped to boost the UK breeding population. The 1988–91 Breeding Atlas identified a considerable expansion of the breeding range into southern Britain ([Gibbons *et al.* 1993](#)). More CBC plots became occupied during the 1970s and 1980s, but samples were still insufficient for annual monitoring until BBS began in 1994. Results since then show extraordinary fluctuations, in both England and Scotland, which have been largely in parallel. To some extent, this may reflect the occasional large continental influxes affecting numbers on a broad UK scale.



Population changes

BBS UK 1994–2007

Siskin





Table of population changes for Siskin

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	11	1995-2006	120	-15	-32	4		
	10	1996-2006	123	-21	-37	-3		
	5	2001-2006	123	4	-15	24		
BBS England	11	1995-2006	39	9	-39	129		
	10	1996-2006	40	-4	-47	92		
	5	2001-2006	43	38	-10	130		
BBS Scotland	11	1995-2006	54	-25	-43	-9	>25	
	10	1996-2006	55	-30	-48	-15	>25	
	5	2001-2006	50	-9	-27	7		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB

BBS England 1994—2007



BBS Scotland 1994—2007



Productivity trends

Productivity information is not currently available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

LINNET

Carduelis cannabina

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 2, declining

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

England: rapid decline

UK population size

556,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BiE04](#), [APEP06](#))

Status summary

Linnet abundance fell rapidly in the UK between the mid 1970s and mid 1980s. Numbers have subsequently changed little overall, although with further decrease in England and Wales and possibly some increase in Northern Ireland. CES has shown declines continuing strongly in recent years. Nest failure rates rose during the principal period of population decline, and this represents the most likely demographic mechanism driving the observed decreases in abundance ([Siriwardena et al. 1999, 2000b](#)). CES and nest record results suggest that low productivity is still a problem for the species, possibly due to reductions in hedgerow quality leaving nests more exposed and therefore at greater risk of predation. Recent decreases in clutch and brood sizes, and in nest survival at the chick stage, raise NRS concern ([Leech & Barimore 2008](#)). Nestling diet incorporates a high proportion of oilseed rape seeds, suggesting that the inclusion of this crop in arable rotations may be important in maintaining Linnet populations ([Moorcroft et al. 2006](#)). Linnets are estimated to have declined by 54% across Europe during 1980–2005 ([PECBMS 2007](#)). Following widespread declines across Europe during the 1990s, the European status of this species is no longer considered 'secure' ([BirdLife International 2004](#)).



Population changes

CBC/BBS England 1966–2007

Linnet




Table of population changes for Linnet


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS England	39	1967-2006	367	-75	-81	-66	>50	
	25	1981-2006	501	-47	-57	-32	>25	
	10	1996-2006	967	-27	-32	-21	>25	
	5	2001-2006	953	-22	-25	-16		
CES adults	22	1984-2006	20	-92	-97	-84	>50	
	10	1996-2006	18	-62	-84	-35	>50	Small sample
	5	2001-2006	14	-14	-54	40		Small sample
CES juveniles	22	1984-2006	14	-96	-99	-87	>50	Small sample
	10	1996-2006	14	-67	-91	-38	>50	Small sample
	5	2001-2006	12	-6	-64	71		Small sample
BBS UK	11	1995-2006	1122	-22	-29	-16		
	10	1996-2006	1145	-20	-26	-14		
	5	2001-2006	1193	-16	-22	-10		
BBS England	11	1995-2006	909	-30	-35	-23	>25	
	10	1996-2006	925	-26	-31	-19	>25	

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS Scotland	10	1996-2006	85	1	-24	21		
	5	2001-2006	88	-3	-20	18		
BBS Wales	11	1995-2006	91	-14	-34	14		
	10	1996-2006	94	-17	-37	7		
	5	2001-2006	106	-26	-40	-8	>25	
BBS N.Ireland	11	1995-2006	31	48	-7	109		
	10	1996-2006	33	37	-10	93		
	5	2001-2006	37	4	-19	36		




The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CES adult abundance 1983—2007




CES juvenile abundance 1983—2007




BBS UK 1994—2007




BBS England 1994—2007



BBS Scotland 1994—2007



BBS Wales 1994—2007




Productivity trends


Table of productivity changes for Linnet

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	108	Curvilinear	4.7 eggs	4.57 eggs	-2.7%	
Brood size	38	1968-2006	122	Curvilinear	4.07 chicks	4.08 chicks	0.3%	
Daily failure rate (eggs)	38	1968-2006	154	Curvilinear	1.65% nests/day	2.29% nests/day	38.8%	
Daily failure rate (chicks)	38	1968-2006	110	Linear increase	1.52% nests/day	2.24% nests/day	47.4%	
Laying date	38	1968-2006	109	None				
Juvenile to Adult ratio (CES)	22	1984-2006	22	Smoothed trend	474 Index value	100 Index value	-79% >50	
Juvenile to Adult ratio (CES)	10	1996-2006	21	Smoothed trend	177 Index value	100 Index value	-44%	
Juvenile to Adult ratio (CES)	5	2001-2006	18	Smoothed trend	186 Index value	100 Index value	-46%	Small sample


Clutch size 1966—2007




Brood size 1966—2007




Egg stage nest failure rate




Chick stage nest failure rate



Laying date 1966—2007



CES productivity 1983—2007



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

BULLFINCH *Pyrrhula pyrrhula*

• Population changes

• Productivity trends

• Additional information



Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK: moderate decline

England: rapid decline

UK population size

166,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

The UK Bullfinch population entered a long period of decline in the mid 1970s, following a period of relative stability. The decline was initially very steep, and more so in farmland than in wooded habitats, but has been shallower since the early 1980s. CES and CBC/BBS both suggest there are large fluctuations around the overall downward trend. The demographic mechanism of decline remains unclear ([Siriwardena et al. 1999, 2000b, 2001](#)), although agricultural intensification and a reduction in the structural and floristic diversity of woodland are suspected to have played a part through losses of food resources and nesting cover ([Fuller et al. 2005](#)). Alongside these factors, [Proffitt et al. \(2004\)](#) and [Marquiss \(2007\)](#) mention the constraints on survival outside the breeding season and the possible role of increasing Sparrowhawk populations on the ability of Bullfinches to exploit resources in some habitats. Recent decreases in brood size and in nest survival have raised NRS concern ([Leech & Barimore 2008](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).

Population changes

CBC/BBS UK 1966–2007 Bullfinch




Table of population changes for Bullfinch

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	271	-50	-61	-37	>25	
	25	1981-2006	340	-28	-39	-14	>25	
	10	1996-2006	585	-11	-17	-3		
	5	2001-2006	584	7	1	17		
CBC/BBS England	39	1967-2006	221	-52	-62	-39	>50	
	25	1981-2006	271	-31	-42	-19	>25	
	10	1996-2006	452	-9	-17	-2		
	5	2001-2006	446	1	-7	8		
CES adults	22	1984-2006	80	-19	-36	2		
	10	1996-2006	85	-20	-33	-7		
	5	2001-2006	77	8	-7	25		
CES juveniles	22	1984-2006	62	-21	-43	21		
	10	1996-2006	67	-14	-29	3		


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
BBS UK	10	1996-2006	530	-9	-18	-2		
	5	2001-2006	584	5	-3	15		
BBS England	11	1995-2006	396	-12	-19	-3		
	10	1996-2006	406	-7	-15	0		
BBS Scotland	5	2001-2006	446	1	-6	8		
	11	1995-2006	32	-12	-39	14		
BBS Wales	10	1996-2006	34	-23	-45	-5		
	5	2001-2006	32	-5	-25	19		
BBS UK	11	1995-2006	60	-3	-21	24		
	10	1996-2006	62	7	-11	35		
	5	2001-2006	75	15	0	35		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007

Bullfinch




CES adult abundance 1983–2007

Bullfinch




CES juvenile abundance 1983–2007

Bullfinch




BBS UK 1994–2007

Bullfinch




BBS England 1994–2007

Bullfinch



BBS Wales 1994–2007

Bullfinch



Productivity trends


Table of productivity changes for Bullfinch

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	35	None				
Brood size	38	1968-2006	36	Curvilinear	4.11 chicks	3.79 chicks	-8%	
Daily failure rate (eggs)	38	1968-2006	50	Curvilinear	3.33% nests/day	4.13% nests/day	24%	
Daily failure rate (chicks)	38	1968-2006	34	Curvilinear	3.36% nests/day	4.05% nests/day	20.5%	
Laying date	38	1968-2006	32	None				
Juvenile to Adult ratio (CES)	22	1984-2006	84	Smoothed trend	97 Index value	100 Index value	3%	

Juvenile CES	Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Juvenile to Adult ratio (CES)									


Clutch size 1966–2007

Bullfinch




Brood size 1966–2007

Bullfinch




Egg stage nest failure rate

Bullfinch




Chick stage nest failure rate

Bullfinch




Laying date 1966–2007

Bullfinch



CES productivity 1983–2007

Bullfinch



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)



Page not found

[home](#) » [birdtrends2008](#) » Page not found



Terribly sorry...


The page you are looking for could not be found.

- Try searching using the search bar (top right)
- Navigate to the area you are looking for using the main menu (above)
- [Return to our homepage](#)
- Email us about what you are trying to find, or to tell us about a broken link - info@bto.org

Share this page



Related content



ARTICLE

Results

See live NBC results and watch the breeding season unfold



CASE STUDY

Climate change in a warming world

BTO science contributes to our understanding of future scenarios, and informing policies and conservation management strategies to help species adapt.



EVENT

Northern Ireland Seabird Report Launch

The NI Seabird Report is unique in the UK, presenting the most up-to-date seabird monitoring data collected by volunteers, NGOs and NIEA from around Northern Ireland. Each March we present freshly produced hard copies...

BTO - Looking out for birds

[About us](#) [Jobs](#) [National offices](#) [Community](#) [Events](#) [Blog](#) [Shop](#)

✉ info@bto.org

[Press releases](#) [Terms & Conditions](#) [Privacy](#) [Cookies](#)



Registered Charity Number 216652 (England & Wales), SC039193 (Scotland)

© British Trust for Ornithology, BTO, The Nunnery, Thetford, Norfolk IP24 2PU, Tel: +44 (0)1842 750050 Fax: +44 (0)1842 750030

Seen a ringed / dead / diseased bird?

[REPORT IT TO US ►](#)

Keep up to date with BTO

[E-NEWSLETTER SIGN UP](#)

Flock together





May we use cookies to monitor performance and improve your experience?

These optional cookies are used for collecting anonymous visit data only.

[READ OUR COOKIE POLICY](#)

ACCEPT ALL COOKIES

[Set preferences](#)

YELLOWHAMMER

Emberiza citrinella

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: no SPEC category (concentrated in Europe, conservation status favourable)

UK: red (>50% population decline)

UK Biodiversity Action Plan: priority species

Long-term trend

UK, England: rapid decline

UK population size

792,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Yellowhammer abundance began to decline on farmland in the mid 1980s and, except in Scotland, the decline has continued ever since. The species, listed as green in 1996, is now red listed. While there is some evidence that survival rates have decreased during the period of decline, Yellowhammer breeding performance has tended to improve ([Siriwardena et al. 1998b, 2000b](#)). However, recent declines in clutch size, brood size and nest success are of NRS concern ([Leech & Barimore 2008](#)). Overall nest failure rates are relatively high, probably because later nests, which tend to be more successful ([Kyrkos 1997](#)), are under-represented in the NRS data set, but this is unlikely to affect overall trends. Reductions in winter seed food availability as a result of agricultural intensification (for example, the loss of winter stubbles and a reduction in weed densities) are widely believed to have contributed to the population decline. [Gillings et al. \(2005\)](#) have identified better population performance in areas with extensive winter stubble, presumably because overwinter survival is relatively high. The local availability of winter setaside is a good predictor of sites chosen for breeding territories the next year ([Whittingham et al. 2005](#)). Numbers have fallen widely in Europe since 1980 ([PECBMS 2007](#)).



Population changes

CBC/BBS UK 1966–2007

Yellowhammer




Table of population changes for Yellowhammer

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	440	-55	-65	-47	>50	
	25	1981-2006	605	-53	-59	-48	>50	
	10	1996-2006	1148	-11	-15	-5		
	5	2001-2006	1123	-4	-8	1		
CBC/BBS England	39	1967-2006	385	-57	-66	-46	>50	
	25	1981-2006	529	-57	-62	-50	>50	
	10	1996-2006	1003	-15	-18	-11		
	5	2001-2006	986	-9	-12	-5		
CES adults	22	1984-2006	18	-74	-91	-28	>50	Small sample
	10	1996-2006	13	-55	-83	1		Small sample
	5	2001-2006	10	-32	-69	44		Small sample
BBS UK	11	1995-2006	1084	-14	-18	-9		
	10	1996-2006	1102	-11	-16	-7		
	5	2001-2006	1123	-4	-7	1		


BBS E	Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
		5	2001-2006	986	-8	-11	-5		
BBS Scotland		11	1995-2006	93	4	-16	24		
		10	1996-2006	95	5	-14	23		
		5	2001-2006	93	11	-2	26		
BBS Wales		11	1995-2006	38	-36	-53	-16	>25	
		10	1996-2006	38	-32	-48	-12	>25	
		5	2001-2006	37	-10	-27	11		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Yellowhammer




CES adult abundance 1983—2007

Yellowhammer




CES juvenile abundance 1983—2007

Yellowhammer




BBS UK 1994—2007

Yellowhammer




BBS England 1994—2007

Yellowhammer




BBS Scotland 1994—2007

Yellowhammer



BBS Wales 1994—2007

Yellowhammer




Productivity trends

Table of productivity changes for Yellowhammer

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	44	Curvilinear	3.35 eggs	3.39 eggs	1.2%	
Brood size	38	1968-2006	67	Curvilinear	2.96 chicks	3.03 chicks	2.3%	
Daily failure rate (eggs)	38	1968-2006	66	Curvilinear	5% nests/day	3.8% nests/day	-24%	
Daily failure rate (chicks)	38	1968-2006	52	Curvilinear	4.62% nests/day	4.23% nests/day	-8.4%	
Laying date	38	1968-2006	27	Linear increase	May 30	Jun 7	8 days	Small sample
Juvenile to Adult ratio (CES)	22	1984-2006	20	Smoothed trend	87 Index value	100 Index value	15%	
Juvenile to Adult ratio (CES)	10	1996-2006	15	Smoothed trend	158 Index value	100 Index value	-37%	Small sample
Juvenile to Adult ratio (CES)	5	2001-2006	12	Smoothed trend	73 Index value	100 Index value	38%	Small sample


Clutch size 1966—2007

Yellowhammer




Brood size 1966—2007

Yellowhammer




Egg stage nest failure rate

Yellowhammer




Chick stage nest failure rate

Yellowhammer




Laying date 1966—2007

Yellowhammer



CES productivity 1983—2007

Yellowhammer



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

REED BUNTING *Emberiza schoeniclus*

- Population changes
- Productivity trends
- Additional information

Conservation listings

Europe: no SPEC category (favourable conservation status in Europe, not concentrated in Europe)

UK: red (>50% population decline)

UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK, England: shallow decline

UK population size

192,000–211,000 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS and WBS trends:

[BiE04](#), [APEP06](#))

Status summary

Red-listing for this species is based on a 62% decline on CBC plots between 1974 and 1999. Both CBC/BBS and WBS indices declined rapidly during the 1970s, but Reed Bunting abundance subsequently remained remarkably stable. In recent years, BBS results indicate significant population increase. The early increase in the CBC index was associated with a gradual spread into drier habitats, especially farmland, and it is likely that the subsequent decline was related to agricultural intensification. Detailed demographic analyses suggest that the decline was driven by decreasing survival rates and that a subsequent population recovery may have been prevented by increased nest losses ([Peach et al. 1999](#)). This is supported by a moderate decline in CES productivity and by a significant increase in failure rates at the egg stage, which has raised NRS concern ([Leech & Barimore 2008](#)). Farmland densities are four times higher in oilseed rape than in cereals or setaside and this crop is crucial in reducing the dependency of the species on wetlands ([Gruar et al. 2006](#)).



Population changes

CBC/BBS UK 1966–2007 Reed Bunting




Table of population changes for Reed Bunting

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	194	-19	-36	3		
	25	1981-2006	242	9	-11	31		
	10	1996-2006	441	30	19	44		
	5	2001-2006	478	26	15	38		
CBC/BBS England	39	1967-2006	153	-22	-41	-1		
	25	1981-2006	187	7	-14	28		
	10	1996-2006	332	30	19	44		
	5	2001-2006	353	29	20	37		
WBS waterways	31	1975-2006	51	-61	-75	-33	>50	
	25	1981-2006	53	-31	-54	17		
	10	1996-2006	53	10	-14	66		
	5	2001-2006	44	16	-1	41		
CES adults	22	1984-2006	59	-51	-64	-36	>50	
	10	1996-2006	68	-25	-36	-12		
	5	2001-2006	59	-2	-12	12		


CES juv Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
	5	2001-2006	43	-26	-40	-9	>25	
BBS UK	11	1995-2006	406	28	17	41		
	10	1996-2006	416	30	18	42		
	5	2001-2006	478	24	16	36		
BBS England	11	1995-2006	303	25	12	34		
	10	1996-2006	309	31	18	40		
	5	2001-2006	353	27	18	36		
BBS Scotland	11	1995-2006	48	42	16	83		
	10	1996-2006	48	41	10	79		
	5	2001-2006	56	21	-7	53		



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966—2007

Reed Bunting




Waterways Bird Survey 1974—2007

Reed Bunting




CES adult abundance 1983—2007

Reed Bunting




CES juvenile abundance 1983—2007

Reed Bunting




BBS UK 1994—2007

Reed Bunting




BBS England 1994—2007

Reed Bunting



BBS Scotland 1994—2007

Reed Bunting




Productivity trends


Table of productivity changes for Reed Bunting

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Clutch size	38	1968-2006	44	None				
Brood size	38	1968-2006	61	None				
Daily failure rate (eggs)	38	1968-2006	52	Linear increase	0.73% nests/day	2.74% nests/day	275.3%	
Daily failure rate (chicks)	38	1968-2006	52	None				
Laying date	38	1968-2006	48	None				
Juvenile to Adult ratio (CES)	22	1984-2006	62	Smoothed trend	217 Index value	100 Index value	-54% >50	
Juvenile to Adult ratio (CES)	10	1996-2006	71	Smoothed trend	171 Index value	100 Index value	-42% >25	
Juvenile to Adult ratio (CES)	5	2001-2006	62	Smoothed trend	127 Index value	100 Index value	-21%	


Clutch size 1966—2007




Brood size 1966—2007




Egg stage nest failure rate
Reed Bunting




Chick stage nest failure rate
Reed Bunting



Laying date 1966—2007



CES productivity 1983—2007



Additional information

- [Maps and statistics from British and Irish atlases](#)
- [BirdFacts page on species biology](#)
- [BirdTrack results](#)
- [Garden BirdWatch results](#)

CORN BUNTING

Emberiza calandra

• Population changes

• Productivity trends

• Additional information

Conservation listings

Europe: SPEC category 2 (declining)
 UK: red (>50% population decline, historical decline)
 UK Biodiversity Action Plan: [click here](#)

Long-term trend

UK, England: rapid decline

UK population size

8,500–12,200 territories in 2000 (1988–91 Atlas estimate updated using CBC/BBS trend: [BIE04](#), [APEP06](#))

Status summary

Following an earlier, historical decrease, Corn Buntings declined very steeply between the mid 1970s and mid 1980s, with local extinctions across large sections of their former range. Subsequently the decline has continued, but at a much-reduced rate. Breeding performance per nesting attempt has increased considerably over this period ([Crick 1997](#)), but it is also reported that fewer birds now raise a second brood, thus reducing productivity overall ([Brickle & Harper 2002](#)). Brood size and nest survival at the chick stage are currently of NRS concern ([Leech & Barimore 2008](#)). Ring-recovery sample sizes do not permit an analysis of survival rates ([Siriwardena et al. 1998b, 2000b](#)). Any decrease there has been in survival rates is probably a result of the deleterious effects of agricultural intensification on seed availability in winter ([Donald 1997](#)). The isolated Corn Bunting population on the Western Isles is still declining rapidly, probably because agricultural change has reduced the supply of winter grain ([Wilson et al. 2007](#)). Targeted restoration of lower-intensity cultivation, but without hedgerows, might help prevent further local extinctions ([Mason & Macdonald 2006](#)). Management interventions in eastern Scotland were shown to have the potential to halt, or perhaps reverse, the Corn Bunting decline there ([Perkins et al. 2008](#)). Corn Buntings are estimated to have declined by 61% across Europe during 1980–2005 ([PECBMS 2007](#)). With declines across much of its European range, this previously 'secure' species is now provisionally evaluated as 'declining' ([BirdLife International 2004](#)).



Population changes

CBC/BBS UK 1966–2007 Corn Bunting




Table of population changes for Corn Bunting

Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
CBC/BBS UK	39	1967-2006	63	-87	-94	-77	>50	
	25	1981-2006	80	-84	-92	-74	>50	Small CBC sample
	10	1996-2006	144	-27	-38	-15	>25	
	5	2001-2006	126	-6	-21	8		
CBC/BBS England	39	1967-2006	60	-84	-93	-74	>50	
	25	1981-2006	76	-83	-92	-71	>50	Small CBC sample
	10	1996-2006	138	-22	-36	-8		
	5	2001-2006	120	0	-16	20		
BBS UK	11	1995-2006	138	-32	-45	-21	>25	
	10	1996-2006	138	-26	-39	-16	>25	
	5	2001-2006	126	-7	-21	6		
BBS England	11	1995-2006	132	-28	-40	-18	>25	


Source	Period (yrs)	Years	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
--------	--------------	-------	-----------	------------	-------------	-------------	-------	---------



The Breeding Bird Survey is jointly funded by the BTO, JNCC & RSPB


CBC/BBS England 1966–2007

Corn Bunting




BBS UK 1994–2007

Corn Bunting



BBS England 1994–2007

Corn Bunting



Productivity trends


Table of productivity changes for Corn Bunting

Variable	Period (yrs)	Years	Mean annual sample	Trend	Modelled in first year	Modelled in 2006	Change	Comment
Brood size	38	1968-2006	12	Curvilinear	3.07 chicks	2.89 chicks	-5.9%	Small sample
Daily failure rate (eggs)	38	1968-2006	11	None				Small sample
Daily failure rate (chicks)	38	1968-2006	11	Curvilinear	4.5% nests/day	2.24% nests/day	-50.2%	Small sample
Laying date	38	1968-2006	13	None				Small sample

Insufficient data on clutch size available for this species


Brood size 1966–2007

Corn Bunting




Egg stage nest failure rate

Corn Bunting




Chick stage nest failure rate

Corn Bunting



Laying date 1966–2007

Corn Bunting



Insufficient data on CES available for this species

Additional information

- [Maps and statistics from British and Irish atlases](#)

- [BirdFacts page on species biology](#)
- [BirdTrack results](#)

3. Help on species accounts

The species in this report can be accessed in any order, but the species index and drop-down list use the taxonomic sequence established by the British Ornithologists' Union in its [British List](#). The vernacular and scientific names we use are also drawn from that list. Given this report's limited geographical scope, we have used the British rather than the international English names. Depending on the availability of data (not every species is covered by each scheme), the following will be found beneath each species heading:

- 1) **Conservation listings:** First, the European conservation category is given, according to current listings by BirdLife International in [Birds in Europe \(BirdLife International 2004\)](#). These update the original listings of [Tucker & Heath \(1994\)](#). For SPECs (Species of European Conservation Concern), the European Threat Status is also given. The current SPEC categories are as follows:

- SPEC 1** Species of global conservation concern, according to the latest assessments by BirdLife International (www.birdlife.org/datazone/species/index.html)
- SPEC 2** Species with an unfavourable European conservation status, and with more than half of the global breeding or wintering population concentrated in Europe
- SPEC 3** Species with an unfavourable European conservation status, but with less than half of the global breeding or wintering population within Europe

Other species, not considered to be of European conservation concern, and assessed as 'secure', have no SPEC category but are placed into two further groupings:

- Species with a favourable European conservation status, and with less than half of the breeding or wintering population within Europe (**Non-SPEC**)
- Species with a favourable European conservation status, but with more than half of the global breeding or wintering population concentrated in Europe (**Non-SPEC^E**)

The UK conservation listing, given next, is taken from *The Population Status of Birds in the UK* ([Gregory et al. 2002](#); see [PSoB](#) pages). These supersede the previous *Birds of Conservation Concern* listings ([Gibbons et al. 1996](#)), and cover the period 2002–07. There are three categories, as follows:

- Red** high conservation concern
- Amber** medium conservation concern
- Green** all other species (except introduced species, which are not classified)

The main reason or reasons for listing as red or amber are also given. NB:

- SPEC 1 (globally threatened) species are automatically red listed, and SPEC 2 or 3 species are amber listed (unless they are introduced or a red-list criterion applies)
- Red or amber listing may stem from decline, localisation or importance of non-breeding as well as breeding populations in the UK
- Rates of population decline used to assess red and amber listing are generally derived from CBC results for the 25-year period 1974–99
- Range declines are generally calculated from the numbers of 10-km squares occupied in the two published [breeding atlases \(Gibbons et al. 1993\)](#)
- Historical decline (in UK over the period 1800–1995) is assessed by literature review

Following the signing of the Convention on Biological Diversity at the 'Earth Summit' in Rio de Janeiro in 1992, the statutory conservation bodies in the UK compiled [Biodiversity Action Plans \(BAPs\)](#) for 26 rare or threatened bird species, of which 12 are covered by this report. A [BAP review](#) published in 2007 has concluded that 56 UK bird species now qualify for BAPs and has recommended that certain subspecies (e.g. Fair Isle and St Kilda Wrens) should now be included. This report covers 31 of those species.

Where a UK BAP exists, we give the link to the latest available version. You will find onward links, for example to local BAPs for that species. For species newly nominated, we record that a BAP is 'in preparation'.

- 2) **Long-term trend:** This summarises the trend in population size since 1975 from WBS data, 1984 from CES data, or 1967 from CBC/BBS, with reference to any CBC/BBS, WBS or CES data that may be tabulated. If there are no data available from these schemes, any assessment of trends covers the period since about the mid 1960s, but may also take historical data into account. Increases and declines that are qualified as 'shallow', 'moderate' or 'rapid' are generally statistically significant. The following terms are used:

- **Rapid decline:** >50% population decline from CBC/BBS, WBS or CES
- **Moderate decline:** 25–50% population decline from CBC/BBS, WBS or CES
- **Shallow decline:** 10–25% population decline from CBC/BBS, WBS or CES
- **Decline/Increase:** information has been derived from other sources
- **Probable/Possible increase/decline:** as above, but the information is not as certain - see the status summary for reasons
- **Stable/Fluctuating, with no long-term trend:** no overall change, or change <10%
- **Uncertain:** where the information from two monitoring schemes conflicts or if the data are

unrepresentative of the species' total UK population

- **Unknown:** no information on the UK population trend is available
- **Shallow increase:** 10–50% population increase from CBC/BBS, WBS or CES
- **Moderate increase:** 50–100% population increase from CBC/BBS, WBS or CES
- **Rapid increase:** >100% population increase from CBC/BBS, WBS or CES

- 3) **UK population size:** Periodic reports on population sizes of birds in Britain and in the UK, for the breeding season and for winter, are agreed by the Avian Population Estimates Panel (APEP), on which BTO, GCT, JNCC, RSPB and WWT are represented. Extracts from the Panel's second report (Baker *et al.* 2006) are given for each of our species, with a shortened reference (APEP06). The second edition of *Birds in Europe* (BirdLife International 2004) was published while APEP06 was in preparation. Their figures are also given, referenced as BiE04. The units and reference year (or period) is given for each estimate, and where possible its derivation is also described briefly or referenced. BiE04 and APEP06 estimates are usually identical, but may differ because:
- one or other is updated to a new reference year
 - the two publications apply different rules for inclusion of introduced species
 - BiE04 figures include the Channel Islands (but for most species this has no effect on the estimate)
 - different methods of rounding or range estimation have been applied to the same original data
 - sources used for BiE04, but not APEP06, included papers in preparation

Information too recent to have been included in either of these publications is also given, pending ratification by APEP. Readers should note that the wide ranges given for many species reflect the considerable uncertainty that applies to all but a few of the current estimates. The application of distance sampling methods to BBS data (Newson *et al.* 2005, 2008), or future surveys, including the current 2007–11 Atlas, may well result in substantial challenge to the presently accepted figures.

- 4) **Status summary:** This section provides a brief summary of the trends detailed for the species and indicates why such changes might have occurred, with reference to any published information, if this is known.
- 5) **Population trend graphs:** The first, large graph shows the most representative long-term trend in abundance for the species, and is followed after the table by further graphs from other schemes, including BBS graphs for separate UK countries, as available. If no suitable long-term trend is available then the BBS trend for the UK is shown. **Methods (Section 2)** provides details about how the trend data are calculated for each scheme. For BBS, CBC/BBS, CBC, WBS and CES, the graphs show a smoothed line (in blue) and its 85% confidence limits (in green); for the Heronries Census, annual estimates are shown in blue, 85% confidence limits in green, and a smoothed trend in red.
- 6) **Population trends table:** This table provides details of summarised percentage changes in population size, over the maximum period from each source, and from the past 25 years, 10 years and 5 years, where these figures are available. Further columns indicate the years included, the average number of census plots included in the analysis for each year, the percentage change (an increase if presented with no sign) and the upper and lower 90% confidence limits of that change. Where the confidence interval does not include zero change, population declines are regarded as statistically significant. The '**Alert**' column indicates where a statistically significant population decline is estimated to be of 50% or more (>50) or between 25% and 50% (>25) (see **Alerts, Section 2.8** for further details). The '**Comment**' column lists any caveats that must be considered when interpreting the estimates. The caveats include:
- **Small sample:** For CBC, WBS and CES data, a mean sample size of less than 20 (but more than 10) census plots was available; for BBS data from individual countries, a mean sample of less than 40 (but more than 30) plots was available.
 - **Unrepresentative?:** Where joint CBC/BBS trends are reported, the trends are always considered to be representative for the region concerned. The CBC data may inadequately represent the population as a whole. This judgment was made either because the species' average abundance in 10-km squares containing CBC plots was less than that in other occupied 10-km squares, as measured by Breeding Atlas timed counts or frequency indices (Gibbons *et al.* 1993), or, where these figures could not be calculated, on expert opinion.
- 7) **Productivity graphs:** Graphs from Constant Effort Sites Scheme or Nest Record Scheme data illustrate trends in productivity. For NRS data, annual means (averages) are shown in green, with error bars to denote ± 1 standard error; quadratic or linear regression lines (in black) and the upper and lower 95% confidence limits of these lines (in blue) are also shown. For CES data, the smoothed trends are plotted (in blue) with their 85% confidence limits (in green) (see **Section 2.6** for details).
- 8) **Productivity trends table:** This provides details of changes in productivity since 1968 (or a more recent year, depending on the availability of data). It lists the period of years concerned, the mean annual sample, the type of trend ('curvilinear' is for a significant quadratic trend, 'linear' is for a significant linear trend, 'none' is where the linear trend is not significantly different from horizontal), the modelled values (from the appropriate regression) for the first and last years and their difference (where the trend is significant), and any caveats that must be considered when interpreting the data. Changes are presented either in the units given or as percentages, and are increases unless a minus sign is shown. The caveat '**Small sample**' is given when the mean number of nest record cards contributing annually was in the range 10–30, or when the mean annual number of CES plots recording the species was less than 20 (but more than 10).
- 9) **Additional information:** Provides links to atlas maps and tables from previous atlas surveys, and the relevant pages of BirdFacts, BirdTrack and Garden BirdWatch, as available, from the BTO web site. Atlas maps from earlier surveys are not yet available online for Red-throated Diver, Goosander, Hen Harrier, Buzzard, Hobby and Peregrine, for which some of the original data were confidential (see **previous atlases species help**).

Tip: use the 'Species quick links' box at top of each page to navigate the species pages

Section 4 – Discussion

[Back to Contents](#)

4. Discussion

- 4.1 [The alert system](#)
- 4.2 [Latest long-term alerts](#)
- 4.3 [Ten-year trends and evidence of species recovery](#)
- 4.4 [Increasing species](#)
- 4.5 [Changes in breeding performance](#)
- 4.6 [Conclusion](#)

[Return to Contents](#)

4.1 The alert system

This report uses a system of 'alerts' that has been agreed between the providers and users of population monitoring information in the UK. The system provides alerts to population declines of 25–50% and of >50% over short, medium and longer terms (5 years, 10 years and 25+ years respectively). These help to highlight the scale and timing of declines, and act as an aid to interpreting the trend graphs presented. Our main emphasis is on long-term declines measured over the longest period available (usually 39 years) and over 25 years, which is the period that is normally used to determine red and amber listing (Gregory *et al.* 2002). Alerts triggered over the short term for individual species should be considered as early warnings, indicating that conservation issues may be developing for these species. However, it is possible that such declines may be due to chance fluctuations in abundance from which the population is able to recover without assistance. The rapid, short-term decline of a suite of similar species should be considered as a stronger indication that potential problems may be developing. Details of the alerts and methodology used in this report are given in the [methods section](#).

These alerts are therefore important for the conservation practitioners who need to set priorities for conservation action, but we also hope that they will prove of more general use to other readers of the report. Similar [alerts for wetland birds](#) are now provided by the Wetland Bird Survey (Maclean & Austin 2008).

In this discussion we:

- 1) Review the latest population change measures and alerts for species that are currently on the [Population Status of Birds](#) (PSoB) [red](#) or [amber](#) lists (declines only) for the UK (Gregory *et al.* 2002).
- 2) Identify species that are not currently on these PSoB lists that have raised alerts on account of long-term declines, and also those species on the list where recovery may be sufficient to downgrade their listing status in the future.
- 3) Briefly review declines along waterways and in scrub and wetland habitats as shown by the WBS and CES schemes.
- 4) Review trends over the last 10 years in species that have shown long-term declines, to identify the extent of ongoing declines and any evidence of recovery.
- 5) Identify those species that have shown rapid long-term population increases.
- 6) Discuss patterns of changes in breeding performance and relationships between trends in abundance and breeding performance.
- 7) Summarise the overall patterns found.

Except where otherwise indicated our discussion is based on the best long-term trend that is available for each species. These are the trends presented as the main trend graph for each species. Details of estimating and comparing trends are given in the [methods section](#). Full details of all trends available for each species are given on the [species pages](#). Summary tables of all alerts raised by each scheme are presented in the [appendices](#).

It should be noted that a number of species included in the PSoB red and amber lists are not covered by this report. Thus tables relating to PSoB list status do not include every species on the relevant PSoB list.

[Return to previous page](#)

[Next page – 4.2 Latest long-term alerts](#)

4.2 Latest long-term alerts

4.2.1 Long-term trends of PSoB red-listed species

The species considered here were **red-listed** due to long-term declines of more than 50% over the 25-year period 1974–99. The latest long-term population changes and alerts over the maximum period available (usually 39 years) and over 25 years are shown in Table 4.2.1. The species are listed in descending order of long-term percentage change.

The results confirm the declining status of all of the 16 species concerned (although the 25-year change for **Reed Bunting** is now a non-significant increase). All these updated changes fire alerts, except for the 25-year change for **Song Thrush**, both changes for **Reed Bunting** and the 31-year change for **Lesser Spotted Woodpecker** (which, although apparently large, has very wide confidence limits and is thus not statistically significant). **Linnet**, **Marsh Tit**, **Skylark** and **Song Thrush** now show declines of less than 50% over the most recent 25-year period, partly reflecting the fact that their long-term declines started more than 25 years ago. **Bullfinch** and **Reed Bunting** now have both long-term and 25-year declines of below 50%. Populations of both species increased between the late 1960s and the mid 1970s, before the rapid declines that gave rise to their current conservation listing.

Table 4.2.1 Latest trends for red-listed species

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Tree Sparrow	39	CBC/BBS England	-97	-99	-94	>50	
Tree Sparrow	25	CBC/BBS England	-93	-97	-88	>50	
Grey Partridge	39	CBC/BBS UK	-88	-91	-83	>50	
Grey Partridge	25	CBC/BBS UK	-78	-85	-71	>50	
Corn Bunting	39	CBC/BBS UK	-87	-94	-77	>50	
Corn Bunting	25	CBC/BBS UK	-84	-92	-74	>50	
Spotted Flycatcher	39	CBC/BBS UK	-86	-90	-80	>50	
Spotted Flycatcher	25	CBC/BBS UK	-81	-86	-75	>50	
Turtle Dove	39	CBC/BBS UK	-85	-90	-76	>50	
Turtle Dove	25	CBC/BBS UK	-82	-88	-75	>50	
Willow Tit	39	CBC/BBS UK	-85	-93	-72	>50	
Willow Tit	25	CBC/BBS UK	-82	-90	-69	>50	
Starling	39	CBC/BBS England	-83	-88	-77	>50	
Starling	25	CBC/BBS England	-78	-83	-73	>50	
Linnet	39	CBC/BBS England	-75	-81	-66	>50	
Linnet	25	CBC/BBS England	-47	-57	-32	>25	
House Sparrow	29	CBC/BBS England	-70	-79	-60	>50	
House Sparrow	25	CBC/BBS England	-59	-71	-49	>50	
Marsh Tit	39	CBC/BBS UK	-68	-77	-56	>50	
Marsh Tit	25	CBC/BBS UK	-38	-51	-16	>25	
Lesser Spotted Woodpecker	31	CBC to 1999	-60	-81	40		Small sample
Lesser Spotted Woodpecker	25	CBC to 1999	-73	-86	-31	>50	Small sample
Skylark	39	CBC/BBS England	-59	-66	-51	>50	
Skylark	25	CBC/BBS England	-47	-54	-40	>25	
Yellowhammer	39	CBC/BBS UK	-55	-65	-47	>50	
Yellowhammer	25	CBC/BBS UK	-53	-59	-48	>50	
Song Thrush	39	CBC/BBS UK	-51	-58	-43	>50	
Song Thrush	25	CBC/BBS UK	-16	-27	-5		
Bullfinch	39	CBC/BBS UK	-50	-61	-37	>25	
Bullfinch	25	CBC/BBS UK	-28	-39	-14	>25	
Reed Bunting	39	CBC/BBS UK	-19	-36	3		
Reed Bunting	25	CBC/BBS UK	9	-11	31		

See [PSoB](#) pages for information on red and amber criteria

4.2.2 Long-term trends of PSoB amber-listed species

Most of the species considered here were **amber-listed** due to long-term declines of more than 25%, but less than 50%, over the 25-year period 1974–99. In addition, three species where the best trend estimate was a decline of more than 50% were listed amber, rather than red, on the grounds that the census data were sparse or maybe unrepresentative. The latest long-term population changes and alerts over the maximum period available (usually 39 years) and over 25 years are shown in Table 4.2.2. The species are listed in descending order of long-term percentage change. These results confirm the declining status of most of the 15 species concerned.

Table 4.2.2 Latest trends for declining amber-listed species

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Lesser Redpoll	39	CBC/BBS England	-90	-96	-78	>50	
Lesser Redpoll	25	CBC/BBS England	-95	-98	-91	>50	
Tree Pipit	39	CBC/BBS England	-83	-92	-71	>50	
Tree Pipit	25	CBC/BBS England	-82	-91	-69	>50	
Woodcock	31	CBC to 1999	-74	-88	-49	>50	Small sample
Woodcock	25	CBC to 1999	-76	-88	-51	>50	Small sample
Yellow Wagtail	39	CBC/BBS UK	-74	-88	-46	>50	
Yellow Wagtail	25	CBC/BBS UK	-70	-82	-55	>50	
Cuckoo	39	CBC/BBS England	-61	-71	-48	>50	
Cuckoo	25	CBC/BBS England	-61	-67	-54	>50	
Willow Warbler	39	CBC/BBS England	-58	-70	-46	>50	
Willow Warbler	25	CBC/BBS England	-58	-67	-49	>50	
House Martin	39	CBC/BBS England	-56	-86	42		
House Martin	25	CBC/BBS England	-60	-88	105		
Redshank	31	WBS waterways	-49	-85	-16	>25	Small sample
Redshank	25	WBS waterways	-45	-76	-22	>25	Small sample
Meadow Pipit	39	CBC/BBS England	-46	-74	-24	>25	
Meadow Pipit	25	CBC/BBS England	-43	-61	-26	>25	
Mistle Thrush	39	CBC/BBS UK	-43	-51	-31	>25	
Mistle Thrush	25	CBC/BBS UK	-38	-45	-31	>25	
Lapwing	39	CBC/BBS UK	-34	-64	-5	>25	
Lapwing	25	CBC/BBS UK	-53	-65	-37	>50	
Dunnock	39	CBC/BBS UK	-33	-42	-22	>25	
Dunnock	25	CBC/BBS UK	-6	-17	5		
Grey Wagtail	31	WBS waterways	-23	-44	0		
Grey Wagtail	25	WBS waterways	35	9	60		
Kestrel	39	CBC/BBS England	6	-23	53		
Kestrel	25	CBC/BBS England	-21	-35	-1		
Goldcrest	39	CBC/BBS England	42	-20	197		
Goldcrest	25	CBC/BBS England	-18	-38	14		

See [PSoB](#) pages for information on red and amber criteria

Six species show significant declines of greater than 50% and could thus be candidates for red listing at the next review. Three of these, **Lesser Redpoll**, **Tree Pipit** and **Woodcock**, were listed amber rather than red in 2002 as a result of uncertainty about the reliability of the data, and there has been no substantial change in the information available on their declines. The others, **Yellow Wagtail**, **Willow Warbler** and **Cuckoo**, have been subject to ongoing declines that have since passed the 50% threshold (although it should be noted that for the last two species the long-term data are from England only). The serious nature of the **Yellow Wagtail** decline is supported by data from both WBS (-97% over 31 years) and BBS (-44% over 11 years). BBS data indicate that **Willow Warblers** and **Cuckoos** have continued to decline in England and Wales over the last 11 years, but have shown more stability in Scotland. Scottish Willow Warblers have shown a significant decline over the recent five years, however.

Our best estimate of long-term change in the English **House Martin** population now also shows a decline of more than 50%, but statistically it is not significantly different from no change. Thus no alerts are raised for this species. **House Martin** is probably best regarded as data deficient rather than as a candidate for red listing. BBS data indicate that its numbers have been stable or increasing since 1994.

Dunnock ceased its decline in the early 1990s, and has subsequently shown significant increase: consequently, its 25-year trend shows no significant change. **Grey Wagtails** have also been increasing recently, especially since the late 1990s, and as a result their 25-year change is now +28%, while the decline over the longest period for which we can measure changes in their populations (31 years) is just 23%. If the positive trend continues they might be removed from the amber list at a future revision. **Kestrel** and **Lapwing** show an opposite pattern to **Grey Wagtail**, with smaller declines over 39 years than 25 years, reflecting modest increases prior to the declines that are now a cause of concern. For **Kestrel**, the 39-year trend shows little overall change. For **Lapwing**, the 25-year decline now exceeds 50%, which could make it a candidate for future red-listing, and it has already been promoted to a priority species on the UK Biodiversity Action Plan. **Goldcrest** is a difficult species for status assessments because its populations show such wide fluctuations. Numbers increased by 42% over the 39-year period and decreased by only 18% over 25 years, neither change being statistically significant. More recently, BBS data show that numbers have increased by 25% over the last 11 years, and it is questionable that the status of this species should be of particular concern.

4.2.3 Long-term declines of species that are not currently red or amber listed (for declines)

We have identified eight species that are currently showing long-term declines of greater than 25% but are not currently included on either the red or amber lists (Table 4.2.3). Should their downward trends continue, these species may be possible candidates for red or amber listing at the next review.

The position of **Whitethroat** at the head of this list is somewhat misleading. Much of the recorded 62% decline is the result of the well-documented crash between 1968 and 1969 ([Winstanley et al. 1974](#)). The 25-year change for this species is a 65% increase, representing a partial recovery to former levels. The **Little Grebe** data should be treated with caution as they are based on a small sample from linear waterways. WBS shows an ongoing decline in this habitat over the last ten years while BBS, which is likely to cover a more representative set of habitat types for this species, shows an increase over the same period.

Evidence for a **Little Owl** decline is statistically significant for both the 39-year and 25-year periods. **Tawny Owls** have shown a very slow decline since the early 1970s, which became more rapid around 1999, and now show a decline of 25% which just reaches statistical significance. These figures must be treated with some caution, however, because CBC and BBS census techniques are not designed with nocturnal and crepuscular species in mind.

Reed Warbler shows a decline of 31% over 21 years in its core habitat of large reedbeds, based on CES data. In clear contrast, however, CBC/BBS and WBS indices, both show strong and ongoing significant increase in this species, and are supported by atlas data showing range expansion. Further investigation is required of how the population changes of this species may differ with respect to region and habitat. For **Sedge Warbler**, a strong decline is indicated by CBC/BBS, WBS and CES schemes but is statistically significant only for CES.

Table 4.2.3 Long-term trends for declining species not on the red or amber list (for declines)

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Whitethroat	39	CBC/BBS UK	-62	-73	-50	>50	
Little Grebe	31	WBS waterways	-59	-87	-9	>50	Small sample
Little Grebe	25	WBS waterways	-64	-87	-23	>50	Small sample
Little Owl	39	CBC/BBS UK	-46	-68	-12	>25	
Little Owl	25	CBC/BBS UK	-46	-63	-29	>25	
Reed Warbler	22	CES adults	-33	-46	-10	>25	
Curlew	39	CBC/BBS England	-29	-68	25		
Curlew	25	CBC/BBS England	-31	-66	10		
Common Sandpiper	25	WBS waterways	-28	-44	-10	>25	
Sedge Warbler	39	CBC/BBS UK	-28	-58	18		
Tawny Owl	25	CBC/BBS UK	-25	-43	-2		

See [PSoB](#) pages for information on red and amber criteria

Two wader species, **Common Sandpiper** and **Curlew**, also appear in the table. Declines for Curlew have wide confidence intervals and are not matched by WBS data, which show increase. Common Sandpipers were recorded by WBS as increasing up to the mid 1980s, and a smaller decline (22%) is therefore estimated for the 31-year period.

4.2.4 Declines on WBS plots

The **Waterways Bird Survey** supplements the results from more broadly based schemes, such as CBC and BBS, by measuring trends in the bird populations alongside linear waterways. For a few waterways habitat specialists, such as **Grey Wagtail** and **Common Sandpiper**, WBS provides our best information on population trends but for 20 or so others it provides supplementary information from this sensitive habitat. Long-term declines of greater than 25% recorded from WBS plots are listed in Table 4.2.4.

Table 4.2.4 Population declines of greater than 25% recorded by the Waterways Bird Survey between 1975 and 2006

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Yellow Wagtail	31	WBS waterways	-97	-99	-93	>50	Small sample
Reed Bunting	31	WBS waterways	-61	-75	-33	>50	
Little Grebe	31	WBS waterways	-59	-87	-9	>50	Small sample
Pied Wagtail	31	WBS waterways	-53	-66	-43	>50	
Redshank	31	WBS waterways	-49	-85	-16	>25	Small sample

The trends for **Yellow Wagtail** and **Reed Bunting** are consistent in direction with those reported from CBC/BBS, but in each case are more severe. The trend for **Little Grebe** is discussed in [section 4.2.3](#). The **Pied Wagtail** decline of 53% is intriguing because it contrasts markedly with recent increases as measured by CBC/BBS. Over the 25-year period 1981–2006, **Pied Wagtails** declined by 39% on linear waterways, compared with no change in the UK as shown by the CBC/BBS trend. The cause of its decline along waterways is currently unknown. For **Redshank**, WBS has provided valuable data to support the amber listing of this species.

A full set of alerts raised by WBS, and long-term increases detected by that scheme, are tabulated in [section 7.2](#).

4.2.5 Declines on CES plots

The **Constant Effort Sites Scheme** provides trends from standardised ringing in scrub and wetland habitats. It is our best scheme for monitoring certain bird populations inhabiting reed beds but its main objective is to collect integrated data on relative abundance, productivity and survival for a suite of species. The longest trends currently available from the CES cover a period of 22 years (Table 4.2.5).

Table 4.2.5 Population declines of greater than 25% recorded by the Constant Effort Sites Scheme between 1984 and 2006

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Lesser Redpoll	22	CES adults	-96	-99	-91	>50	Small sample
Linnet	22	CES adults	-92	-97	-84	>50	
Yellowhammer	22	CES adults	-74	-91	-28	>50	Small sample
Willow Warbler	22	CES adults	-61	-70	-52	>50	
Lesser Whitethroat	22	CES adults	-59	-78	-40	>50	
Willow Tit	22	CES adults	-58	-89	-25	>50	
Reed Bunting	22	CES adults	-51	-64	-36	>50	
Whitethroat	22	CES adults	-36	-57	-22	>25	
Sedge Warbler	22	CES adults	-33	-49	-11	>25	
Reed Warbler	22	CES adults	-33	-46	-10	>25	

Most of the species that are declining on CES sites also show similar trends from CBC/BBS data. **Linnet**, **Yellowhammer**, **Willow Tit**, **Reed Bunting** and **Song Thrush** are already red listed while **Lesser Redpoll** and **Willow Warbler** are amber listed. The decline of **Whitethroat** has also been discussed above ([section 4.2.3](#)). Both **Whitethroat** and **Lesser Whitethroat** are doing less well on CES sites than in the UK as a whole. Over the ten-year period 1996–2006, **Whitethroats** increased by 14% in the UK but decreased by 37% at CES sites. Similarly **Lesser Whitethroats** increased by 8% in the UK as a whole but decreased by 49% on CES sites. Longer-term comparisons show a similar picture. Numbers of juveniles captured at CES sites show very similar patterns of decline to adult captures for both species. It is unclear why these two species are doing so poorly on CES sites as many of these are located in the good-quality scrub habitats that are preferred by these species.

Reed Warbler and **Sedge Warbler** present further similar cases. CES has both as declining enough to raise an alert, whereas over similar time periods CBC/BBS records little change for **Sedge Warbler** and substantial increase for **Reed Warbler**. Increase for the latter species, also noted by WBS, accords better with the considerable range extensions recorded by [atlas projects](#).

A full set of alerts raised by CES, and long-term increases detected by that scheme, are tabulated in [section 7.3](#).

[Return to previous page](#)

[Next page – 4.3 Ten-year trends and evidence of species recovery](#)

4.3 Ten-year trends and evidence for species recovery

If the status of species that have shown long-term declines were now improving, we would expect to find trends to be more positive in recent years compared with the earlier part of the time series. To examine this, we list in Table 4.3.1 the best change estimates over the most recent ten-year period for which we have data (1996–2006) for all of the declining species listed in the previous section of this report (section 4.2). The table also includes four species, **Wood Warbler** (amber), **Red Grouse** (amber), **Grasshopper Warbler** (red) and **Snipe** (amber) for which we can report ten-year trends but lack reliable data covering longer periods.

Table 4.3.1 Ten year trends for species that have shown long-term declines

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Tit	10	CBC/BBS UK	-64	-72	-51	>50	
Turtle Dove	10	CBC/BBS UK	-58	-62	-51	>50	
Wood Warbler	10	BBS UK	-53	-67	-36	>50	
Lesser Spotted Woodpecker	10	CBC to 1999	-51	-75	-22	>50	Small sample
Cuckoo	10	CBC/BBS England	-46	-49	-43	>25	
Yellow Wagtail	10	CBC/BBS UK	-42	-53	-31	>25	
Woodcock	10	CBC to 1999	-40	-62	-11	>25	Small sample
Grey Partridge	10	CBC/BBS UK	-39	-48	-31	>25	
Starling	10	CBC/BBS England	-36	-39	-32	>25	
Lesser Redpoll	10	CBC/BBS England	-33	-51	13		
Redshank	10	WBS waterways	-33	-51	-12	>25	Small sample
Spotted Flycatcher	10	CBC/BBS UK	-32	-38	-16	>25	
Little Grebe	10	WBS waterways	-30	-72	22		Small sample
Tree Pipit	10	CBC/BBS England	-30	-47	-7	>25	
Willow Warbler	10	CBC/BBS England	-30	-36	-24	>25	
Corn Bunting	10	CBC/BBS UK	-27	-38	-15	>25	
Linnet	10	CBC/BBS England	-27	-32	-21	>25	
Reed Warbler	10	CES adults	-26	-36	-13	>25	
Little Owl	10	CBC/BBS UK	-20	-33	-7		
Red Grouse	10	BBS UK	-20	-31	-7		
Curlew	10	CBC/BBS England	-19	-26	-12		
Marsh Tit	10	CBC/BBS UK	-16	-28	-2		
Tawny Owl	10	CBC/BBS UK	-15	-29	5		
House Sparrow	10	CBC/BBS England	-13	-18	-7		
Skylark	10	CBC/BBS England	-13	-17	-9		
Lapwing	10	CBC/BBS UK	-12	-23	-3		
Bullfinch	10	CBC/BBS UK	-11	-17	-3		
Common Sandpiper	10	WBS waterways	-11	-28	5		
Yellowhammer	10	CBC/BBS UK	-11	-15	-5		
Grasshopper Warbler	10	BBS UK	-6	-32	13		
Mistle Thrush	10	CBC/BBS UK	-6	-11	2		
Meadow Pipit	10	CBC/BBS England	-5	-15	6		
Sedge Warbler	10	CBC/BBS UK	-3	-18	11		
House Martin	10	CBC/BBS England	5	-4	14		
Kestrel	10	CBC/BBS England	5	-2	14		
Whitethroat	10	CBC/BBS UK	14	7	20		
Goldcrest	10	CBC/BBS England	22	11	35		
Dunnock	10	CBC/BBS UK	23	18	28		
Song Thrush	10	CBC/BBS UK	23	17	29		
Tree Sparrow	10	CBC/BBS England	24	7	54		
Snipe	10	BBS UK	29	14	47		
Reed Bunting	10	CBC/BBS UK	30	19	44		
Grey Wagtail	10	WBS waterways	32	14	48		

See [PSoB](#) pages for information on red and amber criteria

The 43 species listed include 17 from the red list, 18 declining species that are amber listed on account of population declines and eight species that are not formally listed as declining. The eight species

include **Curlew** which is already amber listed for reasons concerned with its European status.

Eight species at the foot of the table, **Grey Wagtail, Reed Bunting, Snipe, Tree Sparrow, Song Thrush, Dunnock, Goldcrest** and **Whitethroat**, show clear positive trends over the last ten years. The increases in the red-listed **Reed Bunting** and **Song Thrush** are particularly encouraging, as are the positive trends for the amber-listed **Grey Wagtail** and **Dunnock**. However, the most recent figures for **Song Thrush** and **Grey Wagtail** suggest that their recoveries may be levelling off well short of their previous population levels. Similarly while the BBS shows a 29% increase in **Snipe** over the last ten years, much of the former range across lowland Britain lost since the 1960s remains unoccupied, and moreover the population has been declining again since 2003. **Whitethroat** numbers have increased steadily since the mid 1980s but are still far below the population level prior to the 1968/69 crash. The increase in **Tree Sparrow** numbers is very welcome but is coming from such a low level that numbers remain far below those of the mid 1970s, with the population trend graph still showing little sign of a clear recovery.

The rate of decline of 25% over 25 years that is used as a threshold for amber listing is equivalent to a decline of 10.9% over ten years (assuming constant annual rates of change). A further nine species, **Yellowhammer, Sedge Warbler, Common Sandpiper, Meadow Pipit, Bullfinch, Mistle Thrush, Kestrel, Grasshopper Warbler** and **House Martin** have population changes of between +11% and -11% over the last ten years. Where these changes are not statistically significant (eight species), these populations are best regarded as stable. Thus our data suggest that the declines of these species appear to be levelling off, although on this time scale there is as yet no indication of recovery. The position of **Yellowhammer**, with a significant decrease of 11%, is less certain.

Ten-year changes for the remaining 26 species in Table 4.3.1 indicate ongoing declines, with rates equivalent to at least 25% over 25 years. Four species, **Willow Tit, Turtle Dove, Wood Warbler** and **Lesser Spotted Woodpecker**, have declined by more than 50% over the last ten years alone. For **Lesser Spotted Woodpecker**, the ten years in question are 1989–99, since when the species has become too scarce for BBS to monitor. A further 12 species have declined by more than 25% over the last ten years alone. The ongoing declines of so many of the species listed in Table 4.3.1 must be a cause of serious conservation concern.

[Return to previous page](#)

[Next page – 4.4 Increasing species](#)

4.4 Increasing species

Population changes of species for which our best trend estimate from CBC/BBS (usually over 39 years) or from WBS (usually over 31 years) shows an increase of more than 50% are shown in Table 4.4.1 below. There are 25 species included, of which 17 have at least doubled their population size over the decades under review. Four groups of species stand out: corvids – **Carrion Crow**, **Magpie** and **Jackdaw**; doves – **Collared Dove**, **Stock Dove** and **Woodpigeon**; insectivores; and some waterbirds. Corvids appear to have benefited from the decrease of predator control by gamekeepers in recent years, and the increased use of brassica crops (particularly oilseed rape) has probably been beneficial to the larger doves.

The majority of increasing insectivores are woodland species that are also common in gardens: **Great Spotted Woodpecker**, **Green Woodpecker**, **Nuthatch**, **Blackcap**, **Great Tit**, **Wren**, **Long-tailed Tit** and **Coal Tit**. The reasons for these increases are presently unclear. **Pied Wagtail** has increased in numbers by 80% on CBC/BBS plots over 39 years, but declined by 53% on WBS plots over the past 31 years. The former survey is likely to be more representative of the UK population as a whole. The increase in **Pheasants** is driven largely by the scale of releases for shooting.

Table 4.4.1 Long-term population increases of greater than 50% from CBC/WBS (1967-2006) or WBS (1975-2006) using the best survey for each species

Species	Period (yrs)	Source	Change (%)	Lower limit	Upper limit	Alert	Comment
Buzzard	39	CBC/BBS UK	419	251	1372		
Collared Dove	34	CBC/BBS UK	415	236	646		
Great Spotted Woodpecker	39	CBC/BBS UK	349	233	603		
Shelduck	31	CBC to 1999	300	94	787		Small sample
Green Woodpecker	39	CBC/BBS England	205	133	327		
Mute Swan	39	CBC/BBS UK	199	33	473		
Nuthatch	39	CBC/BBS UK	177	98	282		
Mallard	39	CBC/BBS UK	166	111	226		
Sparrowhawk	31	CBC/BBS England	166	78	360		
Stock Dove	39	CBC/BBS England	163	79	293		
Blackcap	39	CBC/BBS UK	149	104	213		
Woodpigeon	39	CBC/BBS UK	146	22	445		
Goosander	25	WBS waterways	144	18	405		
Carrion Crow	39	CBC/BBS England	119	77	179		Includes Hooded Crow
Oystercatcher	31	WBS waterways	113	67	213		
Great Tit	39	CBC/BBS UK	110	90	142		
Magpie	39	CBC/BBS UK	101	65	152		
Pheasant	39	CBC/BBS England	99	53	187		
Wren	39	CBC/BBS UK	98	74	120		
Jackdaw	39	CBC/BBS UK	92	30	206		
Canada Goose	25	WBS waterways	90	-9	698		
Pied Wagtail	39	CBC/BBS UK	80	36	147		
Long-tailed Tit	39	CBC/BBS England	77	28	168		
Coot	31	WBS waterways	66	-16	223		
Coal Tit	39	CBC/BBS England	50	-17	188		

A number of species associated with freshwater habitats are also becoming more abundant, although differences between their ecological requirements make it unlikely that a common causal factor is involved. For **Mallard**, the CBC/BBS increase was matched by a WBS increase of 165% over 31 years. The growth of this population is still continuing, with CBC/BBS recording a 16% increase over the most recent ten-year period. The increases recorded for **Mute Swan** on both CBC/BBS and WBS plots are likely to be the result of banning the use of lead weights by anglers. **Oystercatchers** have increased by 122% on WBS plots over the last 30 years. This finding is consistent with the results of the most recent survey of **Breeding Waders of Wet Meadows** which found that numbers of Oystercatchers using these habitats in England and Wales increased by 51% between 1982 and 2002. **Grey Heron** is not listed in Table 4.4.1 because it is covered by a separate survey that spans a much longer period.

The population of this species is probably not increasing quite as fast as the species listed in the table, with only a 16% increase over the last 25 years. Nevertheless this population has undergone a sustained increase of 58% over the last 77 years (1929–2006).

Two widespread raptors have shown remarkable recoveries from low population levels caused by pesticides in the 1950s and 1960s, assisted by a relaxation of predator control. **Buzzards** increased by a remarkable 419% between 1967 and 2006, with a rapid increase of 39% over the last ten years alone. **Sparrowhawks**, too scarce for CBC to monitor until the mid 1970s, showed a 166% increase over the 31-year period from 1975 to 2006. However, their recovery appears to have been completed earlier than for **Buzzard**, with the population having been relatively stable since the early 1990s.

[Return to previous page](#)

[Next page – 4.5 Changes in breeding performance](#)

4.5 Changes in breeding performance

Changes in a range of aspects of breeding performance can be measured under the **Nest Record Scheme** (NRS) and the **Constant Effort Sites** (CES) scheme. The former provides information on components of breeding performance per nesting attempt. The latter provides an index of breeding performance accrued over all nesting attempts in a particular year, combined with the effect of changes in the survival of fledglings once they have left the nest but before they are caught as juveniles – a period when losses of young can be high.

Breeding performance may be influenced by a variety of factors, including food availability, predation pressure and weather conditions. Variation in breeding performance may help to influence a population, and may even be the main demographic factor responsible for determining its size. Conversely, the breeding performance of a population may be negatively related to its size, with productivity decreasing as the number of individuals increases, and vice versa. This relationship may be due to the action of density-dependent factors, such as competition for resources: as numbers increase, competition for resources is likely to increase, possibly resulting in poorer productivity. Alternatively, increases in abundance may be accompanied by range expansion into new, suboptimal habitats where breeding performance is poorer, thus reducing the average productivity of the population; conversely, where declines result from the loss of individuals from these suboptimal habitats, there may be a subsequent increase in average productivity.

4.5.1 Changes in clutch and brood size from Nest Record Scheme data

Those species exhibiting statistically significant trends in clutch and brood size over the past 38 years (1968–2006) are shown in Tables 4.5.1.1 and 4.5.1.2. More species showed decreases than increases in clutch size (19 decreases, 15 increases) while the reverse was true for brood size (18 decreases, 22 increases).

Table 4.5.1.1

Significant trends in Clutch size measured between 1968-2006

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Long-tailed Tit	38	34	Linear decline	7.63 eggs	6.46 eggs	-1.17 eggs	
Magpie	38	45	Curvilinear	5.62 eggs	4.56 eggs	-1.06 eggs	
Great Tit	38	152	Linear decline	8.32 eggs	7.38 eggs	-0.94 eggs	
Hen Harrier	38	13	Curvilinear	5.48 eggs	4.73 eggs	-0.75 eggs	Small sample
Moorhen	38	91	Linear decline	6.52 eggs	5.96 eggs	-0.56 eggs	
Blue Tit	38	156	Linear decline	9.28 eggs	8.74 eggs	-0.54 eggs	
Peregrine	38	16	Linear decline	3.58 eggs	3.09 eggs	-0.49 eggs	Small sample
Twite	38	12	Curvilinear	5.43 eggs	5.1 eggs	-0.33 eggs	Small sample
Mute Swan	38	22	Curvilinear	5.89 eggs	5.64 eggs	-0.25 eggs	Small sample
Pied Wagtail	38	59	Linear decline	5.12 eggs	4.92 eggs	-0.2 eggs	
Nightjar	38	17	Linear decline	2.02 eggs	1.82 eggs	-0.2 eggs	Small sample
Greenfinch	38	93	Curvilinear	4.73 eggs	4.54 eggs	-0.19 eggs	
Chaffinch	38	86	Curvilinear	4.22 eggs	4.03 eggs	-0.19 eggs	
Linnet	38	108	Curvilinear	4.7 eggs	4.57 eggs	-0.13 eggs	
Common Sandpiper	38	11	Curvilinear	3.99 eggs	3.88 eggs	-0.11 eggs	Small sample
Sedge Warbler	38	37	Curvilinear	4.95 eggs	4.85 eggs	-0.1 eggs	
Spotted Flycatcher	38	81	Curvilinear	4.22 eggs	4.15 eggs	-0.07 eggs	
Rook	38	13	Curvilinear	4.15 eggs	4.12 eggs	-0.03 eggs	Small sample
Grey Wagtail	38	38	Curvilinear	4.68 eggs	4.65 eggs	-0.03 eggs	
Carrion Crow	38	33	Curvilinear	4.08 eggs	4.09 eggs	0.01 eggs	Includes Hooded Crow
Dipper	38	72	Curvilinear	4.48 eggs	4.5 eggs	0.02 eggs	
Yellowhammer	38	44	Curvilinear	3.35 eggs	3.39 eggs	0.04 eggs	
Swallow	38	220	Curvilinear	4.46 eggs	4.51 eggs	0.05 eggs	
Lapwing	38	122	Linear increase	3.69 eggs	3.82 eggs	0.13 eggs	
Mistle Thrush	38	35	Linear increase	3.88 eggs	4.09 eggs	0.21 eggs	
Jackdaw	38	43	Linear increase	4.35 eggs	4.57 eggs	0.22 eggs	
Little Owl	38	18	Linear increase	3.39 eggs	3.62 eggs	0.23 eggs	Small sample
Dunnock	38	99	Linear increase	3.94 eggs	4.2 eggs	0.26 eggs	
Skylark	38	38	Linear increase	3.37 eggs	3.68 eggs	0.31 eggs	
Redstart	38	48	Curvilinear	5.89 eggs	6.23 eggs	0.34 eggs	
Sand Martin	38	32	Curvilinear	4.68 eggs	5.06 eggs	0.38 eggs	
Tree Sparrow	38	155	Curvilinear	4.71 eggs	5.11 eggs	0.4 eggs	
Starling	38	76	Linear increase	4.42 eggs	4.97 eggs	0.55 eggs	
Barn Owl	38	22	Linear increase	4.53 eggs	5.15 eggs	0.62 eggs	Small sample

See [Help](#) for help with interpretation

Nine species (**Nightjar**, **Pied Wagtail**, **Spotted Flycatcher**, **Long-tailed Tit**, **Blue Tit**, **Great Tit**, **Magpie**, **Chaffinch** and **Greenfinch**) exhibited decreases in both clutch size and brood size over the period, whilst another eight species (**Barn Owl**, **Skylark**, **Swallow**, **Dipper**, **Dunnock**, **Redstart**, **Tree Sparrow** and **Yellowhammer**) exhibited increases in both clutch size and brood size. **Moorhen**, **Grey Wagtail**, **Rook** and **Linnet** all showed a decline in average clutch size and an increase in average brood size, although the magnitude of the change in **Rook** clutch sizes and in **Grey Wagtail** and **Linnet**

brood sizes was small (<0.04 eggs/chicks). **Jackdaw** and **Carrion Crow** showed the opposite pattern, with clutch sizes increasing while brood sizes decreased, although again the magnitude of the change in **Carrion Crow** clutch size was very small (0.01 eggs); note that, for historical reasons, **Carrion Crow** figures include a small proportion of data from **Hooded Crow**.

Table 4.5.1.2

Significant trends in Brood size measured between 1968-2006

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Great Tit	38	287	Linear decline	7.57 chicks	6.24 chicks	-1.33 chicks	
Blue Tit	38	270	Linear decline	8.32 chicks	7.35 chicks	-0.97 chicks	
Long-tailed Tit	38	28	Curvilinear	6.68 chicks	6.17 chicks	-0.51 chicks	Small sample
Yellow Wagtail	38	13	Linear decline	4.83 chicks	4.34 chicks	-0.49 chicks	Small sample
House Sparrow	38	111	Curvilinear	3.39 chicks	2.92 chicks	-0.47 chicks	
Greenfinch	38	115	Curvilinear	4.03 chicks	3.61 chicks	-0.42 chicks	
Raven	38	67	Linear decline	3.23 chicks	2.83 chicks	-0.4 chicks	
Carrion Crow	38	79	Curvilinear	2.87 chicks	2.48 chicks	-0.39 chicks	Includes Hooded Crow
Chiffchaff	38	33	Linear decline	5.12 chicks	4.74 chicks	-0.38 chicks	
Bullfinch	38	36	Curvilinear	4.11 chicks	3.79 chicks	-0.32 chicks	
Magpie	38	79	Curvilinear	3.09 chicks	2.84 chicks	-0.25 chicks	
Pied Wagtail	38	114	Linear decline	4.53 chicks	4.34 chicks	-0.19 chicks	
Corn Bunting	38	12	Curvilinear	3.07 chicks	2.89 chicks	-0.18 chicks	Small sample
Nightjar	38	24	Curvilinear	1.82 chicks	1.7 chicks	-0.12 chicks	Small sample
Jackdaw	38	91	Curvilinear	2.69 chicks	2.6 chicks	-0.09 chicks	
Chaffinch	38	136	Curvilinear	3.57 chicks	3.49 chicks	-0.08 chicks	
Robin	38	169	Curvilinear	4.41 chicks	4.36 chicks	-0.05 chicks	
Spotted Flycatcher	38	130	Curvilinear	3.61 chicks	3.6 chicks	-0.01 chicks	
Grey Wagtail	38	80	Curvilinear	3.95 chicks	3.95 chicks	0 chicks	
Linnet	38	122	Curvilinear	4.07 chicks	4.08 chicks	0.01 chicks	
Stock Dove	38	111	Curvilinear	1.82 chicks	1.84 chicks	0.02 chicks	
Blackbird	38	151	Curvilinear	3.35 chicks	3.38 chicks	0.03 chicks	
Swallow	38	378	Curvilinear	4.07 chicks	4.12 chicks	0.05 chicks	
Yellowhammer	38	67	Curvilinear	2.96 chicks	3.03 chicks	0.07 chicks	
Buzzard	38	94	Curvilinear	1.87 chicks	1.95 chicks	0.08 chicks	
Kestrel	38	121	Curvilinear	3.74 chicks	3.83 chicks	0.09 chicks	
Collared Dove	38	70	Linear increase	1.76 chicks	1.85 chicks	0.09 chicks	
Barn Owl	38	149	Curvilinear	3.01 chicks	3.12 chicks	0.11 chicks	
Rook	38	85	Curvilinear	2.22 chicks	2.34 chicks	0.12 chicks	
Reed Warbler	38	127	Linear increase	3.45 chicks	3.58 chicks	0.13 chicks	
Duncock	38	106	Linear increase	3.48 chicks	3.63 chicks	0.15 chicks	
Skylark	38	67	Curvilinear	3.1 chicks	3.29 chicks	0.19 chicks	
Merlin	38	55	Linear increase	3.52 chicks	3.79 chicks	0.27 chicks	
Dipper	38	135	Curvilinear	3.4 chicks	3.69 chicks	0.29 chicks	
Sparrowhawk	38	71	Curvilinear	3.11 chicks	3.45 chicks	0.34 chicks	
Tree Pipit	38	28	Linear increase	4.38 chicks	4.74 chicks	0.36 chicks	Small sample
Redstart	38	85	Curvilinear	5.1 chicks	5.48 chicks	0.38 chicks	
Tree Sparrow	38	197	Curvilinear	3.78 chicks	4.21 chicks	0.43 chicks	
Nuthatch	38	62	Curvilinear	4.05 chicks	4.87 chicks	0.82 chicks	
Moorhen	38	77	Curvilinear	3.12 chicks	4.3 chicks	1.18 chicks	

See [Help](#) for help with interpretation

Long-term changes in clutch or brood size are associated with long-term population trends in a number of species. Here we highlight those changes that are both statistically significant and likely to be of biological importance.

Declines in population size and productivity were identified for **Spotted Flycatcher** (clutch and brood size), **House Sparrow** and **Bullfinch** (brood size). The mean number of eggs and chicks produced by **Spotted Flycatcher** began to fall in the mid 1980s, a worrying trend for a species that, by this point, had been in rapid decline for at least two decades. Population modelling work undertaken by [Freeman & Crick \(2003\)](#) suggests that reduced survival rates of first-year birds drove the initial population decline, but an additive effect of reduced reproductive output may now be a possibility. Declines in **Bullfinch** populations are also thought to have begun due to falling survival rates ([Proffitt et al. 2004](#), [Marquiss 2007](#)), although the mechanism is not clear-cut ([Siriwardena et al. 2001](#)) and a reduction in brood size over the last 25 years may again have had detrimental effects at a population level.

In the case of the **House Sparrow**, population modelling based on BTO data has shown that declines in rural areas were caused by reduced survival rates but that these declines were mainly halted due to improvements in breeding performance ([Crick et al. 2002](#)). The apparently accelerating reduction in brood size is therefore of some concern. [Peach et al. \(2008\)](#) suggested that insect food for the chicks may be limited in certain situations and recent brood size reductions may be a manifestation of this at a wider scale. However, it should be noted that, over the long term, some of the reduction in brood size might have been compensated for by reduced nest failure rates at the egg and chick stages.

It is worth noting that **Pied Wagtail** and **Grey Wagtail** are also exhibiting significant declines in both clutch and brood sizes – data from [WBS](#) suggest that populations using rivers and canals declined significantly between the 1970s and the 1990s, although [BBS](#) results suggest that **Pied Wagtail** populations in the wider countryside are stable.

Several increasing species exhibit concurrently increasing brood sizes, particularly **Sparrowhawk**, **Collared Dove**, **Redstart** and **Nuthatch**. **Sparrowhawk** has returned to eastern areas of the UK, where populations of songbird prey are greater, which may have impacted positively on breeding success, although both the population and the brood size trends began to level out in the early to mid 1990s. **Collared Dove** has spread rapidly since colonising the UK in the 1950s, and brood sizes have

exhibited a steady increase over the past 35 years. **Redstart** is one of the only long-distance migrant passerines currently exhibiting significant population growth, and increasing clutch and brood sizes over the last three decades may have contributed to this. The UK **Nuthatch** population, which has been expanding northwards and has increased considerably in size since the 1970s, currently produces 0.75 more young per nesting attempt than it did on average during the late 1960s. It would seem likely that this has helped to drive the population increase of this species; brood sizes have started to fall in the last decade, however, possibly indicating the onset of density dependent-reductions in productivity (see below).

Inverse associations between clutch or brood size and population trend are found in some 23 species (i.e. they show lower productivity with higher population size). Such relationships may arise through density-dependent processes, whereby increased competition leads to reduced clutch or brood sizes at higher population densities. Eleven increasing species and 12 decreasing species show such associations. Notable examples amongst increasing species include **Magpie**, **Blue Tit**, **Great Tit**, **Long-tailed Tit**, **Chaffinch**, **Greenfinch** (clutch and brood size) **Robin** and **Chiffchaff** (brood size). Amongst declining species examples include **Skylark**, **Duncock**, **Tree Sparrow** (clutch and brood size), **Lapwing**, **Starling** and **Mistle Thrush** (clutch size).

4.5.2 Changes in nest failure rates from Nest Record Scheme data

Statistically significant trends in the daily nest failure rates at the egg and chick stages over the past 38 years (1968–2006) are shown in Tables 4.5.2.1 and 4.5.2.2. The number of species exhibiting declines in failure rates at the egg stage (39) was treble the number displaying increases (13), and while 26 species exhibited declines in chick-stage failure rates, only 13 displayed increases. Thus, the general picture is one of improving nesting success.

Table 4.5.2.1

Significant trends in Daily failure rate (eggs) measured between 1968-2006

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Woodlark	38	20	Curvilinear	0.0683 nests/day	0.0275 nests/day	-0.0408 nests/day	Small sample
Long-tailed Tit	38	52	Linear decline	0.0358 nests/day	0.0084 nests/day	-0.0274 nests/day	
Magpie	38	52	Linear decline	0.0273 nests/day	0.0028 nests/day	-0.0245 nests/day	
Redshank	38	33	Linear decline	0.0398 nests/day	0.0169 nests/day	-0.0229 nests/day	
Dipper	38	100	Curvilinear	0.0257 nests/day	0.0033 nests/day	-0.0224 nests/day	
Snipe	38	16	Linear decline	0.033 nests/day	0.0136 nests/day	-0.0194 nests/day	Small sample
Tree Pipit	38	12	Curvilinear	0.0487 nests/day	0.0312 nests/day	-0.0175 nests/day	Small sample
Robin	38	188	Curvilinear	0.0247 nests/day	0.0103 nests/day	-0.0144 nests/day	
Carrion Crow	38	50	Linear decline	0.0159 nests/day	0.0022 nests/day	-0.0137 nests/day	Includes Hooded Crow
Treecreeper	38	22	Linear decline	0.0191 nests/day	0.0062 nests/day	-0.0129 nests/day	Small sample
Sand Martin	38	24	Linear decline	0.0127 nests/day	0.0001 nests/day	-0.0126 nests/day	Small sample
Yellowhammer	38	66	Curvilinear	0.05 nests/day	0.038 nests/day	-0.012 nests/day	
Wood Warbler	38	21	Linear decline	0.0191 nests/day	0.0079 nests/day	-0.0112 nests/day	Small sample
Redstart	38	73	Linear decline	0.0116 nests/day	0.0031 nests/day	-0.0085 nests/day	
Stock Dove	38	78	Curvilinear	0.012 nests/day	0.0036 nests/day	-0.0084 nests/day	
Pied Wagtail	38	83	Curvilinear	0.0152 nests/day	0.0068 nests/day	-0.0084 nests/day	
Starling	38	118	Linear decline	0.0112 nests/day	0.003 nests/day	-0.0082 nests/day	
Tawny Owl	38	55	Linear decline	0.0095 nests/day	0.0018 nests/day	-0.0077 nests/day	Nocturnal species
House Sparrow	38	92	Linear decline	0.0115 nests/day	0.0041 nests/day	-0.0074 nests/day	
Wheatear	38	18	Curvilinear	0.0082 nests/day	0.0009 nests/day	-0.0073 nests/day	Small sample
Sedge Warbler	38	44	Linear decline	0.013 nests/day	0.0061 nests/day	-0.0069 nests/day	
Greenfinch	38	132	Linear decline	0.0248 nests/day	0.018 nests/day	-0.0068 nests/day	
Barn Owl	38	18	Linear decline	0.0075 nests/day	0.001 nests/day	-0.0065 nests/day	Small sample
Marsh Tit	38	20	Linear decline	0.0076 nests/day	0.0014 nests/day	-0.0062 nests/day	Small sample
Buzzard	38	26	Linear decline	0.007 nests/day	0.0012 nests/day	-0.0058 nests/day	Small sample
Jackdaw	38	54	Linear decline	0.0076 nests/day	0.002 nests/day	-0.0056 nests/day	
Kestrel	38	40	Linear decline	0.0055 nests/day	0.0009 nests/day	-0.0046 nests/day	
Merlin	38	26	Linear decline	0.0067 nests/day	0.0024 nests/day	-0.0043 nests/day	Small sample
Tree Sparrow	38	207	Curvilinear	0.0074 nests/day	0.0032 nests/day	-0.0042 nests/day	
Wren	38	141	Linear decline	0.0183 nests/day	0.0143 nests/day	-0.004 nests/day	
Sparrowhawk	38	34	Linear decline	0.0047 nests/day	0.0009 nests/day	-0.0038 nests/day	
Stonechat	38	35	Curvilinear	0.0052 nests/day	0.0027 nests/day	-0.0025 nests/day	
Great Tit	38	265	Curvilinear	0.0068 nests/day	0.0043 nests/day	-0.0025 nests/day	
Blue Tit	38	259	Linear decline	0.0046 nests/day	0.0025 nests/day	-0.0021 nests/day	
Raven	38	22	Curvilinear	0.0021 nests/day	0.0009 nests/day	-0.0012 nests/day	Small sample
Duncock	38	143	Curvilinear	0.0251 nests/day	0.0248 nests/day	-0.0003 nests/day	
Spotted Flycatcher	38	122	Curvilinear	0.0177 nests/day	0.0188 nests/day	0.0011 nests/day	
Peregrine	38	21	Curvilinear	0.0014 nests/day	0.0028 nests/day	0.0014 nests/day	Small sample
Hen Harrier	38	11	Curvilinear	0.0003 nests/day	0.0022 nests/day	0.0019 nests/day	Small sample
Ringed Plover	38	125	Linear increase	0.0236 nests/day	0.0285 nests/day	0.0049 nests/day	
Linnet	38	154	Curvilinear	0.0165 nests/day	0.0229 nests/day	0.0064 nests/day	
Chaffinch	38	165	Curvilinear	0.0297 nests/day	0.0368 nests/day	0.0071 nests/day	
Willow Warbler	38	69	Linear increase	0.0093 nests/day	0.0166 nests/day	0.0073 nests/day	
Lapwing	38	133	Curvilinear	0.0167 nests/day	0.0245 nests/day	0.0078 nests/day	
Moorhen	38	110	Curvilinear	0.0135 nests/day	0.0215 nests/day	0.008 nests/day	
Mute Swan	38	30	Curvilinear	0.0061 nests/day	0.0141 nests/day	0.008 nests/day	Small sample
Bullfinch	38	50	Curvilinear	0.0333 nests/day	0.0413 nests/day	0.008 nests/day	
Blackbird	38	179	Curvilinear	0.0257 nests/day	0.0365 nests/day	0.0108 nests/day	
Oystercatcher	38	112	Curvilinear	0.0143 nests/day	0.0272 nests/day	0.0129 nests/day	
Whinchat	38	15	Linear increase	0.0065 nests/day	0.0217 nests/day	0.0152 nests/day	Small sample
Reed Bunting	38	52	Linear increase	0.0073 nests/day	0.0274 nests/day	0.0201 nests/day	
Nightjar	38	22	Linear increase	0.0138 nests/day	0.0355 nests/day	0.0217 nests/day	Small sample

Increases in both egg-stage and chick-stage failure rates were observed for four species: **Nightjar**, **Spotted Flycatcher**, **Linnet** and **Bullfinch**. Use of nest cameras to identify potential nest predators of **Nightjar** nests is currently under way in Theford Forest as part of a collaborative project between BTO and the University of East Anglia, which is also monitoring **Woodlark**. Researchers have also used nest cameras to observe predation events at **Spotted Flycatcher** nests, identifying **Jay** as the main predator in their study population in SW England ([Stevens et al. 2008](#)). Predation is not necessarily the only cause of nest failure, but it is a major cause for open-nesting bird species.

Eighteen species exhibited declines in both egg-stage and chick-stage failure rates: **Kestrel**, **Merlin**, **Barn Owl**, **Tawny Owl**, **Stock Dove**, **Sand Martin**, **Pied Wagtail**, **Robin**, **Redstart**, **Stonechat**, **Magpie**, **Jackdaw**, **Carrion Crow**, **Raven**, **Starling**, **House Sparrow**, **Tree Sparrow** and **Yellowhammer**. For a further six species (**Tree Pipit**, **Dipper**, **Whinchat**, **Blackbird**, **Great Tit**, **Blue Tit** and **Long-tailed Tit**), better success at one stage was partly cancelled out by increases in failure rates at the other, suggesting that different factors may influence productivity at egg and chick stages.

Table 4.5.2.2

Significant trends in Daily failure rate (chicks) measured between 1968-2006

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Grey Heron	38	26	Linear decline	0.054 nests/day	0.0004 nests/day	-0.0536 nests/day	Non-breeders included
Corn Bunting	38	11	Curvilinear	0.045 nests/day	0.0224 nests/day	-0.0226 nests/day	Small sample
Meadow Pipit	38	60	Linear decline	0.0271 nests/day	0.0108 nests/day	-0.0163 nests/day	
Magpie	38	50	Linear decline	0.0173 nests/day	0.0013 nests/day	-0.016 nests/day	
Sand Martin	38	36	Linear decline	0.0157 nests/day	0.0008 nests/day	-0.0149 nests/day	
Reed Warbler	38	110	Curvilinear	0.0174 nests/day	0.005 nests/day	-0.0124 nests/day	
Blackbird	38	154	Linear decline	0.0303 nests/day	0.0192 nests/day	-0.0111 nests/day	
Grey Wagtail	38	58	Linear decline	0.0206 nests/day	0.0098 nests/day	-0.0108 nests/day	
Jackdaw	38	52	Linear decline	0.0126 nests/day	0.0024 nests/day	-0.0102 nests/day	
House Sparrow	38	88	Linear decline	0.013 nests/day	0.0036 nests/day	-0.0094 nests/day	
Redstart	38	52	Linear decline	0.0123 nests/day	0.0039 nests/day	-0.0084 nests/day	
Collared Dove	38	55	Linear decline	0.0182 nests/day	0.0106 nests/day	-0.0076 nests/day	
Stonechat	38	54	Curvilinear	0.0116 nests/day	0.004 nests/day	-0.0076 nests/day	
Robin	38	162	Curvilinear	0.0249 nests/day	0.0177 nests/day	-0.0072 nests/day	
Merlin	38	29	Linear decline	0.0094 nests/day	0.0024 nests/day	-0.007 nests/day	Small sample
Tree Sparrow	38	151	Curvilinear	0.0125 nests/day	0.0064 nests/day	-0.0061 nests/day	
Carrion Crow	38	42	Linear decline	0.0074 nests/day	0.0014 nests/day	-0.006 nests/day	Includes Hooded Crow
Stock Dove	38	57	Linear decline	0.012 nests/day	0.0071 nests/day	-0.0049 nests/day	
Starling	38	136	Linear decline	0.0063 nests/day	0.0019 nests/day	-0.0044 nests/day	
Pied Wagtail	38	91	Linear decline	0.0127 nests/day	0.0084 nests/day	-0.0043 nests/day	
Yellowhammer	38	52	Curvilinear	0.0462 nests/day	0.0423 nests/day	-0.0039 nests/day	
Great Spotted Woodpecker	38	31	Linear decline	0.0037 nests/day	0.0002 nests/day	-0.0035 nests/day	
Tawny Owl	38	82	Curvilinear	0.0031 nests/day	0.0011 nests/day	-0.002 nests/day	Nocturnal species
Barn Owl	38	69	Linear decline	0.0021 nests/day	0.0003 nests/day	-0.0018 nests/day	
Kestrel	38	65	Linear decline	0.0022 nests/day	0.0009 nests/day	-0.0013 nests/day	
Raven	38	29	Curvilinear	0.0003 nests/day	0.0002 nests/day	-0.0001 nests/day	Small sample
Whinchat	38	26	Curvilinear	0.0261 nests/day	0.0261 nests/day	0 nests/day	Small sample
Twite	38	13	Curvilinear	0.0061 nests/day	0.0062 nests/day	0.0001 nests/day	Small sample
Dipper	38	78	Curvilinear	0.0059 nests/day	0.0062 nests/day	0.0003 nests/day	
Blue Tit	38	194	Linear increase	0.0065 nests/day	0.0087 nests/day	0.0022 nests/day	
Swallow	38	251	Linear increase	0.0028 nests/day	0.0053 nests/day	0.0025 nests/day	
Great Tit	38	197	Linear increase	0.0057 nests/day	0.0083 nests/day	0.0026 nests/day	
Spotted Flycatcher	38	109	Linear increase	0.0098 nests/day	0.0143 nests/day	0.0045 nests/day	
Nightjar	38	21	Curvilinear	0.0005 nests/day	0.0067 nests/day	0.0062 nests/day	Small sample
Bullfinch	38	34	Curvilinear	0.0336 nests/day	0.0405 nests/day	0.0069 nests/day	
Linnet	38	110	Linear increase	0.0152 nests/day	0.0224 nests/day	0.0072 nests/day	
Tree Pipit	38	19	Curvilinear	0.0337 nests/day	0.0411 nests/day	0.0074 nests/day	Small sample
Long-tailed Tit	38	37	Linear increase	0.008 nests/day	0.0171 nests/day	0.0091 nests/day	

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
---------	--------------	--------------------	-------	-------------------------	------------------------	--------	---------

See [Help](#) for help with interpretation

Long-term changes in egg-stage or chick-stage nest failure rates are associated with long-term population trends in a number of species. Here we highlight those changes that are both statistically significant and likely to be of biological importance.

Increased nest failure rates were associated with long-term decreases in population size for ten species, and may have contributed to the observed population declines of **Nightjar**, **Spotted Flycatcher**, **Linnet**, **Bullfinch** (egg- and chick-stage failure rates increasing), **Lapwing**, **Willow Warbler** and **Reed Bunting** (egg-stage failure rates increasing). Although **Nightjar** has shown a large historical decline and is red-listed because of this, it should be noted that recent surveys show a population increase ([Conway et al. 2007](#)). Reductions in breeding performance at the egg stage have been implicated in a detailed analysis of the population declines of the **Linnet** ([Siriwardena et al. 2000b](#)), but the extent to which decreased productivity has influenced Bullfinch population trends is still not clearly understood ([Siriwardena et al. 2001](#)). It has also been suggested that poor breeding performance may be preventing the recovery of **Reed Bunting** populations ([Peach et al. 1999](#)). The increasing trend in egg- and chick-stage failure rates of **Spotted Flycatcher** has only recently become significant and previous work suggested that other demographic factors were more important in the decline of this species ([Freeman & Crick 2003](#)). Researchers have failed to find any trend in survival rates that might explain declines in **Lapwing** numbers ([Peach et al. 1994](#), [King et al. 2008](#)), and a fall in productivity is thought to have been a major factor ([Galbraith 1988](#)), although a recent study failed to find any correlation between nest failure rates and changes in abundance at a regional scale ([Sharpe et al. 2008](#)). The population decline of Willow Warbler is much more pronounced in the south of the UK than in Scotland, and collaborative project between BTO and the University of East Anglia is currently investigating whether demographic parameters, including productivity, exhibit the similar patterns of spatial variation. Failure rates are also thought to be increasing at the egg and chick stages for the declining **Hen Harrier** and **Twite**, although sample sizes for both are extremely limited, and at the egg stage for **Moorhen**, a species for which the current population trajectory is unclear but which did decline in abundance during the 1980s.

Sixteen species showed clear associations between long-term increases in abundance and long-term reductions in nest failure rates. **Sparrowhawk**, **Buzzard**, **Wren** and **Greenfinch** experienced reduced nest failure rates at the egg stage, while **Grey Heron**, **Great Spotted Woodpecker** and **Collared Dove** exhibited a reduction in failure rates at the chick stage. The remaining nine species (**Barn Owl**, **Stock Dove**, **Robin**, **Redstart**, **Stonechat**, **Maggie**, **Jackdaw**, **Corrion Crow** and **Raven**) displayed reduced failure rates at both the egg and chick stages. Corvids, such as **Maggie**, **Jackdaw**, **Corrion Crow** and **Raven**, appear to have benefited from improvements in nesting success at the egg stage, as have owls and raptors such as **Barn Owl**, **Sparrowhawk** and **Buzzard**. Decreased persecution and reduction in the use of pesticides are likely to have been important factors in the recovery of these species. The improvements in the nesting success of **Stock Dove** could have had a major impact on the size of the population, given the high number of breeding attempts made by this species each year, and the decreased chick-stage failure rates of **Collared Dove** may have aided the rapid growth of the UK population over the last 38 years. **Grey Heron** populations have increased over the last 70 years, and improvements in chick-stage nest survival may have played a part in this increase. **Greenfinch** has rapidly adapted to the provision of supplementary food in gardens and is now very much associated with human habitats. Such provisioning may lead to increased productivity because of its positive impact on adult body condition. Causes of the reduction in failure rates for **Wren**, **Robin**, **Redstart** and **Stonechat** are less clear, although all feed primarily on ground-dwelling invertebrates and changes in arthropod abundance or activity could be responsible.

Inverse associations between changes in egg- or chick-stage nest survival and population trends are found in some 23 species. Such relationships may arise through density-dependent processes where increased competition leads to increased failure rates at higher population densities. Four increasing species (**Mute Swan**, **Peregrine**, **Oystercatcher** and **Chaffinch**) showed long-term increases in egg-stage failure rates, while 19 declining species showed evidence of improving nesting success. **Ringed Plover**, **Snipe**, **Redshank**, **Woodlark**, **Duncock**, **Wheatear**, **Sedge Warbler**, **Wood Warbler** and **Marsh Tit** showed decreasing failure at the egg stage while decreasing chick-stage failure rates were identified for **Meadow Pipit**, **Grey Wagtail** and **Corn Bunting**. The remaining seven species, **Kestrel**, **Merlin**, **Starling**, **House Sparrow**, **Tree Sparrow**, **Yellowhammer** and **Pied Wagtail** (which is thought to be declining in waterway habitats), exhibited decreasing failure rates at both stages.

Several species demonstrated a decrease in failure rates at one stage but a compensatory increase at the other, including **Dipper**, **Great Tit**, **Blue Tit**, **Long-tailed Tit**, **Tree Pipit**, **Whinchat** (declining egg failure, increasing chick failure) and **Blackbird** (increasing egg-stage failure, declining chick-stage failure).

4.5.3 Changes in productivity from Constant Effort Scheme ringing data

The CES started monitoring populations in 1983, so the changes in productivity shown in Table 4.5.3 cover roughly half the time period of the Nest Record Scheme results. The CES data set is unique in providing relative measures of adult abundance and productivity from the same set of sites in wetland and scrub habitats. While the NRS data set monitors the productivity of individual nesting attempts, the proportion of juveniles in the CES catch provides a relative measure of annual variation in productivity that integrates the effects of the number of fledglings produced per attempt, number of nesting attempts and immediate post-fledging survival. Use of these two techniques in combination provides a powerful method of determining which factors are responsible for observed declines in recruitment of young birds into the breeding population.

Overall, ten species exhibit declines of greater than 25% in the proportion of juveniles captured, while only **Chaffinch** shows an increase of greater than 25% in the ratio of juveniles to adults. Six of these

species, **Nightingale**, **Sedge Warbler**, **Blue Tit**, **Linnet**, **Goldfinch** and **Reed Bunting**, all exhibit declines in the proportion of juveniles captured over the last 20 years of greater than 50%, although it should be noted that **Nightingale** occurs on a relatively small number of plots. A further four species show reductions in relative productivity of between 25% and 50%: **Song Thrush**, **Blackcap**, **Willow Warbler** and **Great Tit**. Of the nine of these species for which sufficient Nest Record Scheme data are available for comparison (**Nightingale** is excluded), six have also been identified as exhibiting negative trends in either clutch size, brood size or nest survival (**Sedge Warbler**, **Willow Warbler**, **Blue Tit**, **Great Tit**, **Linnet** and **Reed Bunting**).

Six of the ten species exhibiting productivity declines greater than 25% (**Nightingale**, **Song Thrush**, **Sedge Warbler**, **Willow Warbler**, **Linnet** and **Reed Bunting**) have experienced significant population declines, either on CES sites or more widely (based on CBC/BBS figures). For **Linnet** there is good evidence that variation in productivity has been important in driving the decline ([Siriwardena et al. 2000b](#)), but for **Song Thrush**, **Willow Warbler** and **Reed Bunting** other work indicates that variation in survival rates is likely to have been a more important contributor to population changes ([Peach et al. 1995a](#), [Peach et al. 1999](#), [Robinson et al. 2004](#), [Baillie et al. 2008](#)). The large decline in **Nightingale** productivity may have contributed to the complex changes in its distribution shown by the **1999 survey**, which identified decreases in abundance over large parts of the species' range. The four other species (**Blackcap**, **Great Tit**, **Blue Tit**, and **Goldfinch**) demonstrating marked reductions in productivity on CES sites have not experienced related declines in abundance, either on CES sites or more widely. These productivity declines may be driven by density-dependent processes, whereby increased competition for resources in an expanding population reduces the mean breeding success per pair.

Taking the CES data set as a whole, 20 species show some decline in productivity over the last 22 years while only five show increases. The strong preponderance of trends towards lower productivity requires urgent and more detailed investigation.

Table 4.5.3

Changes in productivity indices (percentage juveniles) for CES 1984-2006 (22 years) calculated from smoothed trend

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Comment
Nightingale	22	11	-85	-99	-27	Small sample
Linnet	22	22	-79	-94	-39	
Goldfinch	22	38	-63	-86	-14	
Sedge Warbler	22	69	-55	-75	-34	
Blue Tit	22	99	-54	-64	-41	
Reed Bunting	22	62	-54	-75	-18	
Song Thrush	22	88	-37	-55	-14	
Great Tit	22	97	-32	-51	-1	
Willow Warbler	22	95	-31	-49	-9	
Blackcap	22	94	-25	-44	-5	
Chaffinch	22	84	61	3	192	

See [Help](#) for help with interpretation

4.5.4 Changes in average laying dates from Nest Record Scheme data

Over the past 25 years, many species have exhibited a trend towards progressively earlier clutch initiation ([Crick et al. 1997](#)) with laying dates showing curvilinear responses over the past 50 years as spring temperatures have cooled and then warmed ([Crick & Sparks 1999](#)). Table 4.5.4 confirms that over the past 38 years the majority of species exhibiting significant trends show an advancement of laying dates rather than a delay. Thus 40 species are laying between 31 days and 1 day earlier, on average, than they were 38 years ago. Three species, **Nightjar**, **Twite** and **Wood Warbler**, are added to the list of earlier layers published in the previous report in this series. There are no taxonomic or ecological associations between the species showing such changes, and they seem to occur across a wide range of species ([Crick et al. 1997](#)).

Only two species, **Skylark** and **Yellowhammer**, show significant changes towards later laying, and sample sizes are small. Both of these species are multi-brooded, however, and it may be that differences in the ratio of first to repeat broods initiated may be obscuring advances in laying date. Evidence of temporal and spatial variation in **Yellowhammer** repeat-brooding over time is currently being investigated in a collaborative research project between BTO and Aberdeen University. It is likely that the laying dates of the majority of those species that do not show a significant trend in timing of laying are related to some aspect of weather, but that those aspects do not show any trend over time ([Crick & Sparks 1999](#)).

The significance of the changes in phenology for breeding performance not well understood but is an active research area within several research groups. Earlier average laying may be beneficial for birds because earlier fledging is often related to improved survival to the following year – early-nesting parents have an increased chance of having their offspring recruited into the next generation ([Visser et al. 1998](#)). However, the timing of leaf emergence and the speed of caterpillar development is also changing under increased temperatures ([Buse et al. 1999](#), [Visser & Holleman 2001](#)) and the results of several recent studies have suggested that some birds may be unable to advance their phenology sufficiently to match phenological changes in their food supply, such that later-nesting birds are suffering from poorer productivity. [Both et al. \(2006\)](#) demonstrated that mismatches between periods of food availability and chick demand can affect abundance in Dutch **Pied Flycatcher** populations, with those demonstrating the largest mismatches between arrival in spring and peak caterpillar abundance exhibiting the greatest

declines. As a consequence of climate change there may be an increasing mismatch between predator activities and the availability of their food supplies at different trophic levels within ecosystems ([Both et al. 2009](#)). The conservation significance of such phenological disjunction remains an active research area with potentially important policy implications for conservation.

Table 4.5.4

Significant trends in Laying date measured between 1968-2006

Species	Period (yrs)	Mean annual sample	Trend	Predicted in first year	Predicted in last year	Change	Comment
Magpie	38	35	Curvilinear	Apr 21	Mar 21	-31 days	
Grey Heron	38	26	Linear decline	Apr 8	Mar 11	-28 days	Non-breeders included
Long-tailed Tit	38	44	Linear decline	Apr 21	Apr 6	-15 days	
Greenfinch	38	95	Linear decline	May 25	May 10	-15 days	
Chiffchaff	38	44	Linear decline	May 17	May 3	-14 days	
Redstart	38	61	Curvilinear	May 21	May 9	-12 days	
Nuthatch	38	26	Linear decline	May 2	Apr 21	-11 days	Small sample
Carrion Crow	38	32	Curvilinear	Apr 16	Apr 5	-11 days	Includes Hooded Crow
Tree Pipit	38	19	Linear decline	May 25	May 16	-9 days	Small sample
Dipper	38	59	Linear decline	Apr 18	Apr 9	-9 days	
Blackcap	38	37	Curvilinear	May 20	May 11	-9 days	
Marsh Tit	38	14	Linear decline	Apr 28	Apr 19	-9 days	Small sample
Treecreeper	38	13	Linear decline	May 7	Apr 28	-9 days	Small sample
Swallow	38	105	Curvilinear	Jun 20	Jun 12	-8 days	
Garden Warbler	38	21	Curvilinear	May 25	May 17	-8 days	Small sample
Chaffinch	38	107	Linear decline	May 11	May 3	-8 days	
Oystercatcher	38	46	Linear decline	May 17	May 10	-7 days	
Tawny Owl	38	14	Linear decline	Mar 29	Mar 22	-7 days	Nocturnal species
Whitethroat	38	18	Curvilinear	May 26	May 19	-7 days	Small sample
Blue Tit	38	197	Linear decline	May 3	Apr 26	-7 days	
Great Tit	38	176	Linear decline	May 4	Apr 27	-7 days	
House Sparrow	38	51	Linear decline	May 25	May 18	-7 days	
Goldfinch	38	22	Linear decline	Jun 6	May 30	-7 days	Small sample
Lesser Redpoll	38	11	Curvilinear	May 26	May 19	-7 days	Small sample
Kestrel	38	21	Linear decline	May 4	Apr 28	-6 days	Small sample
Robin	38	120	Linear decline	Apr 28	Apr 22	-6 days	
Stonechat	38	35	Curvilinear	May 3	Apr 27	-6 days	
Ring Ouzel	38	24	Linear decline	May 14	May 8	-6 days	Small sample
Sedge Warbler	38	49	Curvilinear	May 29	May 23	-6 days	
Reed Warbler	38	159	Curvilinear	Jun 16	Jun 10	-6 days	
Willow Warbler	38	84	Linear decline	May 20	May 14	-6 days	
Moorhen	38	68	Linear decline	May 10	May 5	-5 days	
Wren	38	87	Linear decline	May 14	May 9	-5 days	
Whinchat	38	27	Curvilinear	May 28	May 23	-5 days	Small sample
Wood Warbler	38	31	Curvilinear	May 23	May 18	-5 days	
Jackdaw	38	23	Curvilinear	Apr 23	Apr 18	-5 days	Small sample
Tree Sparrow	38	167	Linear decline	May 29	May 24	-5 days	
Twite	38	16	Linear decline	Jun 2	May 28	-5 days	Small sample
Dunnock	38	79	Linear decline	May 3	Apr 29	-4 days	
Nightjar	38	18	Curvilinear	Jun 17	Jun 16	-1 days	Small sample
Skylark	38	20	Curvilinear	May 25	Jun 1	7 days	Small sample
Yellowhammer	38	27	Linear increase	May 30	Jun 7	8 days	Small sample

See [Help](#) for help with interpretation

[Return to previous page](#)

[Next page – 4.6 Conclusion](#)

5. Acknowledgements

Volunteer fieldwork

Our biggest thankyou is to the volunteers who collected the data on which this website is based. The population trends and other results that we present rely on the sustained, long-term fieldwork of many thousands of BTO volunteers. Our knowledge of the conservation status of the UK's bird populations is possible only as a result of their dedication and enthusiasm. The conservation community owes them an enormous debt of gratitude for their work. We are also very grateful to the many farmers, land managers and landowners who permitted census work, nest recording and ringing to take place on their land.

Report production and analysis

This website presents the latest in a series of reports, prepared within the partnership between the British Trust for Ornithology (BTO) and the Joint Nature Conservation Committee (JNCC) (on behalf of Natural England, Scottish Natural Heritage, the Countryside Council for Wales and the Northern Ireland Environment Agency), as part of its programme of research into nature conservation.

Mr and Mrs J A Pye's Charitable Settlement provided additional support towards the development of the website.

This report includes results from the Breeding Bird Survey, which is funded jointly by BTO, JNCC and RSPB. The BBS partners are very grateful to the Environment and Heritage Service in Northern Ireland (now Northern Ireland Environment Agency) and to the Royal Society for the Protection of Birds in Scotland for supporting professional surveys in areas that would otherwise be difficult to cover.

Helen Baker and Ian McLean of JNCC provided helpful discussions, comments and support during the production of the report. David Stroud, Rowena Langston, David Gibbons, Jacquie Clark, Nigel Clark, Jeremy Greenwood and Malcolm Vincent provided helpful comments on earlier editions of this publication.

The analyses would not have been possible without the hard work of many past and present BTO staff who have organised schemes, collated data sets or overseen analyses, including: Sue Adams, Dawn Balmer, Jeremy Blackburn, Jacquie Clark, Mark Collier, Rachel Coombes, Humphrey Crick, Steve Freeman, Bridget Griffin, Mike Raven, Brenda Read, Richard Thewlis, Anne Trewhitt and Jane Waters. The work is also heavily dependent on the BTO's computer and database systems operated by Peter Lack and Karen Wright. Iain Downie and Susan Waghorn exercised great skill and effort in designing and building the website, and in creating the programs that produce the figures and tables. The site is now maintained by Mandy Andrews.

We are very grateful to all of the above organisations and individuals for their contributions to this report.

[Section 6 – References](#)

[Back to Contents](#)



-
-
-
-







FR Registered with
**FUNDRAISING
REGULATOR**





REFERENCES

Clicking on the short form of any reference in the text of this report will bring you to its full details in this section: the reference sought will be at the very top of your view. Where possible, we provide an onward link either to an abstract or, where it is freely available, to the full text. Alternatively, your own web search will often take you to the summary of an article and the opportunity to purchase the text in full.

Most of the listed publications are available in printed form to BTO members and other bona fide researchers through the [Chris Mead Library](#) at BTO headquarters in Thetford.

Aebischer, N.J. (1999) Multi-way comparisons and generalised linear models of nest success: extensions of the Mayfield method. *Bird Study* 46: S22–S31.

Aebischer, N.J. & Ewald, J.A. (2004) Managing the UK Grey Partridge *Perdix perdix* recovery: population change, reproduction, habitat and shooting. *Ibis* 146 (S2): 181–191.

Aebischer, N.J., Evans, A.D., Grice, P.V. & Vickery, J.A. (2000) *The Ecology and Conservation of Lowland Farmland Birds*. British Ornithologists' Union, Tring.

Amar, A. & Redpath, S.M. (2005) Habitat use by Hen Harriers *Circus cyaneus* on Orkney: implications of land-use change for this declining population. *Ibis* 147: 37–47.

Amar, A., Picozzi, N., Meek, E.R., Redpath, S.M. & Lambin, X. (2005) Decline of the Orkney Hen Harrier *Circus cyaneus* population: do changes to demographic parameters and mating system fit a declining food hypothesis? *Bird Study* 52: 18–24.

Amar, A., Arroyo, B., Meek, E., Redpath, S. & Riley, H. (2008) Influence of habitat on breeding performance of Hen Harriers *Circus cyaneus* in Orkney. *Ibis* 150: 400–404.

Anganuzzi, A.A. (1993) A comparison of tests for detecting trends in abundance indices of dolphins. *Fishery Bulletin* 91: 183–194.

Anonymous (1995) *Biodiversity: the UK Steering Group report. Vol. 1: Meeting the Rio Challenge. Vol. 2: Action Plans*. HMSO, London.

Anonymous (1998) *UK Biodiversity Group Tranche 2 Action Plans. Vol. 1: Vertebrates & Vascular Plants*. English Nature, Peterborough.

Austin, G.E., Rehfisch, M.M., Allan, J.R. & Holloway, S.J. (2007) Population size and differential population growth of introduced Greater Canada Geese *Branta canadensis* and re-established Greylag Geese *Anser anser* across habitats in Great Britain in the year 2000. *Bird Study* 54: 343–352. [Abstract](#)

Austin, G.E., Collier, M.P., Calbrade, N.A., Hall, C. & Musgrove, A.J. (2008) *Waterbirds in the UK 2006/07: the Wetland Bird Survey*. BTO/WWT/RSPB/JNCC, Thetford. [Full text](#)

Baillie, S.R. (1990) Integrated population monitoring of breeding birds in Britain and Ireland. *Ibis* 132: 151–166.

Baillie, S.R. (1991) Monitoring terrestrial breeding bird populations. In Goldsmith, F.B. (ed.) *Monitoring for Conservation and Ecology*: 112–132. Chapman & Hall, London.

Baillie, S.R. & Peach, W.J. (1992) Population limitation in Palaearctic–African migrant passerines. *Ibis* 134 Suppl. 1: 120–132.

Baillie, S.R. & Rehfisch, M.M. (eds) (2006) *National and site-based alert systems for UK birds*. Research Report 226. BTO, Thetford. [Abstract/Full text](#)

Baillie, S.R., Crick, H.Q.P., Balmer, D.E., Beaven, L.P., Downie, I.S., Freeman, S.N., Leech, D.I., Marchant, J.H., Noble, D.G., Raven, M.J., Simpkin, A.P., Thewlis, R.M. & Wernham, C.V. (2002) *Breeding Birds in the Wider Countryside: their conservation status 2001*. Research Report 278. BTO, Thetford. (www.bto.org/birdtrends2001)

Baillie, S.R., Brooks, S.P., King, R. & Thomas, L. (2008) Using a state–space model of the British Song Thrush *Turdus philomelos* population to diagnose the causes of a population decline. Pp 541–561 in: Thomson, D.L., Cooch, E.G. & Conroy, M.J. (eds) *Modeling Demographic Processes in Marked Populations*. Series: Environmental and Ecological Statistics, Vol 3. Springer. ISBN 978-0-387-78150-1.

Baker, H., Stroud, D.A., Aebischer, N.J., Cranswick, P.A., Gregory, R.D., McSorley, C.A., Noble, D.G. & Rehfisch, M.M. (2006) Population estimates of birds in Great Britain and the United Kingdom. *British Birds* 99: 25–44. (APEP06) [Full text](#)

Balmer, D.E., Adams, S.Y. & Crick H.Q.P. (2000) Report on Barn Owl Release Scheme: Monitoring Project Phase II. Research Report 250. BTO, Thetford. [Abstract](#)

- Banks, A.N., Coombes, R.H. & Crick, H.Q.P. (2003) *The Peregrine Falcon breeding population of the UK & Isle of Man in 2002*. Research Report 330. BTO, Thetford. [Abstract](#)
- Beale, C.M., Burfield, I.J., Sim, I.M.W., Rebecca, G.W., Pearce-Higgins, J.W. & Grant, M.C. (2006) Climate change may account for the decline in British ring ouzels *Turdus torquatus*. *Journal of Animal Ecology* 75: 826–835.
- Besbeas, P., Freeman, S.N., Morgan, B.J.T. & Catchpole, E.A. (2002) Integrating mark–recapture–recovery and census data to estimate animal abundance and demographic parameters. *Biometrics* 58: 540–547.
- Bibby, C.J. (1989) A survey of breeding Wood Warblers *Phylloscopus sibilatrix*, in Britain 1984–85. *Bird Study* 36: 56–72.
- Bibby, C.J. & Etheridge, B. (1993) Status of the Hen Harrier *Circus cyaneus* in Scotland in 1988–89. *Bird Study* 40: 1–11.
- BirdLife International (2004) *Birds in Europe: population estimates, trends and conservation status*. BirdLife Conservation Series No. 12. BirdLife International, Cambridge. (BiE04) (www.birdlife.org/action/science/species/birds_in_europe/species_search.html)
- Bolton, M., Tyler, G., Smith, K. & Bamford, R. (2007) The impact of predator control on lapwing *Vanellus vanellus* breeding success on wet grassland nature reserves. *Journal of Applied Ecology* 44: 534–544.
- Both, C. (2002) Nemen Bonte Vliegenvangers *Ficedula hypoleuca* af door klimaatsverandering? [Decrease of European Pied Flycatchers due to climate change?] *Limosa* 75: 73–78.
- Both, C., Bouwhuis, S., Lessells, C.M. & Visser, M.E. (2006) Climate change and population declines in a long-distance migratory bird. *Nature* 441 (4): 81–83. [Full text](#)
- Both, C., van Asch, M., Bijlsma, R.G., van den Burg, A.B. & Visser, M.E. (2009) Climate change and unequal phenological changes across four trophic levels: constraints or adaptations? *Journal of Animal Ecology* 78: 73–83.
- Brickle, N.W. & Harper, D.G. (2002) Agricultural intensification and the timing of breeding of Corn Buntings *Miliaria calandra*. *Bird Study* 49: 219–228.
- Brindley, E., Norris, K., Cook, T., Babbs, S., Forster-Browne, C. & Yaxley, R. (1998) The abundance and conservation status of redshank (*Tringa totanus*) nesting on saltmarshes in Great Britain. *Biological Conservation* 86: 289–297.
- Brooke, M. de L. & Davies, N.B. (1987) Recent changes in host usage by cuckoos *Cuculus canorus* in Britain. *Journal of Animal Ecology* 56: 873–883.
- Brown, A.F., Crick, H.Q.P. & Stillman, R.A. (1995) The distribution, numbers and breeding ecology of Twite *Acanthis flavirostris* in the south Pennines of England. *Bird Study* 42: 107–121.
- Browne, S.J. & Aebischer, N.J. (2004) Temporal changes in the breeding ecology of European Turtle Doves *Streptopelia turtur* in Britain, and implications for conservation. *Ibis* 146: 125–137.
- Browne, S. & Aebischer, N. (2005) Studies of West Palearctic birds: Turtle Dove. *British Birds* 98: 58–72.
- Browne, S., Vickery, J.A. & Chamberlain, D.E. (2000). Densities and population estimates of breeding Skylarks *Alauda arvensis* in Britain in 1997. *Bird Study* 47: 52–65. [Abstract](#)
- Browne, S.J., Aebischer, N.J. & Crick, H.Q.P. (2005) The breeding ecology of Turtle Doves *Streptopelia turtur* in Britain during the period 1941 to 2000: an analysis of BTO Nest Record Cards. *Bird Study* 52: 1–9. [Abstract](#)
- Buchanan, G.M., Pearce-Higgins, J.W., Wotton, S.R., Grant, M.C. & Whitfield, D.P. (2003) Correlates of the change in Ring Ouzel *Turdus torquatus* abundance in Scotland from 1988–91 to 1999. *Bird Study* 50: 97–105.
- Buckland, S.T., Cattanach, K.L. & Anganuzzi, A.A. (1992) Estimating trends in abundance of dolphins associated with tuna in the eastern tropical Pacific Ocean, using sightings data collected on commercial tuna vessels. *Fishery Bulletin* 90: 1–20.
- Burfield, I.J. & Brooke, M. de L. (2005) The decline of the Ring Ouzel *Turdus torquatus* in Britain: evidence from bird observatory data. *Ringling & Migration* 22: 199–204.
- Burton, N.H.K. (2007) Influences of restock age and habitat patchiness on Tree Pipits *Anthus trivialis* breeding in Breckland pine plantations. *Ibis* 149 (suppl. 2): 193–204. [Abstract](#)
- Burton, N.H.K. & Conway, G.J. (2008) *Assessing population change of breeding Ringed Plovers in the UK between 1984 & 2007*. Research Report 503. BTO, Thetford.
- Buse, A., Dury, S.J., Woodburn, R.J.W., Perrins, C.M. & Good, J.E.G. (1999) Effects of elevated temperature on multi-species interactions: the case of pedunculate oak, winter moth and tits. *Functional Ecology* 13 (suppl.): 74–82.
- Butler, C. (2002) Breeding parrots in Britain. *British Birds* 95: 345–348.
- Butler, C.J. (2003) Population biology of the introduced Rose-ringed Parakeet *Psittacula krameri* in the UK. PhD thesis, University of Oxford.

- Catchpole, E.A., Morgan, B.J.T., Freeman, S.N. & Peach, W.J. (1999) Modelling the survival of British Lapwings *Vanellus vanellus* using ring-recovery data and weather covariates. *Bird Study* 46 (suppl.): 5–13.
- Chamberlain, D.E. & Crick, H.Q.P. (1999) Population declines and reproductive performance of skylarks *Alauda arvensis* in different regions and habitats of Great Britain. *Ibis* 141: 38–51.
- Chamberlain, D.E. & Crick, H.Q.P. (2003) Temporal and spatial associations in aspects of reproductive performance of Lapwings *Vanellus vanellus* in the United Kingdom, 1962–99. *Ardea* 91: 183–196. [Abstract](#)
- Chamberlain, D.E. & Siriwardena, G.M. (2000) The effects of agricultural intensification on Skylarks *Alauda arvensis*: evidence from monitoring studies in Great Britain. *Environmental Reviews* 8: 95–113.
- Chamberlain, D.E., Toms, M.P., Cleary-McHarg, R. & Banks, A.N. (2007) House Sparrow (*Passer domesticus*) habitat use in urbanized landscapes. *Journal of Ornithology* 148: 453–462. [Abstract](#)
- Clements, R. (2001) The Hobby in Britain: a new population estimate. *British Birds* 94: 402–408.
- Clements, R. (2002) The Common Buzzard in Britain: a new population estimate. *British Birds* 95: 377–383.
- Clements, R. (2008) The Common Kestrel population in Britain. *British Birds* 101: 228–234.
- Conway, G. & Burton, N. (2009) Changing fortunes for breeding plovers. *BTO News* 280: 10–11.
- Conway, G., Wotton, S., Henderson, I., Langston, R., Drewitt, A. & Currie, F. (2007) Status and distribution of European Nightjars *Caprimulgus europaeus* in the UK in 2004. *Bird Study* 54: 98–111. [Abstract](#)
- Conway, G.J., Burton, N.H.K., Handschuh, M. & Austin, G.E. (2008) *UK population estimates from the 2007 Breeding Little Ringed Plover and Ringed Plover Surveys*. Research Report 510. BTO, Thetford.
- Cowley E. & Siriwardena, G.M. (2005) Long-term variation in survival rates of Sand Martins *Riparia riparia*: dependence on breeding and wintering ground weather, age and sex, and their population consequences. *Bird Study* 52: 237–251. [Abstract](#)
- Crawley, M.J. (1993) *GLIM for Ecologists*. Blackwell Science, Oxford, UK.
- Crick, H.Q.P. (1992) *Trends in the breeding performance of Golden Plover in Britain*. Research Report 76. BTO, Thetford.
- Crick, H.Q.P. (1993) Trends in breeding success of Merlins (*Falco columbarius*) in Britain from 1937–1989. In Nicholls, M.K. & Clarke, R. (eds.) *Biology and Conservation of Small Falcons*, pp 30–38. Hawk & Owl Trust, London.
- Crick, H.Q.P. (1997) Long-term trends in Corn Bunting *Miliaria calandra* productivity in Britain. In Donald, P.F. & Aebischer, N.J. (eds.) *The Ecology and Conservation of Corn Buntings Miliaria calandra*: 52–64. UK Nature Conservation No. 13. JNCC, Peterborough.
- Crick, H.Q.P. (1998) Decline in clutch size of Hen Harriers: reply. *BTO News* 218: 23.
- Crick, H.Q.P. & Ratcliffe, D.A. (1995) The Peregrine *Falco peregrinus* population of the United Kingdom in 1991. *Bird Study* 42: 1–19.
- Crick, H.Q.P. & Sparks, T.H. (1999) Climate change related to egg-laying trends. *Nature* 399: 423–424.
- Crick, H.Q.P., Dudley, C. & Glue, D.E. (1993) Breeding birds in 1991. *BTO News* 185: 15–18.
- Crick, H.Q.P., Dudley, C., Evans, A.D. & Smith, K.W. (1994) Causes of nest failure among buntings in the UK. *Bird Study* 41: 88–94.
- Crick, H.Q.P., Dudley, C., Glue, D.E. & Thomson, D.L. (1997) UK birds are laying eggs earlier. *Nature* 388: 526.
- Crick, H.Q.P., Baillie, S.R., Balmer, D.E., Bashford, R.I., Beaven, L.P., Dudley, C., Glue, D.E., Gregory, R.D., Marchant, J.H., Peach, W.J. & Wilson, A.M. (1998) *Breeding birds in the wider countryside: their conservation status (1972–1996)*. Research Report 198. BTO, Thetford.
- Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D. (eds) (2002) *Investigations into the causes of the decline of Starlings and House Sparrows in Great Britain*. BTO Research Report 290. 305 pp. DEFRA, Bristol.
- Crick, H.Q.P., Baillie, S.R. & Leech, D.I. (2003) The UK Nest Record Scheme: its value for science and conservation. *Bird Study* 50: 254–270.
- Cross, T. (2002) Common Raven (Raven) *Corvus corax*. In *The Migration Atlas: movements of the birds of Britain and Ireland* (eds C.V. Wernham, M.P. Toms, J.H. Marchant, J.A. Clark, G.M. Siriwardena & S.R. Baillie), pp 626–628. T. & A.D. Poyser, London.
- Delany, S., Greenwood, J.J.D. & Kirby, J. (1992) *National Mute Swan Survey 1990*. Report to the Joint Nature Conservation Committee. Wildfowl & Wetlands Trust, Slimbridge. [Full text](#)
- Dixon, A., Richards, C., Haffield, P., Roberts, G., Thomas, M. & Lowe, A. (2008) The National Peregrine Survey 2002: how accurate are the published results for Wales? *Welsh Birds* 5: 276–283.
- Dobson, A.P. & Hudson, P.J. (1992) Regulation and stability of a free-living host–parasite system

- Trichostrongylus tenuis* in Red Grouse. II Population models. *Journal of Animal Ecology* 61: 487–500.
- Dolton, C.S. & Brooke M. de L. (1999) Changes in the biomass of birds breeding in Great Britain, 1968–88. *Bird Study* 46: 274–278.
- Donald, P.F. (1997) The Corn Bunting *Miliaria calandra* in Britain: a review of current status, patterns of decline and possible causes. In Donald, P.F. & Aebischer, N.J. (eds.) *The Ecology and Conservation of Corn Buntings Miliaria calandra*: 11–26. UK Nature Conservation No. 13. Joint Nature Conservation Committee, Peterborough.
- Donald, P.F. & Morris, T.J. (2005) Saving the Sky Lark: new solutions for a declining farmland bird. *British Birds* 98: 570–578.
- Donald, P.F. & Vickery, J.A. (2000) The importance of cereal fields to breeding and wintering skylarks *Alauda arvensis* in the UK. In *Proceedings of the 1999 BOU Spring Conference: Ecology and Conservation of Lowland Farmland Birds* (eds N.J. Aebischer, A.D. Evans, P.V. Grice & J.A. Vickery), pp 140–150. British Ornithologists' Union, Tring.
- Donald, P.F. & Vickery, J.A. (2001) *The ecology and conservation of Skylarks*. RSPB, Sandy.
- Dougall, T.W., Holland, P.K. & Yalden, D.W. (2004) A revised estimate of the breeding population of Common Sandpipers *Actitis hypoleucos* in Great Britain and Ireland. *Wader Study Group Bulletin* 105: 42–49.
- Driver, J. (2006) Raven *Corvus corax* population census of northwest Wales, 1998 to 2005. *Welsh Birds* 4: 442–453.
- Elliott, G.R. & Avery, M.I. (1991) A review of reports of buzzard persecution 1975–1989. *Bird Study* 38: 52–56.
- Eraud, C., Boutin, J.-M., Riviere, M., Brun, J., Barbraud, C. & Lormee, H. (2009) Survival of Turtle Doves *Streptopelia turtur* in relation to western Africa environmental conditions. *Ibis* 151: 186–190.
- Etheridge, B., Summers, R.W. & Green, R.E. (1997) The effects of illegal killing and destruction of nests by humans on the population dynamics of the hen harrier *Circus cyaneus* in Scotland. *Journal of Applied Ecology* 34: 1081–1105.
- Evans, K.L. & Robinson, R.A. (2004) Barn Swallows and agriculture. *British Birds* 97: 218–230.
- Fewster, R.M., Buckland, S.T., Siriwardena, G.M., Baillie, S.R. & Wilson, J.D. (2000) Analysis of population trends for farmland birds using generalized additive models. *Ecology* 81: 1970–1984.
- Freeman, S.N. & Crick, H.Q.P. (2002) Population dynamics of House Sparrows *Passer domesticus* breeding in Britain: an integrated analysis. pp 193–211 in Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D. (eds) *Investigation into the causes of the decline of starlings and house sparrows in Great Britain*. Research Report 290. BTO, Thetford.
- Freeman, S.N. & Crick, H.Q.P. (2003) The decline of the Spotted Flycatcher *Muscicapa striata* in the UK: an integrated population model. *Ibis* 145: 400–412.
- Freeman, S.N., Wernham, C.V. & Balmer, D.E. (2001) *Long-term changes in the productivity of common songbirds in Britain and Ireland from constant effort ringing*. Unpublished. BTO, Thetford.
- Freeman, S.N., Robinson, R.A., Clark, J.A., Griffin, B.M. & Adams, S.Y. (2002) Population dynamics of Starling *Sturnus vulgaris* breeding in Britain: an integrated analysis. Pp 121–139 in Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D. (eds) *Investigation into the causes of the decline of starlings and house sparrows in Great Britain*. Research Report 290. BTO, Thetford.
- Freeman, S.N., Noble, D.G., Newson, S.E. & Baillie, S.R. (2003) *Modelling bird population changes using data from the Common Birds Census and the Breeding Bird Survey*. Research Report 303. BTO, Thetford. [Abstract](#)
- Freeman, S.N., Noble, D.G., Newson, S.E. & Baillie, S.R. (2007a) Modelling bird population changes using data from different surveys: the Common Birds Census and the Breeding Bird Survey. *Bird Study* 54: 61–72. [Abstract](#)
- Freeman, S.N., Robinson, R.A., Clark, J.A., Griffin, B.M. & Adams, S.Y. (2007b) Changing demography and population decline in the Common Starling *Sturnus vulgaris*: a multisite approach to Integrated Population Monitoring. *Ibis* 149: 587–596. [Abstract](#)
- Fuller, R.J. (1996) *Relationships between grazing and birds with particular reference to sheep in the British uplands*. Research Report 164. BTO, Thetford.
- Fuller, R. & Hoodless, A. (2004) Breeding Woodcock Survey, 2003. *BTO News* 253: 4–5.
- Fuller, R.J., Marchant, J.H. & Morgan, R.A. (1985) How representative of agricultural practice in Britain are Common Birds Census farmland plots? *Bird Study* 32: 56–70.
- Fuller, R.J., Gregory, R.D., Gibbons, D.W., Marchant, J.H., Wilson, J.D., Baillie, S.R. & Carter, N. (1995) Population declines and range contractions among lowland farmland birds in Britain. *Conservation Biology* 9: 1425–1441.
- Fuller, R.J., Noble, D.G., Smith, K.W. & Vanhinsbergh, D. (2005) Recent declines in populations of woodland birds in Britain: a review of possible causes. *British Birds* 98: 116–143. [Abstract](#)
- Furness, R.W. & Greenwood, J.J.D. (eds.) (1993) *Birds as Monitors of Environmental Change*.

Chapman & Hall, London.

Galbraith, H. (1988) Effects of agriculture on the breeding ecology of lapwings *Vanellus vanellus*. *Journal of Applied Ecology* 25: 487–503.

Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993) *The New Atlas of Breeding Birds in Britain and Ireland: 1988–1991*. T. & A.D. Poyser, London.

Gibbons, D.W., Avery, M.I., Baillie, S.R., Gregory, R.D., Kirby, J., Porter, R.F., Tucker, G.M. & Williams, G. (1996) Bird species of conservation concern in the United Kingdom, Channel Islands and Isle of Man: revising the Red Data List. *RSPB Conservation Review* 10: 7–18.

Gibbons, D.W., Bainbridge, I.P., Mudge, G.P., Tharme, A.P. & Ellis, P.M. (1997) The status and distribution of the Red-throated Diver *Gavia stellata* in Britain in 1994. *Bird Study* 44: 194–205.

Gill, R.M.A. & Fuller, R.J. (2007) The effects of deer browsing on woodland structure and songbirds in lowland Britain. *Ibis* 149 suppl. 2: 119–127. [Abstract](#)

Gillings, S., Newson, S.E., Noble, D.G. & Vickery, J.A. (2005) Winter availability of cereal stubbles attracts declining farmland birds and positively influences breeding population trends. *Proceedings of the Royal Society B, Biological Sciences* 272: 733–739. doi 10.1098/rspb.2004.3010 [Abstract/Full text](#)

Gillings, S., Newson, S. & Sellers, R.M. (2007) Breeding population estimates for Northern Wheatear in Britain. *British Birds* 100: 179–181.

Glue, D.E. (1990) Breeding biology of the Grasshopper Warbler in Britain. *British Birds* 83: 131–145.

Glue, D.E. (2006) Cuckoos in crisis? *BTO News* 263: 22–23.

Goss-Custard, J.C. (1993) The effect of migration and scale on the study of bird populations: 1991 Witherby Lecture. *Bird Study* 40: 81–96.

Green, R.E. (1999) Applications of large-scale studies of demographic rates to bird conservation. *Bird Study* 46 (Suppl.): S279–S288.

Greenwood, J.J.D. (2000). How BTO's monitoring of birds contributes to conservation. In *European Monitoring for Nature Conservation, Bonn-Bad Godesberg (Bundesamt für Naturschutz)*, ed. by C. Bischoff & R. Droschmeister. *Schriftenreihe für Landschaftspflege und Naturschutz* 62: 105–117.

Gregory, R.D. & Marchant, J.H. (1996) Population trends of Jays, Magpies, Jackdaws and Carrion Crows in the United Kingdom. *Bird Study* 43: 28–37.

Gregory, R.D., Carter, S.P. & Baillie, S.R. (1997) Abundance, distribution and habitat use of breeding Goosanders *Mergus merganser* and Red-breasted Mergansers *Mergus serrator* on British rivers. *Bird Study* 44: 1–12.

Gregory, R.D., Wilkinson, N.I., Noble, D.G., Robinson, J.A., Brown, A.F., Hughes, J., Procter, D., Gibbons, D.W. & Galbraith, C.A. (2002) The population status of birds in the United Kingdom, Channel Islands and Isle of Man: an analysis of conservation concern 2002–2007. *British Birds* 95: 410–448.

Gregory, R.D., Noble, D.G. & Custance, J. (2004) The state of play of farmland birds: population trends and conservation status of lowland farmland birds in the United Kingdom. *Ibis* 146 (suppl. 2): 1–13.

Gruar, D., Barritt, D. & Peach, W.J. (2006) Summer utilization of Oilseed Rape by Reed Buntings *Emberiza schoeniclus* and other farmland birds. *Bird Study* 53: 47–54.

Hardey, J., Crick, H.Q.P., Wernham, C.V., Riley, H.T., Etheridge, B. & Thompson, D.B.A. (2006) *Raptors: a field guide to survey and monitoring*. The Stationery Office, Edinburgh.

Hastie, T.J. & Tibshirani, R.J. (1990) *Generalized additive models*. Chapman & Hall, London.

Henderson, I.G. & Hart, P.J.B. (1993) Provisioning, parental investment and reproductive success in Jackdaws *Corvus monedula*. *Ornis Scandinavica* 24: 142–148.

Henderson, I.G., Wilson, A.M., Steele, D. & Vickery, J.A. (2002) Population estimates, trends and habitat associations of breeding Lapwing *Vanellus vanellus*, Curlew *Numenius arquata* and Snipe *Gallinago gallinago* in Northern Ireland in 1999. *Bird Study* 49: 17–25. [Abstract](#)

Henderson, I., Holt, C. & Vickery, J. (2007) National and regional patterns of habitat association with foraging Barn Swallows *Hirundo rustica* in the UK. *Bird Study* 54: 371–377. [Abstract](#)

Hinsley, S.A., Bellamy, P.E., Newton, I. & Sparks, T.H. (1995) Habitat and landscape factors influencing the presence of individual breeding bird species in woodland fragments. *Journal of Avian Biology* 26: 94–104.

Hinsley, S.A., Carpenter, J.E., Broughton, R.K., Bellamy, P.E., Rothery, P., Amar, A., Hewson, C.M. & Gosler, A.G. (2007) Habitat selection by Marsh Tits *Poecile palustris* in the UK. *Ibis* 149 (suppl. 2): 224–233.

Holland, P.K. & Yalden, D.W. (2002) Population dynamics of Common Sandpipers *Actitis hypoleucos* in the Peak District of Derbyshire – a different decade: a report of the failure of a population to recover from a catastrophic snow storm. *Bird Study* 49: 131–138.

Holling, M. & the Rare Breeding Birds Panel (2007a) Rare breeding birds in the United Kingdom in 2003 and 2004. *British Birds* 100: 321–367. [Full text](#)

- Holling, M. & the Rare Breeding Birds Panel (2007b) Non-native breeding birds in the United Kingdom in 2003, 2004 and 2005. *British Birds* 100: 638–649. [Full text](#)
- Holling, M. & the Rare Breeding Birds Panel (2008) Rare breeding birds in the United Kingdom in 2005. *British Birds* 101: 276–316.
- Holloway, S. (1996) *The Historical Atlas of Breeding Birds in Britain and Ireland 1875–1900*. T. & A.D. Poyser, London.
- Hudson, P.J. (1992) *Grouse in space and time*. Game Conservancy Trust, Fordingbridge.
- Hudson, R. (1972) Collared Doves in Britain and Ireland during 1965–70. *British Birds* 65: 139–155.
- Hudson, R., Tucker, G.M. & Fuller, R.J. (1994) Lapwing *Vanellus vanellus* populations in relation to agricultural changes: a review. In Tucker, G.M., Davies, S.M. & Fuller, R.J. (eds) *The Ecology and Conservation of Lapwings Vanellus vanellus*: 1–33. UK Nature Conservation No 9. JNCC, Peterborough.
- Hughes, S.W.M., Bacon, P., & Flegg, J.J.M. (1979) The 1975 census of the Great Crested Grebe in Britain. *Bird Study* 26: 213–226.
- Hutchinson, C.D. (1989) *Birds in Ireland*. T. & A.D. Poyser, Calton.
- Jackson, D.B. (2007) Factors affecting the abundance of introduced hedgehogs (*Erinaceus europaeus*) to the Hebridean island of South Uist in the absence of natural predators and implications for nesting birds. *Journal of Zoology* 271: 210–217.
- Jackson, D.B., Fuller, R.J. & Campbell, S.T. (2004) Long-term population changes among breeding shorebirds in the Outer Hebrides, Scotland, in relation to introduced hedgehogs (*Erinaceus europaeus*). *Biological Conservation* 117: 151–166.
- JNCC (1996) *Birds of Conservation Importance*. Press release (31 May 1996). Joint Nature Conservation Committee, Peterborough.
- Johnson, D.H. (1979) Estimating nest success: The Mayfield method and an alternative. *Auk* 96: 651–661.
- Johnstone, I., Dyda, J. & Lindley, P. (2007) The population status and hatching success of Curlews *Numenius arquata* in Wales in 2006. *Welsh Birds* 5: 78–87.
- Johnstone, I., Dyda, J. & Lindley, P. (2008) The population status of breeding Golden Plover and Dunlin in Wales in 2007. *Welsh Birds* 5: 300–310.
- Joy, A.C., Noble, D.G. & Baillie, S.R. (2003) *Evaluation of species coverage and precision using the BBS indexing method*. Research Report 317. BTO, Thetford. [Abstract](#)
- King, R., Brooks, S.P., Mazzetta, C., Freeman, S.N. & Morgan, B.J.T. (2008) Identifying and diagnosing population declines: a Bayesian assessment of Lapwings in the UK. *Applied Statistics* 57: 609–632.
- Krebs, J.R., Wilson, J.D., Bradbury R.B. & Siriwardena, G.M. (1999) The second silent spring? *Nature* 400: 611–612.
- Kyrkos, A. (1997) *Behavioural and demographic responses of yellowhammers to variation in agricultural practices*. D.Phil. thesis, University of Oxford.
- Langston, R.H.W., Liley, D., Murison, G., Woodfield, E. & Clarke, R.T. (2007) What effects do walkers and dogs have on the distribution and productivity of breeding European Nightjar *Caprimulgus europaeus*? *Ibis* 149, supplement 1: 27–36.
- Leech, D. & Barimore, C. (2008) Is avian breeding success weathering the storms? *BTO News* 279: 19–20.
- Leech, D. & Crick, H. (2005) Nest Record Scheme breeding trends – latest results. *BTO News* 261: 18–19.
- Leech, D.I., Crick, H.Q.P. & Shawyer, C.R. (2005) *The BTO Barn Owl Monitoring Programme: fourth year 2003*. Research Report 411. BTO, Thetford. [Abstract](#)
- Leech, D., Crick, H. & Shawyer, C. (2006a) Barn Owls and winter weather. *BTO News* 262: 8–9.
- Leech, D., Barimore, C. & Crick, H. (2006b) NRS Concern List – five new species added. *BTO News* 267: 4–5.
- Leech, D., Barimore, C. & Crick, H. (2007) Volunteer boom. *BTO News* 273: 18–19.
- Lewis, A.J.G., Amar, A., Cordi-Piec, D. & Thewlis, R.M. (2007) Factors influencing Willow Tit *Poecile montanus* site occupancy: a comparison of abandoned and occupied woods. *Ibis* 149 (suppl. 2): 205–213. [Abstract](#)
- Liley, D. & Sutherland, W.J. (2007) Predicting the population consequences of human disturbance for Ringed Plovers *Charadrius hiaticula*: a game theory approach. *Ibis* 149, supplement 1: 82–94.
- MacDonald, M.A. & Bolton, M. (2008) Predation of Lapwing *Vanellus vanellus* nests on lowland wet grassland in England and Wales: effects of nest density, habitat and predator abundance. *Journal of Ornithology* 149: 555–563.

- Macleán, I.M.D & Austin, G.E. (2008) *Wetland Bird Survey alerts 2004/2005 Release 2: Changes in numbers of wintering waterbirds in the constituent countries of the United Kingdom, Special Protection Areas (SPAs) and Sites of Special Scientific Interest (SSSIs)*. Research Report 492. British Trust for Ornithology, Thetford. [Full text](#)
- Mallord, J.W., Dolman, P.M., Brown, A.F. & Sutherland, W.J. (2007) Linking recreational disturbance to population size in a ground-nesting passerine. *Journal of Applied Ecology* 44: 185–195. doi 10.1111/j.1365-2664.2006.01242.x
- Manly, B.F.J. (1991) *Randomisation and Monte Carlo Methods in Biology*. Chapman & Hall, London.
- Marchant, J.H. & Gregory, R.D. (1999) Numbers of nesting Rooks *Corvus frugilegus* in the United Kingdom in 1996. *Bird Study* 46: 258–273. [Abstract](#)
- Marchant, J.H., Hudson, R., Carter, S.P. & Whittington, P.A. (1990) *Population Trends in British Breeding Birds*. BTO, Tring.
- Marchant, J.H., Freeman, S.N., Crick, H.Q.P. & Beaven, L.P. (2004) The BTO Heronries Census of England and Wales 1928–2000: new indices and a comparison of analytical methods. *Ibis* 146: 323–334. [Abstract](#)
- Marquiss, M. (2007) Seasonal pattern in hawk predation on Common Bullfinches *Pyrrhula pyrrhula*: evidence of an interaction with habitat affecting food availability. *Bird Study* 54: 1–11.
- Marquiss, M., Newton, I. & Ratcliffe, D.A. (1978) The decline of the Raven *Corvus corax* in relation to afforestation in southern Scotland and northern England. *Journal of Applied Ecology* 15: 129–144.
- Mason, C.F. & Macdonald, S.M. (2006) Recent marked decline in Corn Bunting numbers in northeast Essex. *British Birds* 99: 206–214.
- Mavor, R.A., Parsons, M., Heubeck, M., Pickerell, G. & Schmitt, S. (2003) *Seabirds numbers and breeding success in Britain and Ireland, 2002*. UK Nature Conservation no. 27. JNCC, Peterborough. [Abstract/Full text](#)
- Mavor, R.A., Parsons, M., Heubeck, M. & Schmitt, S. (2004) *Seabird numbers and breeding success in Britain and Ireland, 2003*. UK Nature Conservation no. 28. JNCC, Peterborough. [Abstract/Full text](#)
- Mavor, R.A., Parsons, M., Heubeck, M. & Schmitt, S. (2006) *Seabird numbers and breeding success in Britain and Ireland, 2005*. UK Nature Conservation no. 30. JNCC, Peterborough. [Abstract/Full text](#)
- Mavor, R.A., Heubeck, M., Schmitt, S. & Parsons, M. (2008) *Seabird numbers and breeding success in Britain and Ireland, 2006*. UK Nature Conservation no. 31. JNCC, Peterborough. [Abstract/Full text](#)
- Mayfield, H. (1961) Nesting success calculated from exposure. *Wilson Bulletin* 73: 255–261.
- Mayfield, H. (1975) Suggestions for calculating nest success. *Wilson Bulletin* 87: 456–466.
- Mearns, R. (1983) The status of the Raven in southern Scotland and Northumbria. *Scottish Birds* 12: 211–218.
- Messenger, A. & Roome, M. (2007) The breeding population of the Hobby in Derbyshire. *British Birds* 100: 594–608.
- Milsom, T.P. (2005) Decline of Northern Lapwing *Vanellus vanellus* breeding on arable farmland in relation to loss of spring tillage. *Bird Study* 52: 297–306.
- Mitchell, C., Patterson, D., Boyer, P., Cunningham, P., McDonald, R., Meek, E., Okill, J.D. & Symonds, F. (2000) The summer status and distribution of Greylag Geese in north and west Scotland. *Scottish Birds* 21: 69–77.
- Mitchell, C., King, R. & Cook, T. (2002) Mallard *Anas platyrhynchos*. In *The Migration Atlas: movements of the birds of Britain and Ireland* (eds C.V. Wernham, M.P. Toms, J.H. Marchant, J.A. Clark, G.M. Siriwardena & S.R. Baillie), pp 193–195. T. & A.D. Poyser, London.
- Mitchell, P.I., Newton, S.F., Ratcliffe, N. & Dunn, T.E. (2004) *Seabird Populations of Britain and Ireland*. T. & A.D. Poyser, London. [Summary](#)
- Monaghan, P., Uttley, J.D., Burns, M.D., Thaine, C. & Blackwood, J. (1989) The relationship between food supply, reproductive effort and breeding success in Arctic Terns *Sterna paradisaea*. *Journal of Animal Ecology* 58: 261–274.
- Monaghan, P., Uttley, J.D. & Burns, M.D. (1992) Effect of changes in food availability on reproductive effort in Arctic terns *Sterna paradisaea*. *Ardea* 80: 71–81.
- Moorcroft, D., Wilson, J.D. & Bradbury, R.B. (2006) The diet of nestling Linnets *Carduelis cannabina* on lowland farmland before and after agricultural intensification. *Bird Study* 53: 156–162.
- Morris, A., Burges, D., Fuller, R.J., Evans, A.D. & Smith, K.W. (1994) The status and distribution of Nightjars *Caprimulgus europaeus* in Britain in 1992. *Bird Study* 41: 181–191.
- Moss, D. & Moss, G.M. (1993) Breeding biology of the Little Grebe *Tachybaptus ruficollis* in Britain and Ireland. *Bird Study* 40: 107–114.
- Mountford, M.D. (1982) Estimation of population fluctuations with application to the Common Birds Census. *Applied Statistics* 31: 135–143.

- Mountford, M.D. (1985) An index of population change with an application to the Common Birds Census. In Morgan, B.J.T. & North, P.M. (eds) *Statistics in Ornithology*: 121–132. Springer-Verlag, Berlin.
- Nelson, S.H., Court, I., Vickery, J.A., Watts, P.N. & Bradbury, R.B. (2003) The status and ecology of the Yellow Wagtail in Britain. *British Wildlife* 14: 270–274.
- Newson, S.E., Woodburn, R.J.W., Noble, D.G., Baillie, S.R. & Gregory, R.D. (2005) Evaluating the Breeding Bird Survey for producing national population size and density estimates. *Bird Study* 52: 42–54. [Abstract](#)
- Newson, S.E., Ekins, G.R., Marchant, J.H., Rehfisch, M.M. & Sellers, R.M. (2006) *The status of inland and coastal breeding Great Cormorants Phalacrocorax carbo in England*. Research Report 433. BTO, Thetford. [Abstract](#)
- Newson, S.E., Marchant, J.H., Ekins, G.R. & Sellers, R.M. (2007) The status of inland-breeding Great Cormorants in England. *British Birds* 100: 289–299. [Abstract](#)
- Newson, S.E., Evans, K.L., Noble, D.G., Greenwood, J.J.D. & Gaston, K.J. (2008) Use of distance sampling to improve estimates of national population sizes for common and widespread breeding birds in the UK. *Journal of Applied Ecology* 45: 1330–1338. doi: 10.1111/j.1365-2664.2008.01480x [Abstract](#)
- Newton, I. (1986) *The Sparrowhawk*. T. & A.D. Poyser, Calton.
- Newton, I. (2004) The recent declines of farmland bird populations in Britain: an appraisal of causal factors and conservation actions. *Ibis* 146: 579–600.
- Norris, K., Brindley, E., Cook, T., Babbs, S., Forster-Brown, C. & Yaxley, R. (1998) Is the density of Redshank *Tringa totanus* nesting on saltmarshes in Great Britain declining due to changes in grazing management? *Journal of Applied Ecology* 35: 621–634.
- O'Brien, M. (2005) Estimating the number of farmland waders breeding in the United Kingdom. *International Wader Studies* 14: 135–139.
- O'Connor, R.J. & Marchant, J.H. (1981) *A field validation of some Common Birds Census techniques*. Research Report 4. BTO, Tring.
- O'Connor, R.J. & Mead, C.J. (1984) The Stock Dove in Britain, 1930–1980. *British Birds* 77: 181–201.
- O'Connor, R.J. & Shrubbs, M. (1986) *Farming and Birds*. Cambridge University Press, Cambridge.
- Ormerod, S.J. & Tyler, S.J. (1989) Long-term change in the suitability of Welsh streams for Dippers *Cinclus cinclus* as a result of acidification and recovery: a modelling study. *Environmental Pollution* 62: 171–182.
- Ormerod, S.J. & Tyler, S.J. (1990) Environmental pollutants in the eggs of Welsh Dippers *Cinclus cinclus*: a potential monitor of organochlorine and mercury contamination in upland rivers. *Bird Study* 37: 171–176.
- Ormerod, S.J., O'Halloran, J., Gribbin, S.D. & Tyler, S.J. (1991) The ecology of Dippers *Cinclus cinclus* in relation to stream acidity in upland Wales: breeding performance, calcium physiology and nestling growth. *Journal of Applied Ecology* 28: 419–433.
- Owen, M., Atkinson-Willes, G.L. & Salmon, D.G. (1986) *Wildfowl in Great Britain*. Cambridge University Press, Cambridge.
- Pannekoek, J. & van Strien, A. (1996) *TRIM (TRends and Indices for Monitoring data)*. Research paper 9634. Statistics Netherlands, Voorburg.
- Parkin, D.T., Collinson, M., Helbig, A.J., Knox, A.G. & Sangster, G. (2003) The taxonomic status of Carrion and Hooded Crows. *British Birds* 96: 274–290.
- Parr, S.J. (1994) Changes in the population size and nest sites of Merlins *Falco columbarius* in Wales between 1970 and 1991. *Bird Study* 41: 42–47.
- Peach, W.J. (1993) Combining mark recapture data sets for small passerines. In *Marked Individuals in the Study of Bird Populations* (eds J.-D. Lebreton & P.M. North), pp. 107–122. Birkhauser Verlag, Basel, Switzerland.
- Peach, W.J. & Baillie, S.R. (1994) Implementation of the Mountford indexing method for the Common Birds Census. In Hagemeyer, W. & Verstrael, T. (eds.) *Bird Numbers 1992*. Distribution, Monitoring and Ecological Aspects: 653–662. Proc. 12th Int. Conf. International Bird Census Council and European Ornithological Atlas Committee. SOVON, Beek-Ubbergen.
- Peach, W.J., Baillie, S.R. & Underhill, L. (1991) Survival of British Sedge Warblers *Acrocephalus schoenobaenus* in relation to west African rainfall. *Ibis* 133: 300–305.
- Peach, W.J., Thompson, P.S. & Coulson, J.C. (1994) Annual and long-term variation in the survival rates of British lapwings *Vanellus vanellus*. *Journal of Animal Ecology* 63: 60–70.
- Peach, W.J., Crick, H.Q.P. & Marchant, J.H. (1995a) The demography of the decline in the British Willow Warbler population. *Journal of Applied Statistics* 22: 905–922.
- Peach, W.J., du Feu, C. & McMeeking, J. (1995b) Site tenacity and survival rates of Wrens *Troglodytes troglodytes* and Treecreepers *Certhia familiaris* in a Nottinghamshire wood. *Ibis* 137: 497–507.

- Peach, W.J., Buckland, S.T. & Baillie, S.R. (1996) The use of constant effort mist-netting to measure between-year changes in the abundance and productivity of common passerines. *Bird Study* 43: 142–156.
- Peach, W.J., Baillie, S.R. & Balmer, D.E. (1998) Long-term changes in the abundance of passerines in Britain and Ireland as measured by constant effort mist-netting. *Bird Study* 45: 257–275.
- Peach, W.J., Siriwardena, G.M. & Gregory, R.D. (1999) Long-term changes in the abundance and demography of British reed buntings *Emberiza schoeniclus*. *Journal of Applied Ecology* 36: 798–811. [Abstract/Full text](#)
- Peach, W.J., Robinson, R.A. & Murray, K.A. (2004) Demographic and environmental causes of the decline of rural Song Thrushes *Turdus philomelos* in lowland Britain. *Ibis* 146 (Suppl. 2): 50–59. [Abstract](#)
- Peach, W.J., Vincent, K.E., Fowler, J.A. & Grice, P.V. (2008) Reproductive success of house sparrows along an urban gradient. *Animal Conservation* 11: 493–503. doi 10.1111/j.1469-1795.2008.00209.x
- Pearce-Higgins, J.W., Yalden, D.W. & Whittingham, M.J. (2005) Warmer springs advance the breeding phenology of golden plovers *Pluvialis apricaria* and their prey (Tipulidae). *Oecologia* 143: 470–476.
- PECBMS (2007) *State of Europe's Common Birds, 2007*. CSO/RSPB, Prague. [Full text](#)
- Percival, S.M. (1990) *Population trends in British Barn Owls, Tyto alba, and Tawny Owls, Strix aluco, in relation to environmental change*. Research Report 57. BTO, Tring.
- Perkins, A.J., Maggs, H.E., Wilson, J.D., Watson, A. & Smout, C. (2008) Targeted management intervention reduces rate of population decline of Corn Buntings *Emberiza calandra* in eastern Scotland. *Bird Study* 55: 52–58.
- Perrins, C. (2003) The status of Marsh and Willow Tits in the UK. *British Birds* 96: 418–426.
- Pienkowski, M.W. (1991) Using long-term ornithological studies in setting targets for conservation in Britain. *Ibis* 133 (Suppl. 1): 62–75.
- Potts, G.R. (1986) *The Partridge: pesticides, predation and conservation*. Collins, London.
- Prater, A.J. (1989) Ringed Plover *Charadrius hiaticula* breeding population of the United Kingdom in 1984. *Bird Study* 36: 154–159.
- Prince, P. & Clarke, R. (1993) The hobby's breeding range in Britain. *British Wildlife* 4: 341–346.
- Proffitt, F.M., Newton, I., Wilson, J.D. & Siriwardena, G.M. (2004) Bullfinch *Pyrrhula pyrrhula* breeding ecology in lowland farmland and woodland: comparisons across time and habitat. *Ibis* 146 (Suppl. 2): 78–86. [Abstract](#)
- Ratcliffe, D.A. (1993) *The Peregrine Falcon*. Second Edition. T. & A.D. Poyser, London.
- Rebecca, G.W. & Bainbridge, I.P. (1998) The breeding status of the Merlin *Falco columbarius* in Britain in 1993–94. *Bird Study* 45: 172–187.
- Redpath, S. & Thirgood, S. (1997) *Birds of Prey and Red Grouse*. HMSO, London.
- Reed, T. (1985) Estimates of British breeding wader populations. *Wader Study Group Bulletin* 45: 11–12.
- Rehfishch, M.M., Wernham, C.V. & Marchant, J.H. (eds) (1999) *Population, distribution, movements and survival of fish-eating birds in Great Britain*. DETR, London.
- Rehfishch, M.M., Austin, G.E., Holloway, S.J., Allan, J.R. & O'Connell, M. (2002) An approach to the assessment of change in the numbers of Canada Geese *Branta canadensis* and Greylag Geese *Anser anser* in southern Britain. *Bird Study* 49: 50–59.
- Risely, K., Noble, D.G. & Baillie, S.R. (2008) *The Breeding Bird Survey 2007*. Research Report 508. BTO, Thetford. [Abstract](#)
- Robertson, D. (2003) Eurasian Reed Warblers in Scotland: a review of probable breeding records. *Scottish Birds* 24: 36–39.
- Robertson, P.A., Woodburn, M.I.A., Tapper, S.C. & Stoate, C. (1989) *Estimating game densities in Britain from land-use maps*. Institute of Terrestrial Ecology, Grange-over-Sands.
- Robinson, R.A., Siriwardena, G.M. & Crick, H.Q.P. (2002) Status and population trends of the Starling *Sturnus vulgaris* in Great Britain. Pp 11–32. in Crick, H.Q.P., Robinson, R.A., Appleton, G.F. Clark, N.A. & Rickard, A.D. (eds) *Investigation into the causes of the decline of starlings and house sparrows in Great Britain*. Research Report 290. BTO, Thetford.
- Robinson, R.A., Crick, H.Q.P. & Peach, W.J. (2003) Population trends of Swallows *Hirundo rustica* breeding in Britain. *Bird Study* 50: 1–7. [Abstract](#)
- Robinson, R.A., Green, R.E., Baillie, S.R., Peach, W.J. & Thomson, D.L. (2004) Demographic mechanisms of the population decline of the song thrush *Turdus philomelos* in Britain. *Journal of Animal Ecology* 73: 670–682. [Abstract](#)
- Robinson, R.A., Siriwardena, G.M. & Crick, H.Q.P. (2005a) Status and population trends of Starling *Sturnus vulgaris* in Great Britain. *Bird Study* 52: 252–260. [Abstract](#)

- Robinson, R.A., Siriwardena, G.M. & Crick, H.Q.P. (2005b) Size and trends of the House Sparrow *Passer domesticus* population in Great Britain. *Ibis* 147: 552–562. [Abstract](#)
- Robinson, R.A., Freeman, S.N., Balmer, D.E. & Grantham, M.J. (2007) Cetti's Warbler *Cettia cetti*: analysis of an expanding population. *Bird Study* 54: 230–235. [Abstract](#)
- Robinson, R.A., Balmer, D.E. & Marchant, J.H. (2008) Survival rates of hirundines in relation to British and African rainfall. *Ringing & Migration* 24: 1–6.
- Rowell, H.E. & Spray, C.J. (2004) *The Mute Swan Cygnus olor (Britain and Ireland populations) in Britain and Northern Ireland 1960/61 – 2000/01*. Waterbird Review Series, The Wildfowl & Wetlands Trust/Joint Nature Conservation Committee, Slimbridge.
- Sanderson, F.J., Donald, P.F., Pain, D.J., Burfield, I.J. & van Bommel, F.P.J. (2006) Long-term population declines in Afro–Palearctic migrant birds. *Biological Conservation* 131: 93–105.
- SAS Institute Inc. (1990) *SAS Language: Reference, Version 6, First Edition*. SAS Inst. Inc., Cary, NC.
- Sellers, R.M. (2006) Breeding population estimate for Northern Wheatear in Britain. *British Birds* 99: 533–535.
- Sharpe, F., Clark, J. & Leech, D. (2008) Does variation in demographic parameters account for regional variation in Northern Lapwing *Vanellus vanellus* population declines across Great Britain? *Bird Study* 55: 247–256.
- Shaw, L.M., Chamberlain, D.E. & Evans, M.R. (2008) The house sparrow *Passer domesticus* in urban areas: reviewing a possible link between post-decline distribution and human socioeconomic status. *Journal of Ornithology* 149: 293–299. doi 10.1007/s10336-008-0285-y [Abstract](#)
- Sim, I.M.W., Gibbons, D.W., Bainbridge, I.P. & Mattingley, W.A. (2001) Status of the Hen Harrier *Circus cyaneus* in the UK and the Isle of Man in 1998. *Bird Study* 48: 341–353.
- Sim, I.M.W., Dillon, I.A., Eaton, M.A., Etheridge, B., Lindley, P., Riley, H., Saunders, R., Sharpe, C. & Tickner, M. (2007a) Status of the Hen Harrier *Circus cyaneus* in the UK and Isle of Man in 2004, and a comparison with the 1988/89 and 1998 surveys. *Bird Study* 54: 256–267.
- Sim, I.M.W., Burfield, I.J., Grant, M.C., Pearce-Higgins, J.W. & Brooke, M. de L. (2007b) The role of habitat composition in determining breeding site occupancy in a declining Ring Ouzel *Turdus torquatus* population. *Ibis* 149: 374–385.
- Siriwardena, G.M. (2004) Possible roles of habitat, competition and avian nest predation in the decline of the Willow Tit *Parus montanus* in Britain. *Bird Study* 51: 193–202. [Abstract](#)
- Siriwardena, G.M. (2006) Avian nest predation, competition and the decline of British Marsh Tits *Parus palustris*. *Ibis* 148: 255–265. [Abstract](#)
- Siriwardena, G.M., Baillie, S.R. & Wilson, J.D. (1998a). Variation in the survival rates of British farmland passerines with respect to their population trends. *Bird Study* 45: 276–292.
- Siriwardena, G.M., Baillie, S.R., Buckland, S.T., Fewster, R.M., Marchant, J.H. & Wilson, J.D. (1998b) Trends in the abundance of farmland birds: a quantitative comparison of smoothed Common Birds Census indices. *Journal of Applied Ecology* 35: 24–43. [Abstract/Full text](#)
- Siriwardena, G.M., Baillie, S.R. & Wilson, J.D. (1999) Temporal variation in the annual survival rates of six granivorous birds with contrasting population trends. *Ibis* 141: 621–636.
- Siriwardena, G.M., Baillie, S.R., Crick, H.Q.P., Wilson, J.D. & Gates S. (2000a) The demography of lowland farmland birds. In *Proceedings of the 1999 BOU Spring Conference: Ecology and Conservation of Lowland Farmland Birds* (eds. N.J. Aebischer, A.D. Evans, P.V. Grice & J.A. Vickery), pp 117–133. British Ornithologists' Union, Tring.
- Siriwardena, G.M., Baillie, S.R., Crick, H.Q.P. & Wilson, J.D. (2000b) The importance of variation in the breeding performance of seed-eating birds for their population trends on farmland. *Journal of Applied Ecology* 37: 128–148. [Abstract/Full text](#)
- Siriwardena, G.M., Freeman, S.N. & Crick, H.Q.P. (2001) The decline of the Bullfinch *Pyrrhula pyrrhula* in Britain: is the mechanism known? *Acta Ornithologica* 36: 143–152.
- Siriwardena, G.M., Robinson, R.A. & Crick, H.Q.P. (2002) Status and population trends of the house sparrow *Passer domesticus* in Great Britain. Pp 33–52 in Crick, H.Q.P., Robinson, R.A., Appleton, G.F., Clark, N.A. & Rickard, A.D. (eds) *Investigation into the causes of the decline of starlings and house sparrows in Great Britain*. Research Report 290. BTO, Thetford.
- Sitters, H.P., Fuller, R.J., Hoblyn, R.A., Wright, M.T., Cowie, N. & Bowden, C.G.R. (1996) The Woodlark *Lullula arborea* in Britain: population trends, distribution and habitat occupancy. *Bird Study* 43: 172–187.
- Smart, J., Amar, A., O'Brien, M., Grice, P. & Smith, K. (2008) Changing land management of lowland wet grasslands of the UK: impacts on snipe abundance and habitat quality. *Animal Conservation* 11: 339–351.
- Smith, K.W. (2005) Has the reduction in nest-site competition from Starlings *Sturnus vulgaris* been a factor in the recent increase of Great Spotted Woodpecker *Dendrocopos major* numbers in Britain? *Bird Study* 52: 307–313.
- Smith, K.W. (2006) The implications of nest site competition from starlings *Sturnus vulgaris* and the

- effect of spring temperatures on the timing and breeding performance of great spotted woodpeckers *Dendrocopos major* in southern England. *Annales Zoologici Fennici* 43: 177–185. [Full text](#)
- Snow, D.W. (1965) The relationship between census results and the breeding population of birds on farmland. *Bird Study* 12: 287–304.
- Stevens, D.K., Anderson, G.Q.A., Grice, P.V., Norris, K. & Butcher, N. (2008) Predators of Spotted Flycatcher *Muscicapa striata* nests in southern England as determined by digital nest-cameras. *Bird Study* 55: 179–187.
- Stoate, C. & Szczyr, J. (2006) Potential influence of habitat and predation on local breeding success and population in Spotted Flycatchers *Muscicapa striata*. *Bird Study* 53: 328–330.
- Stone, B.H., Sears, J., Cranswick, P.A., Gregory, R.D., Gibbons, D.W., Rehfisch, M.M., Aebischer, N.J. & Reid, J.B. (1997) Population estimates of birds in Britain and in the United Kingdom. *British Birds* 90: 1–22.
- Stroud, D.A., Reed, T.M., Pienkowski, M.W. & Lindsay, R.A. (1987) *Birds, Bogs and Forestry*. Nature Conservancy Council, Peterborough.
- Strubbe, D. & Matthysen, E. (2007) Invasive ring-necked parakeets *Psittacula krameri* in Belgium: habitat selection and impact on native birds. *Ecography* 30: 578–588.
- Summers, R. (1998) The decline in clutch size of Hen Harriers. *BTO News* 218: 23.
- Summers-Smith, J.D. (2003) The decline of the House Sparrow: a review. *British Birds* 96: 439–446.
- Summers-Smith, J.D. (2007) Is unleaded petrol a factor in urban House Sparrow decline? *British Birds* 100: 558–559.
- Szép, T. (1995) Relationship between west African rainfall and the survival of central European Sand Martins *Riparia riparia*. *Ibis* 137: 162–168.
- Tapper, S. (1999) *A question of balance: game animals and their role in the British countryside*. The Game Conservancy Trust, Hampshire, UK.
- Temple, S.A. & Wiens, J.A. (1989) Bird populations and environmental changes: can birds be bio-indicators? *American Birds* 43: 260–270.
- Thomas, G.E. (1993) Estimating annual total heron population counts. *Applied Statistics* 42: 473–486.
- Thomson, D.L., Baillie, S.R. & Peach, W.J. (1997) The demography and age-specific annual survival of British song thrushes *Turdus philomelos* during periods of population stability and decline. *Journal of Animal Ecology* 66: 414–424.
- Tompkins, D.M., Draycott, R.A.H. & Hudson, P.J. (2002) Field evidence for apparent competition mediated via the shared parasites of two gamebird species. *Ecological Letters* 3: 10–14.
- Toms, M.P., Crick, H.Q.P. & Shawyer, C.R. (2000) *Project Barn Owl Final Report*. BTO Research Report 197/ HOT Research Report 98/1. BTO/Hawk & Owl Trust, Thetford.
- Toms, M.P., Crick, H.Q.P. & Shawyer, C.R. (2001) The status of breeding Barn Owls *Tyto alba* in the United Kingdom 1995–97. *Bird Study* 48: 23–37.
- Tryjanowski, P., Sparks, T.H. & Crick, H.Q.P. (2006) Red-backed Shrike (*Lanius collurio*) nest performance in a declining British population: a comparison with a stable population in Poland. *Ornis Fennica* 83: 181–186.
- Tucker, G.M. & Heath, M.F. (1994) *Birds in Europe: their conservation status*. Conservation Series no. 3. BirdLife International, Cambridge.
- Turner, A. (2009) Climate change: a Swallow's eye view. *British Birds* 102: 3–16.
- Vickery, J.A. (1991) Breeding density of Dippers *Cinclus cinclus*, Grey Wagtails *Motacilla cinerea* and Common Sandpipers *Actitis hypoleucos* in relation to the acidity of streams in south-west Scotland. *Ibis* 133: 178–185.
- Vickery, J.A. (1992) The reproductive success of the Dipper *Cinclus cinclus* in relation to the acidity of streams in south-west Scotland. *Freshwater Biology* 28: 195–205.
- Vickery, J.A., Bradbury, R.B., Henderson, I.G., Eaton, M.A. & Grice, P.V. (2004) The role of agri-environment schemes and farm management practices in reversing the decline of farmland birds in England. *Biological Conservation* 119: 19–39.
- Vincent, K.E. (2005) *Investigating the causes of the decline of the urban House Sparrow *Passer domesticus* population in Britain*. Unpublished PhD thesis, De Montfort University. [Full text](#)
- Visser, M.E. & Holleman, L.J.M. (2001) Warmer springs disrupt the synchrony of oak and winter moth phenology. *Proceedings of the Royal Society B, Biological Sciences* 268: 289–294.
- Visser, M.E., van Noordwijk, A.J., Tinbergen, J.M. & Lessells, C.M. (1998) Warmer springs lead to mistimed reproduction in Great Tits (*Parus major*). *Proceedings of the Royal Society B, Biological Sciences* 265: 1867–1870. [Full text](#)
- Wanless, S., Harris, M.P., Redman, P. & Speakman, J.R. (2005) Low energy values of fish as a probable cause of a major seabird breeding failure in the North Sea. *Marine Ecology Progress Series*

Ward, R.M., Cranswick, P.A., Kershaw, M., Austin, G., Brown, A.W., Brown, L.M., Coleman, J.T., Chisholm, H. & Spray, C. (2007) *National Mute Swan Census 2002*. WWT, Slimbridge. [Full text](#)

Watson, M., Aebischer, N.J., Potts, G.R. & Ewald, J.A. (2007) The relative effects of raptor predation and shooting on overwinter mortality of grey partridges in the United Kingdom. *Journal of Applied Ecology* 44: 972–982.

Whittingham, M.J., Swetnam, R.D., Wilson, J.D., Chamberlain, D.E. & Freckleton, R.P. (2005) Habitat selection by yellowhammers *Emberiza citrinella* on lowland farmland at two spatial scales: implications for conservation management. *Journal of Applied Ecology* 42: 270–290. [Abstract/Full text](#)

Wilson, A.M., Marchant, J.H., Gregory, R.D., Siriwardena, G.M. & Baillie, S.R. (1998) *Enhancements for monitoring of opportunistic bird populations*. Research Report 200. BTO, Thetford.

Wilson, A.M., Vickery, J.A. & Browne, S.J. (2001) Numbers and distribution of Northern Lapwings *Vanellus vanellus* breeding in England and Wales in 1998. *Bird Study* 48: 2–17.

Wilson, A.M., Henderson, A.C.B. & Fuller, R.J. (2002) Status of the Common Nightingale *Luscinia megarhynchos* in England at the end of the 20th century with particular reference to climate change. *Bird Study* 49: 193–204. [Abstract](#)

Wilson, A.M., Vickery, J.A., Brown, A., Langston, R.H.W., Smallshire, D., Wotton, S. & Vanhinsbergh, D. (2005) Changes in the numbers of breeding waders on lowland wet grasslands in England and Wales between 1982 and 2002. *Bird Study* 52: 55–69. [Abstract/Detailed summary](#)

Wilson, J.D., Evans, J., Browne, S.J., & King, J.R. (1997) Territory distribution and breeding success of skylarks *Alauda arvensis* on organic and intensive farmland in Southern England. *Journal of Applied Ecology* 34: 1462–1478. [Abstract](#)

Wilson, J.D., Boyle, J., Jackson, D.B., Lowe, B. & Wilkinson, N.I. (2007) Effect of cereal harvesting method on a recent population decline of Corn Buntings *Emberiza calandra* on the Western Isles of Scotland. *Bird Study* 54: 362–370.

Winstanley, D., Spencer, R. & Williamson, K. (1974) Where have all the Whitethroats gone? *Bird Study* 21: 1–14.

Wotton, S.R. & Gillings, S. (2000) The status of breeding Woodlarks *Lullula arborea* in Britain in 1997. *Bird Study* 47: 212–224. [Abstract](#)

Wotton, S., Gibbons, D.W., Dilger, M. & Grice, P.V. (1998) Cetti's Warblers in the United Kingdom and the Channel Islands in 1996. *British Birds* 91: 77–89.

Wotton, S.R., Langston, R.H.W. & Gregory, R.D. (2002) The breeding status of the Ring Ouzel *Turdus torquatus* in the UK in 1999. *Bird Study* 49: 26–34.

Wright, L.J., Hoblyn, R.A., Sutherland, W.J. & Dolman, P.M. (2007) Reproductive success of Woodlarks *Lullula arborea* in traditional and recently colonized habitats. *Bird Study* 54: 315–323. [Abstract](#)

Wright, P.M. (2005) *Merlins of the south-east Yorkshire Dales*. Tarnmoor, Skipton.

[Section 7 – Appendix](#)

[Back to Contents](#)

7. Appendix – Summary tables of changes in population size and breeding performance

7.1 Tables of alerts and population increases from CBC/BBS

- 1a. [CBC/BBS UK alerts – 39 years](#)
- 1b. [CBC/BBS England alerts – 39 years](#)
- 2a. [CBC/BBS UK alerts – 25 years](#)
- 2b. [CBC/BBS England alerts – 25 years](#)
- 3a. [CBC/BBS UK alerts – 10 years](#)
- 3b. [CBC/BBS England alerts – 10 years](#)
- 4a. [CBC/BBS UK alerts – 5 years](#)
- 4b. [CBC/BBS England alerts – 5 years](#)
- 5a. [CBC/BBS UK population increases of >50% – 39 years](#)
- 5b. [CBC/BBS England population increases of >50% – 39 years](#)

7.2 Tables of alerts and population increases from WBS

1. [WBS alerts – 31 years](#)
2. [WBS alerts – 25 years](#)
3. [WBS alerts – 10 years](#)
4. [WBS alerts – 5 years](#)
5. [WBS population increases of >50% – 31 years](#)

7.3 Tables of alerts and population increases from CES

1. [CES adults alerts – 22 years](#)
2. [CES adults alerts – 10 years](#)
3. [CES adults alerts – 5 years](#)
4. [CES adults population increases of >50% – 22 years](#)

7.4 Tables of population declines and increases from BBS

1. [BBS – UK alerts - 11 years](#)
2. [BBS – England alerts - 11 years](#)
3. [BBS – Scotland alerts - 11 years](#)
4. [BBS – Wales alerts - 11 years](#)
5. [BBS – UK alerts - 10 years](#)
6. [BBS – England alerts - 10 years](#)
7. [BBS – Scotland alerts - 10 years](#)
8. [BBS – Wales alerts - 10 years](#)
9. [BBS – UK alerts - 5 years](#)
10. [BBS – England alerts - 5 years](#)
11. [BBS – Scotland alerts - 5 years](#)
12. [BBS – Wales alerts - 5 years](#)
13. [BBS – UK population increases of >50%](#)
14. [BBS – England population increases of >50%](#)
15. [BBS – Scotland population increases of >50%](#)
16. [BBS – Wales population increases of >50%](#)
17. [BBS – Northern Ireland population increases of >50%](#)

[Back to top](#)

[Return to Contents](#)

7.1 Tables of alerts and population increases from CBC/BBS

- 1a. CBC/BBS UK alerts – 39 years
- 1b. CBC/BBS England alerts – 39 years
- 2a. CBC/BBS UK alerts – 25 years
- 2b. CBC/BBS England alerts – 25 years
- 3a. CBC/BBS UK alerts – 10 years
- 3b. CBC/BBS England alerts – 10 years
- 4a. CBC/BBS UK alerts – 5 years
- 4b. CBC/BBS England alerts – 5 years
- 5a. CBC/BBS UK population increases of >50% – 39 years
- 5b. CBC/BBS England population increases of >50% – 39 years

1a. Table of population alerts for CBC/BBS UK 1967-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Grey Partridge	39	119	-88	-91	-83	>50	
Corn Bunting	39	63	-87	-94	-77	>50	
Spotted Flycatcher	39	121	-86	-90	-80	>50	
Turtle Dove	39	106	-85	-90	-76	>50	
Willow Tit	39	43	-85	-93	-72	>50	
Yellow Wagtail	39	71	-74	-88	-46	>50	
Marsh Tit	39	89	-68	-77	-56	>50	
Whitethroat	39	447	-62	-73	-50	>50	
Yellowhammer	39	440	-55	-65	-47	>50	
Song Thrush	39	678	-51	-58	-43	>50	
Bullfinch	39	271	-50	-61	-37	>25	
Little Owl	39	54	-46	-68	-12	>25	
Mistle Thrush	39	447	-43	-51	-31	>25	
Lapwing	39	231	-34	-64	-5	>25	
Dunnock	39	703	-33	-42	-22	>25	

1b. Table of population alerts for CBC/BBS England 1967-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Tree Sparrow	39	83	-97	-99	-94	>50	
Lesser Redpoll	39	43	-90	-96	-78	>50	
Grey Partridge	39	107	-88	-91	-82	>50	
Spotted Flycatcher	39	93	-87	-92	-82	>50	
Turtle Dove	39	105	-85	-91	-76	>50	
Corn Bunting	39	60	-84	-93	-74	>50	
Tree Pipit	39	44	-83	-92	-71	>50	
Willow Tit	39	40	-83	-92	-70	>50	
Starling	39	492	-83	-88	-77	>50	
Linnet	39	367	-75	-81	-66	>50	
Yellow Wagtail	39	69	-72	-85	-38	>50	
Marsh Tit	39	82	-68	-78	-55	>50	
Whitethroat	39	388	-63	-73	-50	>50	
Cuckoo	39	260	-61	-71	-48	>50	
Skylark	39	455	-59	-66	-51	>50	
Willow Warbler	39	399	-58	-70	-46	>50	
Yellowhammer	39	385	-57	-66	-46	>50	
Bullfinch	39	221	-52	-62	-39	>50	
Song Thrush	39	540	-51	-60	-42	>50	
Mistle Thrush	39	367	-49	-57	-39	>25	
Meadow Pipit	39	136	-46	-74	-24	>25	
Sedge Warbler	39	81	-39	-69	-13	>25	
Dunnock	39	582	-37	-46	-26	>25	

2a. Table of population alerts for CBC/BBS UK 1981-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Corn Bunting	25	80	-84	-92	-74	>50	
Turtle Dove	25	123	-82	-88	-75	>50	
Willow Tit	25	44	-82	-90	-69	>50	
Spotted Flycatcher	25	142	-81	-86	-75	>50	
Grey Partridge	25	142	-78	-85	-71	>50	
Yellow Wagtail	25	91	-70	-82	-55	>50	
Lapwing	25	324	-53	-65	-37	>50	
Yellowhammer	25	605	-53	-59	-48	>50	
Little Owl	25	69	-46	-63	-29	>25	
Mistle Thrush	25	615	-38	-45	-31	>25	
Marsh Tit	25	107	-38	-51	-16	>25	
Bullfinch	25	340	-28	-39	-14	>25	

2b. Table of population alerts for CBC/BBS England 1981-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Lesser Redpoll	25	40	-95	-98	-91	>50	
Tree Sparrow	25	80	-93	-97	-88	>50	
Corn Bunting	25	76	-83	-92	-71	>50	
Turtle Dove	25	122	-82	-88	-74	>50	
Tree Pipit	25	48	-82	-91	-69	>50	
Spotted Flycatcher	25	105	-82	-87	-75	>50	
Willow Tit	25	40	-82	-91	-72	>50	
Grey Partridge	25	126	-78	-84	-71	>50	
Starling	25	702	-78	-83	-73	>50	
Yellow Wagtail	25	89	-68	-81	-49	>50	
Cuckoo	25	352	-61	-67	-54	>50	
House Sparrow	25	587	-59	-71	-49	>50	
Willow Warbler	25	532	-58	-67	-49	>50	
Yellowhammer	25	529	-57	-62	-50	>50	
Skylark	25	644	-47	-54	-40	>25	
Linnet	25	501	-47	-57	-32	>25	
Mistle Thrush	25	500	-46	-52	-38	>25	
Lapwing	25	266	-43	-58	-16	>25	
Meadow Pipit	25	192	-43	-61	-26	>25	
Little Owl	25	66	-37	-55	-8	>25	
Marsh Tit	25	98	-37	-52	-15	>25	
Bullfinch	25	271	-31	-42	-19	>25	

3a. Table of population alerts for CBC/BBS UK 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Tit	10	60	-64	-72	-51	>50	
Turtle Dove	10	199	-58	-62	-51	>50	
Yellow Wagtail	10	164	-42	-53	-31	>25	
Grey Partridge	10	245	-39	-48	-31	>25	
Spotted Flycatcher	10	222	-32	-38	-16	>25	
Corn Bunting	10	144	-27	-38	-15	>25	

3b. Table of population alerts for CBC/BBS England 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Tit	10	53	-62	-71	-51	>50	
Turtle Dove	10	195	-57	-62	-51	>50	
Cuckoo	10	614	-46	-49	-43	>25	
Yellow Wagtail	10	160	-42	-49	-31	>25	
Starling	10	1394	-36	-39	-32	>25	
Grey Partridge	10	220	-33	-41	-25	>25	

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Flycatcher	10	159	-22	-44	-7	>25	
Tree Pipit	10	73	-30	-47	-7	>25	
Willow Warbler	10	938	-30	-36	-24	>25	
Linnet	10	967	-27	-32	-21	>25	

4a. Table of population alerts for CBC/BBS UK 2001-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	5	157	-40	-47	-31	>25	
Willow Tit	5	47	-33	-47	-14	>25	
Little Owl	5	101	-25	-39	-13	>25	
Yellow Wagtail	5	149	-25	-35	-14	>25	

4b. Table of population alerts for CBC/BBS England 2001-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	5	155	-40	-47	-30	>25	
Willow Tit	5	40	-34	-51	-17	>25	

[Back to top](#)

Appendix 7.1 Tables 5a and 5b

7.1 [Tables of alerts and population increases from CBC/BBS](#)

1a.	CBC/BBS UK alerts – 39 years
1b.	CBC/BBS England alerts – 39 years
2a.	CBC/BBS UK alerts – 25 years
2b.	CBC/BBS England alerts – 25 years
3a.	CBC/BBS UK alerts – 10 years
3b.	CBC/BBS England alerts – 10 years
4a.	CBC/BBS UK alerts – 5 years
4b.	CBC/BBS England alerts – 5 years
5a.	CBC/BBS UK population increases of >50% – 39 years
5b.	CBC/BBS England population increases of >50% – 39 years



My BTO

Go to project



VOLUNTEER

JOIN

DONATE

MENU

Page not four

[home](#) » [birdtrends2008](#) » Page not found



Terribly sorry...

The page you are looking for could not be found.

- Try searching using the search bar (top right)
- Navigate to the area you are looking for using the main menu (above)
- [Return to our homepage](#)
- Email us about what you are trying to find, or to tell us about a broken link - info@bto.org

Share this page



Related content



BLOG

Widening BTO's appeal

Andy Clements, BTO's Chief Executive, looks at how BTO can engage new audiences.



ARTICLE

Bird tracking - a masterclass

Phil Atkinson explains the technology behind tracking.



CASE STUDY

Climate change in a warming world

BTO science contributes to our understanding of future scenarios, and informing policies and conservation management strategies to help species adapt.

BTO - Looking out for birds

[About us](#) [Jobs](#) [National offices](#) [Community](#) [Events](#) [Blog](#) [Shop](#)

✉ info@bto.org

[Press releases](#) [Terms & Conditions](#) [Privacy](#) [Cookies](#)



Registered Charity Number 216652 (England & Wales), SC039193 (Scotland)
© British Trust for Ornithology, BTO, The Nunnery, Thetford, Norfolk IP24 2PU, Tel: +44 (0)1842 750050 Fax: +44 (0)1842 750030

Seen a ringed / dead / diseased bird?

REPORT IT TO US ▶

Keep up to date with BTO

E-NEWSLETTER SIGN UP

Flock together



Page not found

home » birdtrends2008 » Page not found



Terribly sorry...


The page you are looking for could not be found.

- Try searching using the search bar (top right)
- Navigate to the area you are looking for using the main menu (above)
- [Return to our homepage](#)
- Email us about what you are trying to find, or to tell us about a broken link - info@bto.org

Share this page



Related content



ARTICLE

Results

See live NBC results and watch the breeding season unfold



EVENT

Northern Ireland Seabird Report Launch

The NI Seabird Report is unique in the UK, presenting the most up-to-date seabird monitoring data collected by volunteers, NGOs and NIEA from around Northern Ireland. Each March we present freshly produced hard copies...



BLOG

Widening BTO's appeal

Andy Clements, BTO's Chief Executive, looks at how BTO can engage new audiences.

BTO - Looking out for birds

[About us](#) [Jobs](#) [National offices](#) [Community](#) [Events](#) [Blog](#) [Shop](#)

✉ info@bto.org

[Press releases](#) [Terms & Conditions](#) [Privacy](#) [Cookies](#)



Registered Charity Number 216652 (England & Wales), SC039193 (Scotland)
© British Trust for Ornithology, BTO, The Nunnery, Thetford, Norfolk IP24 2PU, Tel: +44 (0)1842 750050 Fax: +44 (0)1842 750030

Seen a ringed / dead / diseased bird?

[REPORT IT TO US ▶](#)

Keep up to date with BTO

[E-NEWSLETTER SIGN UP](#)

Flock together



[Back to top](#)

[Appendix 7.2](#)

May we use cookies to monitor performance and improve your experience?

These optional cookies are used for collecting anonymous visit data only.

[READ OUR COOKIE POLICY](#)

ACCEPT ALL COOKIES

[Set preferences](#)

7.2 Tables of alerts and population increases from WBS

1. WBS alerts – 31 years
2. WBS alerts – 25 years
3. WBS alerts – 10 years
4. WBS alerts – 5 years
5. WBS population increases of >50% – 31 years

1. Table of alerts for WBS waterways 1975-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Yellow Wagtail	31	18	-97	-99	-93	>50	Small sample
Reed Bunting	31	51	-61	-75	-33	>50	
Little Grebe	31	16	-59	-87	-9	>50	Small sample
Pied Wagtail	31	64	-53	-66	-43	>50	
Redshank	31	17	-49	-85	-16	>25	Small sample

2. Table of alerts for WBS waterways 1981-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Yellow Wagtail	25	16	-97	-99	-95	>50	Small sample
Little Grebe	25	15	-64	-87	-23	>50	Small sample
Redshank	25	17	-45	-76	-22	>25	Small sample
Pied Wagtail	25	64	-39	-53	-25	>25	
Common Sandpiper	25	26	-28	-44	-10	>25	

3. Table of alerts for WBS waterways 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Sand Martin	10	22	-44	-59	-25	>25	
Redshank	10	13	-33	-51	-12	>25	Small sample

4. Table of population increases for WBS waterways 1975-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Oystercatcher	31	23	113	67	213		
Mallard	31	92	165	88	258		
Whitethroat	31	41	171	15	367		

[Back to top](#)

7.3 Tables of alerts and population increases from CES

1. CES adults alerts – 22 years
2. CES adults alerts – 10 years
3. CES adults alerts – 5 years
4. CES adults population increases of >50% – 22 years

1. Table of alerts for CES adults 1984-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Lesser Redpoll	22	17	-96	-99	-91	>50	Small sample
Linnet	22	20	-92	-97	-84	>50	
Yellowhammer	22	18	-74	-91	-28	>50	Small sample
Willow Warbler	22	88	-61	-70	-52	>50	
Lesser Whitethroat	22	40	-59	-78	-40	>50	
Willow Tit	22	20	-58	-89	-25	>50	
Reed Bunting	22	59	-51	-64	-36	>50	
Whitethroat	22	59	-36	-57	-22	>25	
Sedge Warbler	22	64	-33	-49	-11	>25	
Reed Warbler	22	52	-33	-46	-10	>25	

2. Table of alerts for CES adults 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Lesser Redpoll	10	12	-86	-95	-72	>50	Small sample
Linnet	10	18	-62	-84	-35	>50	Small sample
Lesser Whitethroat	10	37	-49	-61	-32	>25	
Willow Tit	10	15	-49	-82	-4	>25	Small sample
Willow Warbler	10	91	-48	-54	-42	>25	
Sedge Warbler	10	74	-46	-54	-37	>25	
Whitethroat	10	68	-37	-46	-27	>25	
Reed Warbler	10	60	-26	-36	-13	>25	

3. Table of alerts for CES adults 2001-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Sedge Warbler	5	64	-26	-34	-19	>25	

4. Table of population increases for CES adults 1984-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Wren	22	95	58	35	87		
Treecreeper	22	37	60	9	122		
Robin	22	89	63	40	87		
Blackcap	22	86	69	39	112		
Chiffchaff	22	67	161	75	304		

[Back to top](#)

7.4 Tables of population declines and increases from BBS

1. BBS – UK alerts - 11 years
2. BBS – England alerts - 11 years
3. BBS – Scotland alerts - 11 years
4. BBS – Wales alerts - 11 years
5. BBS – UK alerts - 10 years
6. BBS – England alerts - 10 years
7. BBS – Scotland alerts - 10 years
8. BBS – Wales alerts - 10 years
9. BBS – UK alerts - 5 years
10. BBS – England alerts - 5 years
11. BBS – Scotland alerts - 5 years
12. BBS – Wales alerts - 5 years
13. BBS – UK population increases of >50%
14. BBS – England population increases of >50%
15. BBS – Scotland population increases of >50%
16. BBS – Wales population increases of >50%
17. BBS – Northern Ireland population increases of >50%

1. Table of declines >25% for BBS UK 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Tit	11	54	-66	-74	-54	>50	
Turtle Dove	11	182	-60	-66	-54	>50	
Wood Warbler	11	53	-58	-70	-44	>50	
Pied Flycatcher	11	41	-50	-63	-38	>25	
Yellow Wagtail	11	158	-44	-53	-34	>25	
Grey Partridge	11	227	-40	-49	-32	>25	
Curlew	11	453	-38	-43	-33	>25	
Whinchat	11	74	-38	-53	-18	>25	
Spotted Flycatcher	11	202	-37	-46	-23	>25	
Cuckoo	11	729	-33	-38	-27	>25	
Corn Bunting	11	138	-32	-45	-21	>25	
Starling	11	1631	-29	-34	-24	>25	
Swift	11	961	-26	-33	-18	>25	

2. Table of declines >25% for BBS England 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Tit	11	47	-66	-75	-57	>50	
Turtle Dove	11	179	-60	-65	-52	>50	
Cuckoo	11	584	-50	-53	-46	>25	
Yellow Wagtail	11	154	-43	-52	-33	>25	
Starling	11	1335	-38	-41	-34	>25	
Spotted Flycatcher	11	144	-37	-48	-21	>25	
Grey Partridge	11	202	-35	-44	-26	>25	
Tree Pipit	11	67	-35	-51	-11	>25	
Whinchat	11	30	-34	-60	-9	>25	
Lesser Redpoll	11	54	-34	-58	-6	>25	
Willow Warbler	11	866	-32	-37	-27	>25	
Linnet	11	909	-30	-35	-23	>25	
Swift	11	825	-28	-36	-18	>25	
Corn Bunting	11	132	-28	-40	-18	>25	

3. Table of declines >25% for BBS Scotland 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Black-headed Gull	11	72	-69	-83	-45	>50	

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Swift	11	46	-42	-59	18	>25	
Kestrel	11	42	-40	-56	-17	>25	
Lapwing	11	86	-38	-50	-25	>25	
Siskin	11	54	-25	-43	-9	>25	

4. Table of declines >25% for BBS Wales 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Starling	11	83	-46	-64	-23	>25	
Curlew	11	38	-45	-58	-30	>25	
Yellowhammer	11	38	-36	-53	-16	>25	
Cuckoo	11	57	-33	-47	-19	>25	
Garden Warbler	11	53	-29	-46	-5	>25	
Willow Warbler	11	152	-27	-37	-14	>25	
Coal Tit	11	68	-26	-46	-1	>25	
Wheatear	11	50	-25	-43	-8	>25	

1. Table of declines >25% for BBS UK 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Tit	10	53	-64	-72	-51	>50	
Turtle Dove	10	184	-58	-63	-51	>50	
Wood Warbler	10	53	-53	-67	-36	>50	
Pied Flycatcher	10	41	-49	-61	-37	>25	
Yellow Wagtail	10	157	-43	-52	-33	>25	
Grey Partridge	10	230	-38	-47	-30	>25	
Whinchat	10	73	-38	-53	-18	>25	
Curlew	10	460	-36	-40	-31	>25	
Spotted Flycatcher	10	206	-35	-42	-21	>25	
Cuckoo	10	731	-29	-35	-24	>25	
Great Black-backed Gull	10	101	-28	-43	-2	>25	
Starling	10	1665	-28	-32	-23	>25	
Corn Bunting	10	138	-26	-39	-16	>25	

2. Table of declines >25% for BBS England 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Willow Tit	10	46	-63	-72	-54	>50	
Turtle Dove	10	181	-57	-63	-50	>50	
Nightingale	10	30	-47	-61	-31	>25	
Cuckoo	10	582	-46	-49	-42	>25	
Yellow Wagtail	10	153	-42	-51	-32	>25	
Starling	10	1359	-36	-39	-32	>25	
Lesser Redpoll	10	55	-35	-60	-9	>25	
Spotted Flycatcher	10	146	-34	-44	-20	>25	
Whinchat	10	30	-32	-58	-9	>25	
Grey Partridge	10	205	-31	-41	-23	>25	
Willow Warbler	10	873	-31	-36	-26	>25	
Tree Pipit	10	67	-30	-47	-10	>25	
Linnet	10	925	-26	-31	-19	>25	
Swift	10	840	-25	-32	-19	>25	

3. Table of declines >25% for BBS Scotland 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Black-headed Gull	10	72	-63	-79	-39	>50	
Curlew	10	120	-47	-53	-41	>25	
Kestrel	10	42	-34	-50	-12	>25	
Lapwing	10	87	-34	-47	-20	>25	
Great Black-backed Gull	10	37	-33	-49	-8	>25	

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Siskin	10	55	-30	-38	15	>25	

4. Table of declines >25% for BBS Wales 1996-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Curlew	10	39	-43	-55	-29	>25	
Starling	10	85	-41	-61	-20	>25	
Goldcrest	10	83	-38	-56	-7	>25	
Cuckoo	10	59	-32	-46	-21	>25	
Yellowhammer	10	38	-32	-48	-12	>25	
Swift	10	68	-30	-48	-4	>25	
Garden Warbler	10	55	-28	-45	-7	>25	
Whitethroat	10	77	-26	-39	-12	>25	
Willow Warbler	10	157	-26	-36	-14	>25	

1. Table of declines >25% for BBS UK 2001-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	5	157	-40	-46	-32	>25	
Willow Tit	5	47	-36	-51	-21	>25	
Great Black-backed Gull	5	106	-28	-40	-9	>25	

2. Table of declines >25% for BBS England 2001-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Turtle Dove	5	155	-40	-48	-31	>25	
Willow Tit	5	40	-36	-52	-19	>25	
Redshank	5	58	-27	-38	-7	>25	

3. Table of declines >25% for BBS Scotland 2001-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Great Black-backed Gull	5	38	-35	-47	-14	>25	
Curlew	5	110	-31	-40	-24	>25	
Mallard	5	96	-26	-37	-17	>25	
Red Grouse	5	43	-26	-44	-6	>25	

4. Table of declines >25% for BBS Wales 2001-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Goldcrest	5	93	-31	-45	-16	>25	
Curlew	5	37	-26	-40	-10	>25	
Swift	5	77	-26	-41	-11	>25	
Linnet	5	106	-26	-40	-8	>25	

[Back to top](#)

[Appendix 7.4 continued](#)

7.4 Tables of population declines and increases from BBS

1. BBS – UK alerts - 11 years
2. BBS – England alerts - 11 years
3. BBS – Scotland alerts - 11 years
4. BBS – Wales alerts - 11 years
5. BBS – UK alerts - 10 years
6. BBS – England alerts - 10 years
7. BBS – Scotland alerts - 10 years
8. BBS – Wales alerts - 10 years
9. BBS – UK alerts - 5 years
10. BBS – England alerts - 5 years
11. BBS – Scotland alerts - 5 years
12. BBS – Wales alerts - 5 years
13. BBS – UK population increases of >50%
14. BBS – England population increases of >50%
15. BBS – Scotland population increases of >50%
16. BBS – Wales population increases of >50%
17. BBS – Northern Ireland population increases of >50%

5. Table of population increases for BBS UK 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Blackcap	11	1293	55	48	64		
Raven	11	214	80	40	133		
Great Spotted Woodpecker	11	811	112	100	124		
Canada Goose	11	393	115	72	159		
Greylag Goose	11	127	120	21	281		
Stonechat	11	122	200	140	265		
Red Kite	11	43	282	152	599		

6. Table of population increases for BBS England 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Green Woodpecker	11	636	51	42	62		
Buzzard	11	400	90	68	117		
Canada Goose	11	367	102	58	152		
Great Spotted Woodpecker	11	721	105	94	119		
Greylag Goose	11	103	126	64	257		
Stonechat	11	48	167	83	278		
Raven	11	71	172	68	303		
Ring-necked Parakeet	11	34	522	202	1861		

7. Table of population increases for BBS Scotland 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Mistle Thrush	11	66	51	21	101		
Goldcrest	11	79	57	24	82		
Goldfinch	11	73	58	14	126		
Raven	11	39	63	3	138		
Wren	11	194	66	43	80		
House Martin	11	53	84	21	136		
Whitethroat	11	66	87	41	155		
Blackcap	11	36	145	81	246		
Chiffchaff	11	29	158	62	336		
Stonechat	11	29	229	122	359		

8. Table of population increases for BBS Wales 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Blackcap	11	109	56	30	83		
Goldfinch	11	115	56	21	113		
Pheasant	11	82	57	32	97		
House Sparrow	11	113	68	45	100		
Great Spotted Woodpecker	11	61	98	62	148		
Herring Gull	11	76	137	35	294		
Stonechat	11	33	207	117	360		

9. Table of population increases for BBS Northern Ireland 1995-2006

Species	Period (yrs)	Plots (n)	Change (%)	Lower limit	Upper limit	Alert	Comment
Meadow Pipit	11	59	51	21	101		
Jackdaw	11	66	58	11	94		
House Martin	11	36	59	1	146		
Wren	11	82	62	21	95		
Woodpigeon	11	73	64	28	103		
Goldcrest	11	38	84	21	111		
Starling	11	69	86	35	169		
Coal Tit	11	53	87	40	137		
Dunnock	11	62	90	37	136		
Hooded Crow	11	71	111	59	160		
Greenfinch	11	46	115	34	204		
Great Tit	11	61	124	71	160		
Pheasant	11	30	148	46	239		
Goldfinch	11	34	543	.	.		

[Back to top](#)

[Back to Contents](#)



Images: Spotted Flycatcher, by Chris Knights / BTO; Lapwing, by Sarah Kelman / BTO

Breeding Birds in the Wider Countryside: their conservation status 2008

This report is a “one-stop-shop” for information about the population status of our common terrestrial birds. With one page per species, readers can quickly find all the key information about trends in population size and breeding performance as measured by BTO monitoring schemes. It provides an overview of trends for the period 1966-2007.

This report is the third in a series, prepared within the Partnership between the British Trust for Ornithology (BTO) and the Joint Nature Conservation Committee (JNCC) (on behalf of Natural England, Scottish Natural Heritage, Countryside Council for Wales and the Environment & Heritage Service of Northern Ireland) as part of its programme of research into nature conservation.

It is the result of the sustained long-term fieldwork efforts of many thousands of the BTO's volunteer supporters. Without their enthusiasm for collecting these hard-won facts, the cause of conservation in the UK would be very much the poorer.

Baillie, S., Marchant, J., Leech, D., Joys, A., Noble, D., Barimore, C., Grantham, M., Risely, K. & Robinson, R.. 2009. Breeding Birds in the Wider Countryside: their conservation status 2008. *BTO Research Report*, BTO, Thetford, UK.